

Contents

1	Introduction	1
2	Methodological studies on the immunocytochemical detection of isolated residual tumor cells in mesenchymal organs	3
2.1	Bone marrow	3
2.1.1	Specificity of monoclonal antibodies to epithelial marker proteins	3
2.1.2	Sensitivity of tumor cell detection	6
2.1.3	Influence of aspiration site and sample size	8
2.1.4	Influence of downregulated cytokeratin expression	9
2.2	Peripheral blood	11
2.3	Lymph nodes	13
3	Prognostic relevance of individual disseminated tumor cells	14
3.1	Breast cancer	14
3.2	Cancer of the gastrointestinal tract	16
3.2.1	Colorectal cancer	16
3.2.2	Gastric cancer	17
3.2.3	Esophageal cancer	18
3.3	Lung cancer	19
3.3.1	Small cell lung cancer	19
3.3.2	Non-small cell lung cancer	20
3.4	Malignancies of the urogenital tract	23
3.4.1	Prostate cancer	23
3.4.2	Renal cell cancer	25
3.4.3	Urinary bladder cancer	25
4	Immunocytochemical monitoring of therapeutic effects	26
5	Immunocytochemical phenotyping of individual disseminated carcinoma cells	28
5.1	Technical considerations	28
5.2	Detection of histogenetic markers	29
5.3	Expression of proliferation-associated proteins	29
5.4	Expression of oncogenes and tumor suppressor genes	31
5.4.1	The ErbB2 oncogene	31
5.4.2	The p53 tumor suppressor gene	34
5.5	Modulated expression of epithelial cell adhesion molecules	36
5.5.1	Plakoglobin	36
5.5.2	Epithelial glycoprotein 40/17-1A antigen	36
5.6	Modulation of proteins relevant to the immunological anti-tumor defense	38
5.6.1	Major histocompatibility antigens	38
5.6.2	Intercellular adhesion molecule-1 (ICAM-1)	39
6	Development of new diagnostic techniques	40

6.1	Immunobiochemical approach – Cellular enzyme immunoassay	40
6.2	Molecular approaches – Polymerase chain reaction technique	41
6.2.1	Detection of histogenetic marker expression	41
6.2.2	Detection of tumor-associated gene expression	42
6.2.3	Detection of genomic changes	43
7	Concluding remarks	43
	References	50
	Subject index	61