

## Contents

<b>Preface</b>	v
<b>Quantitation in PET and image processing</b>	1
<i>Invited lecture</i>	
Role of human and animal PET studies in drug development <i>A.A. Lammertsma</i>	3
Kinetic analysis of neuroreceptor binding using PET <i>V.J. Cunningham, E.A. Rabiner, J.C. Matthews, R.N. Gunn, S. Zamuner and A.D. Gee</i>	12
<i>Session</i>	
Formation of parametric images with statistical clustering <i>Y. Kimura</i>	25
Development of motion correction technique for PET study using optical tracking system <i>H. Watabe, K. Koshino, P.M. Bloomfield, R.R. Fulton and H. Iida</i>	31
3D deformable image processing and integration <i>H.-D. Lin and K.-P. Lin</i>	39
Impact of inter-subject image registration on group analysis of fMRI data <i>B.A. Ardekani, A.H. Bachman, S.C. Strother, Y. Fujibayashi and Y. Yonekura</i>	49
Rapid protocol for quantitative CMRO <sub>2</sub> and CBF using PET and O-15 labelled compounds <i>K. Nobuyuki, H. Takuya, T. Noboru, W. Hiroshi, K. Naoki, O. Youichirou, K. KyonMim and I. Hidehiro</i>	60
<i>Poster</i>	
Performance evaluation of microPET P4 for rat, rabbit and monkey <i>Y. Wada, A. Matsumura, F. Nakamura, M. Tanaka, H. Mizuma, K. Mizuno, S. Nozaki, H. Nakajima, S. Kondo, K. Ishii, K. Koyama, Y. Hamazawa, T. Okuma, T. Okamura, Y. Inoue and Y. Watanabe</i>	69

Simultaneous quantification of two brain functions with dual tracer injection in PET dynamic study <i>Y. Ikoma, H. Toyama and T. Suhara</i>	74
Improved parametric images of blood flow and vascular volume by cluster analysis in H <sub>2</sub> <sup>15</sup> O brain PET study <i>K.M. Kim, H. Watabe, T. Hayashi, N. Kudomi and H. Iida</i>	79
Impact of image-based scatter correction for <sup>123</sup> IMP-SPECT and SPM analysis <i>M. Shidahara, H. Watabe, K. Kim, S. Kawatsu, R. Kato, T. Kato, H. Iida and K. Ito</i>	84
Development of FDG-PET quantitative procedures without blood sampling <i>L.-C. Wu, Y.-H. Fang, T. Kao and R.-S. Liu</i>	89
New method for the synthesis of <sup>15</sup> O-labeled carbon monoxide and <sup>15</sup> O-labeled dioxide for rapid supply in clinical use <i>Y. Miyake, H. Iida, K. Hayashida and Y. Ishida</i>	93
Neutron reduction of the small cyclotron for production of oxygen-15-labeled gases <i>T. Inomata, M. Fujiwara, H. Iida, N. Kudomi and I. Miura</i>	97
Image improvement in pinhole SPECT using complete data acquisition combined with statistical image reconstruction <i>T. Zeniya, H. Watabe, T. Aoi, K.M. Kim, N. Teramoto, T. Hayashi, A. Sohlberg, H. Kudo and H. Iida</i>	101
Effects of motion correction on quantification of myocardial blood flow with <sup>15</sup> O-H <sub>2</sub> O PET <i>K. Koshino, N. Kawachi, T. Hayashi, H. Watabe, S. Hasegawa, J. Hatazawa and H. Iida</i>	106
<b>Quantitation and hyperpolarized contrast agent in MRI</b>	111
<b><i>Invited lecture</i></b>	
Quantitative T <sub>1</sub> mapping and absolute water content measurement using MRI <i>H. Neeb, T. Dierkes and N.J. Shah</i>	113
<b><i>Session</i></b>	
Hyperpolarized <sup>129</sup> Xe as a novel probe agent of lung functions in MRI and MRS. Experimental results with mice at 9.4 T <i>H. Fujiwara, A. Kimura and T. Wakayama</i>	124

Dynamic spectroscopy of hyperpolarized Xe-129 in rat lung <i>H. Sato, J.-i. Enmi, T. Hayashi, N. Takei, Y. Iwadate, S. Abe, N. Teramoto, N. Kawachi, M. Hattori, H. Watabe, T. Sawada, K. Uchiyama, T. Tsukamoto, K. Nagasawa and H. Iida</i>	131
In vivo MR spectroscopy of hyperpolarized Xe-129 in rat brain <i>A. Wakai, K. Nakamura, J. Kershaw and I. Kanno</i>	139
Developments of apparatus for hyperpolarization of Xe-129: optimized structure of cell for efficient spin exchange at a high concentration of optically pumped rubidium <i>M. Hattori, T. Hiraga, M. Saito, T. Nakai and T. Moriya</i>	144
Measurement of cerebral blood flow with dynamic susceptibility contrast MRI and comparison with O-15 positron emission tomography <i>J.-i. Enmi, T. Hayashi, H. Watabe, H. Moriwaki, N. Yamada and H. Iida</i>	150
Measurement of cerebral circulation by dynamic susceptibility contrast-enhanced MRI: effect of tracer delay <i>M. Ibaraki, E. Shimosegawa, H. Toyoshima, K. Ishigame, S. Miura, K. Takahashi and I. Kanno</i>	159
<b>Poster</b>	
MRI and MRS of mouse head at 9.4 T using hyperpolarized Xenon gas <i>T. Wakayama, T. Oose, M. Narazaki, A. Kimura and H. Fujiwara</i>	167
Practical application of NMR spectra of $^{129}\text{Xe}$ dissolved in red blood cell <i>Y. Kawata, T. Kamiya, H. Miura, A. Kimura and H. Fujiwara</i>	172
MRS of hyperpolarized $^{129}\text{B Xe}$ in the chest of mouse—effect of contrast agents on the MRS <i>M. Narazaki, T. Wakayama, A. Kimura and H. Fujiwara</i>	177
Distortion correction in echo-planar imaging and quantitative $T_2^*$ mapping <i>T. Dierkes, H. Neeb and N.J. Shah</i>	181
<b>Application of quantitative PET and MRI techniques</b>	187
<b>Invited lecture</b>	
Quantitative PET for assessment of cerebral blood flow and glucose consumption under varying physiological conditions <i>G.M. Knudsen, E. Rostrup and S.G. Hasselbalch</i>	189
Cortical layer-dependent CBF changes induced by neural activity <i>S.-G. Kim and S.-P. Lee</i>	201

**Session**

Changes in CBF and CBV during changes in neural activity or $P_a\text{CO}_2$ measured by PET <i>H. Ito, I. Kanno, H. Fukuda and S. Miura</i>	211
Measurement of arterial part of vascular volume ( $V_0$ ) for the evaluation of hemodynamic changes in cerebrovascular disease <i>H. Okazawa, H. Yamauchi and Y. Yonekura</i>	218
A physiological model for cerebral oxygen delivery and consumption and effective oxygen diffusibility evaluated by PET <i>T. Hayashi, N. Kudomi, N. Teramoto, H. Watabe, J.-I. Enmi, K.-M. Kim and H. Iida</i>	228
Cerebral perfusion measurements using continuous arterial spin labeling: accuracy and limits of a quantitative approach <i>H. Kimura, H. Kabasawa, Y. Yonekura and H. Itoh</i>	238
Positron emission tomography—molecular imaging of biological processes <i>J. Knuuti</i>	248
<b>Poster</b>	
[ $^{18}\text{F}$ ]FDG mouse brain PET imaging: absolute quantification of regional cerebral glucose utilization as compared with 2-[ $^{14}\text{C}$ ]DG autoradiography <i>H. Toyama, M. Ichise, J.-S. Liow, K.J. Modell, D.C. Vines, S. Zoghbi, T. Esaki, M. Cook, J. Seidel, K. Katada, L. Sokoloff, M.V. Green and R.B. Innis</i>	255
Development of injectable O-15 oxygen and its application for estimation of OEF <i>T. Temma, Y. Magata, H. Iida, T. Hayashi, M. Ogawa, T. Mukai, Y. Iida, H. Tsukada, J. Konishi and H. Saji</i>	262
Imaging of vulnerable atherosclerotic plaque with [ $^{18}\text{F}$ ]FDG-PET: an animal atherosclerosis model study <i>M. Ogawa, T. Mukai, S. Ishino, N. Teramoto, H. Watabe, N. Kudomi, M. Shiomi, Y. Magata, H. Iida and H. Saji</i>	266
Development of modified method for quantification of regional myocardial blood flow using $^{15}\text{O}$ -water with PET <i>C. Katoh, K. Morita, K. Nakada and N. Tamaki</i>	269
Assessment of the precision in co-registration of structural MR images and PET images with localized binding <i>P. Willendrup, L.H. Pinborg, S.G. Hasselbalch, K.H. Adams, K. Stahr, G.M. Knudsen and C. Svarer</i>	275
<b>Author index</b>	281
<b>Keyword index</b>	285