

|  |                |
|--|----------------|
| List of Figures  | <i>page xi</i> |
| List of Tables   | xiii           |
| Preface  | xv             |
| Acknowledgments  | xix            |
| <b>1 The Persistence of Misery in Europe and America before 1900</b> | 1              |
| <b>2 Why the Twentieth Century Was So Remarkable</b>                 | 20             |
| <b>3 Tragedies and Miracles in the Third World</b>                   | 43             |
| <b>4 Prospects for the Twenty-First Century</b>                      | 66             |
| <b>5 Problems of Equity in Health Care</b>                           | 96             |
| <b>Postscript: How Long Can We Live?</b>                             | 108            |
| Appendix   | 113            |
| Notes  | 127            |
|  | ix             |

|                                    |     |
|------------------------------------|-----|
| <b>Glossary of Technical Terms</b> | 145 |
| <b>Biographical Notes</b>          | 151 |
| <b>References</b>                  | 155 |
| <b>Index</b>                       | 183 |

---

## Figures

|  |                  |
|--|------------------|
| <b>Frontispiece</b>  | <i>page viii</i> |
| <b>1.1 Secular Trends in Mortality Rates in England and France</b>   | 7                |
| <b>1.2 Trend in Mean Final Height of Native-Born White American Males and Trend in Their Life Expectancy at Age 10</b>   | 17               |
| <b>2.1 The Growth of World Population and Some Major Events in the History of Technology</b>   | 22               |
| <b>2.2 Relative Mortality Risk among Union Army Veterans and among Modern Norwegian Males</b>  | 23               |
| <b>2.3 Comparison of Relative Mortality Risk by BMI among Men 50 Years of Age, Union Army Veterans around 1900 and Modern Norwegians</b>                                     | 24               |
| <b>2.4 Iso-Mortality Curves of Relative Risk for Height and Weight among Norwegian Males Aged 50–64, with a Plot of the Estimated French Height and Weight at Four Dates</b> | 26               |

|   |    |
|---|----|
| 2.5 Relationship between Height and Relative Risk of Ill Health in NHIS Veterans Aged 40–59   | 28 |
| 2.6 Mean BMI by Age Group and Year, 1864–1991   | 29 |
| 2.7 Health Improvement Predicted by NHIS 1985–88 Health Surface   | 30 |
| 3.1 Secular Trends in the Average Heights of Male Adolescents in Great Britain, 1748–1993, Relative to Current Dutch Growth Curves  | 48 |
| 3.2 Waaler Surface of Relative Mortality Risk for Height and Weight among Norwegian Males Aged 50–64 with a Plot of the Estimated French and English Heights and Weights since 1705 at Ages 25–39 | 56 |
| 3.3 Perinatal Death Rate by Birth Weight in Ghana, India, and the United States   | 58 |
| 3.4 Efficient Region of Body Build for Health Production on a Waaler Surface in Mortality for Norwegian Males Aged 45–89  | 62 |
| 3.5 Mean Height and Weight of 140 Adult Male Populations in 1990  | 63 |
| 3.6 Iso-Mortality Curves of Relative Risk for Height and Weight among Norwegian Males Aged 50–64, with Two Plots  | 64 |
| 4.1 Relative Burden of Health Care by Age, U.S. Data c. 1996  | 85 |
| 4.2 How Will the Curve of Relative Disease Burden Shift?  | 86 |
| 4.3 Index of Average Annual Health Care Costs by Year before Death  | 87 |

---

## Tables

|   |               |
|---|---------------|
| 1.1 Life Expectancy at Birth in Seven Nations, 1725–2100  | <i>page 2</i> |
| 1.2 Secular Trends in the Daily Caloric Supply in France and Great Britain, 1700–1989   | 9             |
| 1.3 A Comparison of Energy Available for Work Daily per Consuming Unit in France, England and Wales, and the United States, 1700–1994   | 11            |
| 1.4 Estimated Average Final Heights (cm) of Men Who Reached Maturity between 1750 and 1975 in Six European Populations, by Quarter Centuries  | 13            |
| 1.5 A Comparison of the Average Daily Uses of Dietary Energy in England and Wales in 1700 and 1800  | 14            |
| 2.1 Comparison of the Prevalence of Chronic Conditions among Union Army Veterans in 1910, Veterans in 1983 (Reporting Whether They Ever Had Specific Chronic Conditions), and Veterans in NHIS, 1985–88 (Reporting Whether They Had Specific Chronic Conditions during the Preceding 12 Months), Aged 65 and Above, Percentages | 31            |

|     |   |     |
|-----|---|-----|
| 4.1 | Secular Trends in Time Use: The Average Hourly Division of the Day of the Average Male Household Head   | 68  |
| 4.2 | Estimated Trend in the Lifetime Distribution of Discretionary Time  | 71  |
| 4.3 | Annual Rate of Decline in Prevalence Rates of Selected Chronic Conditions among Elderly Veterans between 1910 and the Mid-1980s (in Percent) before and after Alleviating Interventions | 81  |
| 4.4 | The Long-Term Trend in the Structure of Consumption and the Implied Income Elasticities of Several Consumption Categories   | 89  |
| 4.5 | Average Number of Chronic Conditions per U.S. Male in 1900 and in the 1990s   | 91  |
| 4.6 | Average Capacity of Males to Engage in Manual Labor, by Age, c. 1900  | 92  |
| A1  | Relative Risks of Mortality   | 114 |
| A2  | Relative Mortality Risk Table for Norwegian Males Aged 50–64, by Weight (kg) and Height (m)   | 116 |
| A3  | Relative Mortality Risk Table for Norwegian Males Aged 50–64, by BMI and Height, Also Showing the Optimal BMI and Minimum Risk at Each Height   | 123 |