

CONTENTS

ART.	PAGE
INTRODUCTION	1
I. THE RESULTANT	
2. Resultant of two homogeneous polynomials	4
6. Resultant of n homogeneous polynomials	7
8. Resultant isobaric and of weight L	11
8. Coefficient of $a_r^{Lr} \dots a_n^{Ln}$ in R is $R_r^{lr^{l_{r+1}} \dots l_n}$	11
8. The extraneous factor A involves the coefficients of $(F_1, F_2, \dots, F_{n-1})_{x_n=0}$ only	11
9. Resultant is irreducible and invariant	12
10. The vanishing of the resultant is the necessary and sufficient condition that $F_1 = \dots = F_n = 0$ should have a proper solution	13
11. The product theorem for the resultant	15
11. If (F_1, \dots, F_n) contains (F_1', \dots, F_n') , R is divisible by R'	15
12. Solution of equations by means of the resultant	15
12. The u -resultant resolves into linear homogeneous factors in x, u_1, u_2, \dots, u_r	16
II. THE RESOLVENT	
15. Complete resolvent is a member of the module	20
15. Complete resolvent is 1 if there is no finite solution	21
17. Examples on the resolvent	21
18. The complete u -resolvent F_u	24
18. $(F_u)_{x=u_1x_1+\dots+u_nx_n=0} \text{ mod } (F_1, F_2, \dots, F_k)$	24
19. All the solutions of $F_1 = F_2 = \dots = F_k = 0$ are obtainable from true linear factors of F_u	25
20. Any irreducible factor of F_u having a true linear factor is a homogeneous whole function of x, u_1, \dots, u_n	26
21. Irreducible spreads of a module	27
22. Geometrical property of an irreducible spread	28

III. GENERAL PROPERTIES OF MODULES

ART.		PAGE
23.	$M/M' = M/(M, M')$	30
23.	If $M'M''$ contains M , M' contains M/M''	31
24.	Associative, commutative, and distributive laws	31
25.	$(M, M')[M, M']$ contains MM'	32
26.	M/M' and $M/(M/M')$ mutually residual with respect to M	32
28.	$M/(M_1, M_2, \dots, M_k) = [M/M_1, M/M_2, \dots, M/M_k]$	33
28.	$[M_1, M_2, \dots, M_k]/M = [M_1/M, M_2/M, \dots, M_k/M]$	33
30.	Spread of prime or primary module is irreducible	34
31.	Prime module is determined by its spread	34
32.	If M is primary some finite power of the corresponding prime module contains M	35
33.	A simple module is primary	36
34.	There is no higher limit to the number of members that may be required for the basis of a prime module	36
34.	Space cubic curve has a basis consisting of two members	37
35.	The L.C.M. of primary modules with the same spread is a primary module with the same spread	37
36.	If M is primary M/M' is primary	37
37.	Hilbert's theorem	38
38.	Relations between a module and its equivalent H -module	39
38, 42.	Condition that an H -module M may be equivalent to $M_{x_n=1}$	39
38.	Properties of an H -basis	40
39.	Lasker's theorem	40
40.	Method of resolving a module	42
41, 44.	Conditions that a module may be unmixed	43
42.	Deductions from Lasker's theorem	44
42.	When M/M' is M and when not	44
42.	No module has a relevant spread at infinity	44
43.	Properties of the modules $M^{(r)}$, $M^{(s)}$	45
44.	Section of prime module by a plane may be mixed	47
46.	The Hilbert-Netto theorem	48
UNMIXED MODULES		49
48.	Module of the principal class is unmixed	49
49.	Conditions that (F_1, F_2, \dots, F_r) may be an H -basis	50
50.	Any power of module of principal class is unmixed	51
51, 52.	Module with γ -point at every point of M	52

ART.	PAGE
52. When a power of a prime module is unmixed	53
53. Module whose basis is a principal matrix is unmixed	54
SOLUTION OF HOMOGENEOUS LINEAR EQUATIONS	58
NOETHER'S THEOREM	60
56. The Lasker-Noether theorem	61
IV. THE INVERSE SYSTEM	
58. Number of modular equations of an H -module of the principal class	65
59. Any inverse function for degree t can be continued	67
59. Diagram of dialytic and inverse arrays	67
59. The modular equation $1=0$	69
60, 82. The inverse system has a finite basis	69
61. The system inverse to (F_1, F_2, \dots, F_k) is that whose F_i -derivates vanish identically	70
62. Modular equations of a residual module	70
63. Conditions that a system of negative power series may be the inverse system of a module	71
64. Corresponding transformations of module and inverse system	71
65. Noetherian equations of a module	73
65. Every Noetherian equation has the derivate $1=0$	73
65. The Noetherian array	75
66. Modular equations of simple modules	75
PROPERTIES OF SIMPLE MODULES	77
67. A theorem concerning multiplicity	77
69. Unique form of a Noetherian equation	79
71. A simple module of the principal Noetherian class is a principal system	80
72. A module of the principal class of rank n is a principal system	81
73. $\mu = \mu' + \mu''$	82
74. $\mu'_v + \mu''_{v'} = \mu_v = \mu_{v'}$, where $v + v' = \gamma - 1$	83
75. $H_m = 1 + \mu_1 + \mu_2 + \dots + \mu_m$	83
76. $H'_v - H''_{v'} = H'_{v+v'} - H_{v'} = H_v - H''_{v+v'}$, where $v + v' = \gamma - 2$.	84
MODULAR EQUATIONS OF UNMIXED MODULES	85
77. Dialytic array of $M^{(r)}$	86
78. Solution of the dialytic equations of $M^{(r)}$	88

ART.		PAGE
79.	Unique system of r -dimensional modular equations of M	89
79.	The n -dimensional equations	89
80.	Equations of the simple H -module determined by the highest terms of the members of an H -basis of $M^{(r)}$	89
81.	If $R=1$ and M is unmixed, M is perfect	90
82.	If $M^{(r)}$ is a principal system so is M	90
82.	A module of the principal class is a principal system	90
83.	$M^{(r)}$ and M are principal systems if the module determined by the terms of highest degree in the members of an H -basis of $M^{(r)}$ is a principal system ; not conversely	91
84.	Modular equations of an H -module of the principal class	92
85.	Whole basis of system inverse to $M^{(r)}$	93
86.	Modules mutually residual with respect to an H -module of the principal class	94
87.	The theorem of residuation	96
88.	Any module of rank n is perfect	98
88.	An unmixed H -module of rank $n-1$ is perfect	98
88.	An H -module of the principal class is perfect	98
88.	A module of the principal class which is not an H -module is not necessarily perfect	98
88.	A prime module is not necessarily perfect	98
89.	An H -module M of rank r is perfect if the module $M_{x_{r+2}=\dots=x_n=0}$ is unmixed	99
90.	A perfect module is unmixed	99
90.	The L. C. M. of a perfect module of rank r and any module in x_{r+1}, \dots, x_n only is the same as their product	99
91.	Value of H_l for a perfect module	99
92.	If M, M' are perfect H -modules of rank r , and if M contains M' , and $M_{x_{r+1}=\dots=x_n=0}$ is a principal system, M/M' is perfect	100
	NOTE ON THE THEORY OF IDEALS	101