TABLE OF CONTENTS

| CHAPTER | PAGE |
|--|------------------------|
| I. HISTORY OF SCHLICHT FUNCTIONS AND ELEMENTARY PROPERTIES OF THE nTI | Ŧ |
| REGION | . 1 |
| 1.1. Brief history of the theory of schlicht functions | . 1 |
| 1.2. Definition of the nth coefficient region. Description of the 3rd region | . 5 |
| 1.3. Definition of a general class of extremal problems for schlicht function | s 8 |
| 1.4. Elementary properties of the <i>n</i> th coefficient region. Characterization of interior points. Topological equivalence of the <i>n</i> th region and the closed $(2n-2)$ -dimensional full sphere | e |
| II. Variations of schlicht functions | |
| 2.1. Formation of Riemann surfaces of genus zero by the identification o arcs on the circumference of the unit circle. Proof of a particula case of the uniformization principle | r . 14 |
| 2.2. The domain obtained from the plane by identifying the edges of two Jordan arcs with common end-points, one Jordan arc lying in an eneighborhood of the other. Construction of the first variation of the mapping function. Construction of the variation when the domain is obtained from the unit circle instead of the plane. Variation of schlicht function of class § | e s s |
| 2.3. Application of the variational formula to obtain a differential equation satisfied by a function belonging to a boundary point of V_n where a certain function of the first n coefficients is maximal | a. . 36 |
| 2.4. Differential equation satisfied by a function belonging to any boundary point of V_n | . 40 |
| 2.5. The nth coefficient region for functions which are schlicht and bounded by unity | . 41 |
| 2.6. Interior variations | . 43 |
| III. THE CRITICAL POINTS OF THE DIFFERENTIAL EQUATION | |
| 3.1. Hyperelliptic trajectories, or loci in the plane on which the real part of a hyperelliptic integral is constant | |
| 3.2. Local behavior of the loci near zeros and simple poles of the integrand lying in the finite part of the plane. Behavior near a double pole Behavior near a pole of order higher than the second | |
| 3.3. Local behavior of the loci near the point at infinity | . 55 |
| IV. THE Γ-STRUCTURE. BEHAVIOR IN THE LARGE | . 56 |
| 4.1. Definition and classification of the critical points of the loci. Proof tha a domain bounded by finitely many arcs on which the real part of th hyperelliptic integral is constant and by one arc on which the imagi nary part is constant must contain a critical point of negative order | e - . 5 6 |
| 4.2. Investigation of the behavior of the loci defined by $P(w)$, $Q(z)$, and $\Phi(v)$ when prolonged from a point of the plane | |
| 4.3. Definition of the Γ -structures Γ_w , Γ_z , and Γ_v . The connectivity of Γ and Γ_z | |
| 4.4. Dissection of the plane by Γ_w and Γ_s into end domains and strip domains The structure of Γ_w and Γ_s in the large | |
| 4.5. Dissection of the plane by Γ_{\bullet} . The structure of Γ_{\bullet} in the large | |
| V. Geodesics. Continuity theorem | |
| 5.1. The metrics defined by $P(w)$, $Q(z)$, and $\Phi(v)$. Admissible paths | |

| | | | GE |
|-----------|-------------|--|-------|
| CHAPTER | 5.2. | Proof that any two points of the plane different from the origin are | 76 |
| | •• | | |
| | 5.3. | | 80 |
| | •••• | | 00 |
| | 5.4. | | 81 |
| | | | 01 |
| VIE | UNCTIO | " AND SATISFI THE DIFFERENCE | 88 |
| V 2. 1 | EOH | | 88 |
| | 6.1. | | 88 |
| | 6.2. | | 93 |
| VII T | HE LEI | Prototistit I PP S METHOD. | 93 |
| V 11. 1 | 7.1. | | 93 |
| | 7.2. | | 99 |
| | 7.3. | | 109 |
| | | | 100 |
| WIII R | TEL ATTI | | |
| V 111. IV | 8.1. | ONS BETWEEN $P(w)$ AND $Q(z)$. The coefficients of a \mathfrak{D}_n -equation. | 100 |
| | 8.2. | | |
| | 0.2. | | 110 |
| | 8.3. | | |
| | 8.4. | The same is a file of corresponding to any given in (w) | 111 |
| | 8.5. | The standard Riemann surface of genus zero by ruonomy | |
| | 0.0. | | |
| | | the most six defined by $H(z)$. The local uniformizers of the | 110 |
| | | | 112 |
| | 8.6. | The first of Francisco the closed Riemann surface onto | |
| | 0.0. | | 117 |
| | 8.7. | $=$ a (1) $c_{\text{total}} = c_{\text{total}} = c_{$ | |
| | 0.1. | 11 ming function H at noints of 21 - 1 | 118 |
| | 8.8. | - a tit - li-timentiahod evelo 2: Of a canonical identification | |
| | 0.0. | . I I amd of the chara N. Pront that to cath point of | |
| | | g 1 - lalames o unique (1) -timetion | |
| | 8.9 | or the identification of the identification | • |
| | 0.9 | g view which identifies a pair of abiliting ares on the unit off | |
| | | Continuous realization of the identification. A differ | |
| | | u(z, u). The limit of $v(z, u)$ | |
| | | 1 1-4- mana | |
| 7.57 | T V | | |
| IX. | | my T " dobbod by the continuous realization of the recommendation | |
| | 9.1 | c time and the Löwner differential equation for $V(2, \omega) \cdots \cdots$ | |
| | | The Tamper representation for the coefficients of the function $V(z)$, w | • |
| | 9.2 | m_{1} differential equation for $O(z, u)$. Definition of the coefficient | |
| | | a () 1 (1. differential equations for the (1)(1), the system of | |
| | | u x z | . 202 |
| | | b Toll - Tarmor curves lying on the houndary of Vn . Denimon of the | • |
| | 9.3 | f(z, u) The limit of $f(z, u)$ as u tends to zero. The differential | P.7 |
| | | $h(x, y) = h_x + h_y + $ | . 100 |
| | _ | THE PROPERTY OF STREET | . 110 |
| Χ. | | | |
| | 10. | | of |
| | 10. | 2. Proof that a function belonging to a point of 22 maps the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the plane minus a set of arcs without finite critical the unit circle onto the unit circle of the unit circle onto the u | al |
| | | points | . 144 |
| | | points | |

| HAPTER 10 | .3. Proof that the supporting surface K_n is connected and contains the | AGE |
|--------------|---|------------|
| | curve (2e^{iθ}, 3e^{2iθ}, ··· , ne^{(n-1)iθ}). Proof that a function belonging to a point of K_n maps the interior of the unit circle onto the plane minus a single analytic slit. Proof that the convex hull of the Koebe curve (2e^{iθ}, 3e^{2iθ}, ··· , ne^{(n-1)iθ}) does not contain the set V_n in case n > 2. 4. Functions f(z) satisfying more than one D_n-equation. Example of an | 147 |
| | algebraic function $f(z)$ satisfying two \mathfrak{D}_n -equations which maps $ z < 1$ onto the complement of a single analytic arc without finite critical points. Examples of functions satisfying several \mathfrak{D}_n -equations which map $ z < 1$ onto the plane slit along straight lines. Proof that if $f(z)$ satisfies a \mathfrak{D}_n -equation where $n-1$ is a prime, then it satisfies no | 150 |
| | other \mathfrak{D}_n -equation unless it is a rational function of simple type | 153 |
| | PORTION OF THE BOUNDARY OF V_n CORRESPONDING TO SINGLE ANALYTIC | 150 |
| | SLITS | 199 |
| | .1. Definition of the boundary set L_n . Differential equations of the Löwner curves belonging to L_n | 159 |
| 11 | .2. Proof by the method of successive approximation that there are curves lying on the boundary of V_n which begin at points of the Koebe curve and sweep over all of L_n | 160 |
| - 11 | and sweep over all of L_n . 3. Proof that points on the boundary of V_n defined by functions $Q(z)$ | 100 |
| 11 | having only one zero on $ z = 1$ which is of precise order 2 depend analytically on the roots of $Q(z)$ | 165 |
| 11 | .4. Definition of the boundary set M_n . Proof that M_n is not empty, that it is open, and that its closure contains the supporting surface K_n . The one-one correspondence between points of M_n and a class of | |
| | Q-functions | 172 |
| •• | | 112 |
| | .5. Proof of the variational formula for all functions belonging to a neighborhood of a point of M_n | 182 192 |
| | AMETRIZATION OF THE BOUNDARY OF γ_n . 1. Definition of the tree or linear graph defined by a point of the space S_n . | 192 |
| | 2. The b-sequence of a tree. Formula for the number of topologically dif- | |
| | ferent trees having the same number of terminal tips | 194 |
| 12 | sequence of a tree | 199 |
| 12 | 4. The λ-sequence | 201 |
| 12 | .5. The parameters which define a tree | |
| 12 | | 206 |
| 12 | 1.7. The dissection of the space S_n into portions whose points depend on a | 000 |
| | set of real parameters | 206 |
| | REGION V ₃ | 208 |
| 13 | Summary of results to be proved including parametric formulas for the boundary of V ₃ . Discussion of the tables given in the Appendix | 208 |
| 13 | 3.2. Integration of the differential equation in special cases. Integration of the right and left sides of the differential equation in the general case | 211 |
| 4.6 | | 215 |
| | | 210 |
| 18 | to functions which map the interior of the unit circle onto the plane minus a forked slit | 218 |
| 10 | 3.5. The parametric formulas for the portion of the boundary corresponding | |
| 18 | to functions which map the interior of the unit circle onto the plane minus an unforked analytic slit | 223 |
| | minus an umorked analytic site | |

| HAPTER | | AGE |
|--------------|---|-------------|
| XIV. А метно | od for investigating the conjecture $ a_4 \leq 4$ | 226 |
| 14.1. | Brief outline of the method | 226 |
| 14.2. | Formulas expressing a_2 , a_3 , a_4 in terms of the coefficients of the \mathfrak{D}_4 - | |
| | equation. Special case of these formulas when Re (a_4) is maximal | 227 |
| 14.3. | The Prayitz form of the area-principle. Proof that $a_4^{(2)} = a_2^2 + 2a_3$ is | |
| | not zero when Re $(a_*) \ge 4$ | 22 8 |
| 14.4. | Dissection of the Löwner formula for a_4 into terms J_2 , J_4 , J_6 , and | |
| 11.1. | I_1 , I_3 , I_5 | 229 |
| 14.5. | Determination of a lower bound for J_2 in terms of an integral over a | |
| 14.0. | triangular region of the u, v plane. Proof that for points in the tri- | |
| | angular region the integrand is a positive semi-definite quadratic | |
| | form in the values $h(u)$, $h(v)$ | 231 |
| 44.0 | Determination of a lower bound for the integral of the quadratic form | |
| 14.6. | in terms of the integral of the square of the function $h(u)$. Proof of | |
| | the existence of the minimizing function. The Sturm-Liouville sys- | |
| | tem for the determination of the minimizing function. | 234 |
| | Calculation of the smallest eigen-value of the system | 241 |
| 14.7. | Calculation of the smallest eigen-value of the system | 245 |
| 14.8. | Determination of an upper bound for J_4 | 210 |
| 14.9. | Proof that if the function $\alpha(u)$ differs from a constant by at most | |
| | 0.08697 for $0 \le u \le 1$, then the coefficient a_4 given by Löwner's for- | 947 |
| | mula does not exceed 4 in modulus | 24 i |
| 14.10. | Parametric formulas for a_2 , a_3 obtained from the Q-function corre- | 050 |
| | sponding to maximal Re (a ₄) | 250 |
| 14.11. | Elimination of one of the real parameters for a large portion of the | 051 |
| | parameter domain | 251 |
| 14.12. | Integration of a system of differential equations over the remaining | |
| | portion of the parameter domain | 260 |
| XV. THE RE | GION OF VALUES OF THE DERIVATIVE OF A SCHLICHT FUNCTION (by Arthur | |
| Gra | ad) | 263 |
| 15.1. | Definition of the region of values in the plane of $f'(z_0)$ (the region | l |
| | $R(z_0)$). The corresponding region of values for any simply-connected | |
| | domain. The choice of a real $z_0 = r$. The symmetry of $R(r)$. The | ; |
| | dependence of $R(r)$ on r. Interior points of $R(r)$. Definition of the | , |
| | region $L(r)$ in the logarithmic plane | 263 |
| 15.2. | The formulas for the boundary of $L(r)$ | 265 |
| 15.3. | The differential equation satisfied by boundary functions. Investiga- | - |
| | tion of the equation connecting φ, ψ, ρ , and r | 268 |
| 15.4. | The Γ_* -structure | 274 |
| 15.5. | Determination of the portion of the boundary corresponding to un- | - |
| | forked functions | 276 |
| 15.6. | | l |
| 20.0. | functions | 285 |
| 15.7. | | Э |
| 10.1. | boundary curves intersect is equal to π | 2 90 |
| APPENDIX | boundary our top intorpoor to oquar to attended to attended to attended to attended to attended to attend | |
| Table I | Points (a_2, a_3) for which a_2 is real | 292 |
| Table I. I | Points (a_2, a_3) for which a_3 is real. | 2 98 |
| Lable 11. | Y | . 304 |
| DIBLIOGRAPH | Y | . 307 |
| INDEX | | |