

Contents

Preface	xiii
Acknowledgements	xvii

Part A

1 Introduction to Part A	1
2 The example	5
3 Styles of analysis	11
3.1 Preliminary considerations	11
3.2 Scatter Plot Matrices	13
3.3 Within environment analyses for single attributes	17
3.4 Overlap plots	22
3.5 Profiles across environments and B-confidence	27
3.6 Grouping the data	30
3.7 Smoothing the profiles	35
3.8 Profiles across attributes	39
3.9 Choice of expression	39
3.10 This account in a larger framework	43

Part B

4 Introduction to Part B	47
5 Overall behaviour: SPLOMs across attributes and across environments	51

6 Semigraphical comparison of genotypes for each attribute-environment pair	69
7 Profiles of individual genotypes and groups	101
8 SPLOMs for genotype groups	129
9 Re-attribution of the responses on the 43 genotypes	137
9.1 Our specific example	148
9.2 Styles of re-attribution	152
10 Semigraphical displays for the re-attributed data	155
11 Re-environmenting	169
11.1 Two remarks	181
11.2 Overview of position	181
12 Re-expression	185

Part C

13 Introduction to Part C	211
14 Global aspects of the data	213
14.1 Ordering of genotypes	213
14.2 Linear ordering of genotypes	214
14.3 Dispersed inheritance, etc	218
14.4 Characterizing and ordering of environments	219
14.5 Ordering of attributes	222
15 Data laundry	223
15.1 Seeking exotics	224
15.2 Winsorization of anomalies	226
16 Choices of expression	231
16.1 Comparative scaling	231
16.2 Comparative responses	231
16.3 Re-expression of individual attributes	232
17 Seeking exotic values	233

17.1 SPLOMs	233
17.2 Alternative displays	245
17.3 Correlations among environments	254
18 Local analyses and displays	259
18.1 Local analysis within environments	259
18.2 Adjustment of structured sets of s^2 values	259
18.3 Choice of critical levels	262
18.4 Benjamini-Hochberg (FDR) procedure	264
18.5 Overlap plots	266
18.6 Dealing with ties in the BSD procedure	275
18.7 Profiles across environments and B-confidence	278
18.8 Some plausible developments for the BSD	285
19 Combined analyses	289
19.1 Analysis over environments	289
19.2 Analyses over attributes	297
20 Grouping and labelling genotypes	301
20.1 Grouping	301
20.2 Labelling	302
21 Idiolinkage, nearest and centroid	303
21.1 Nearest	303
21.2 Nearest algorithm	303
21.3 Centroid	304
21.4 Mean squared distance	306
22 Other blended forms; xpanded idiolinkage	307
22.1 Other blended forms	307
22.2 Xpanded linkage	307
22.3 Generalized xpansion	308
22.4 Techniques so far proposed	309
23 Robust forms of idiolinkage	311
23.1 Flexible Winsorization	311
24 Idiomax	313
24.1 Expansion	313
24.2 Plausible scope of exploration	315

25 Robust spherling	317
25.1 Robust orthogonalization	318
25.2 Simplification	319
25.3 Robust scaling	320
25.4 Comments	320
25.5 Relative importance?	320
26 A side issue and more careful spherling	323
26.1 Pattern of analysis	323
26.2 Choice of scaling	323
26.3 A possible exception	325
26.4 Postcomment	325
27 A suggestion about ‘G in or out’	327
27.1 Another suggestion	327
28 Detrivialization of one- and two-way tables: an introduction	333
28.1 The one-way case	333
28.2 The two-way case	335
29 Later attributes and subtables	337
29.1 The second attribute	337
29.2 The third attribute	339
29.3 Later attributes	339
29.4 Combination	340
29.5 Postcomment	340
30 Spherling the soybean data	341
30.1 Ordering of attributes	341
30.2 Robust orthogonalization	342
30.3 Robust scaling	347
31 Grouping the soybean genotypes	351
31.1 First-stage grouping	351
31.2 Second-stage grouping	354
31.3 Group responses	354
31.4 Alternative analyses	357
31.5 Partitioning variability within groups	357
32 Ordering the environments	363

32.1 Greedy close ordering of environments	363
33 One approach to plotting	373
33.1 Smoothing in general	374
33.2 Group 4 and attribute 1 as an example	375
33.3 Looking at the seven groups for attribute 1	384
33.4 Looking at the seven groups for all attributes	384
34 Smoothing short sequences (of perhaps 6 to 18 values)	399
34.1 3R 3pR E ₂	399
34.2 3 ⁺ R E ₂	402
35 Presentations across attributes	403
36 Choice of expression: generalities	405
36.1 Matching re-expressions	406
36.2 Purposes of choice of expression	408
36.3 Rarity of conflict between these goals	409
36.4 Tools for diagnosis	410
36.5 Combining <i>Q</i> -diagnoses	413
36.6 Breaking down the example	413
36.7 Weighted regression	413
37 Removable inhomogeneity of variability	417
37.1 Replication within environments (e.g. 2×40)	418
37.2 Replication between environments (e.g. 8×40)	418
37.3 Inhomogeneity of interaction size (e.g. 8×40 or 8×18)	418
37.4 Variability between replications within environments	419
37.5 The special case of two versions	426
37.6 Interaction variability	428
38 Removable non-additivity	433
38.1 Extended fits	433
38.2 Choice of expression to promote additivity of response	434
38.3 General case of several versions	435
39 Linearity of response	441
40 Relative importance and combination	443
40.1 Bowstring plots	444

41 Interpretation and hybridization	451
Appendices	453
Appendix 1	453
Appendix 2	473
Appendix 3	481
Appendix 4	482
Appendix 5	486
Appendix 6	487
Appendix 7	496
Appendix 8	503
Appendix 9	504
Appendix 10	508
Appendix 11	512
Appendix 12	518
Appendix 13	524
Appendix 14	530
Appendix 15	536
Appendix 16	542
Appendix 17	548
Appendix 18	552
Appendix 19	553
Appendix 20	556
Appendix 21	564
Appendix 22	573
References	575
Author Index	579
Subject Index	581