

# Contents

|                             |  |              |
|-----------------------------|--|--------------|
| <i>Special Features</i>     |  | <i>viii</i>  |
| <i>Detailed Contents</i>    |  | <i>ix</i>    |
| <i>Acknowledgments</i>      |  | <i>xxvi</i>  |
| <i>A Note to the Reader</i> |  | <i>xxxii</i> |
| <b>PART I</b>               | <b>INTRODUCTION TO THE CELL</b>  |              |
| 1.                          | Cells and Genomes  | 1            |
| 2.                          | Cell Chemistry and Biosynthesis  | 45           |
| 3.                          | Proteins   | 125          |
| <b>PART II</b>              | <b>BASIC GENETIC MECHANISMS</b>  |              |
| 4.                          | DNA, Chromosomes, and Genomes  | 195          |
| 5.                          | DNA Replication, Repair, and Recombination                                       | 263          |
| 6.                          | How Cells Read the Genome: From DNA to Protein                                   | 329          |
| 7.                          | Control of Gene Expression   | 411          |
| <b>PART III</b>             | <b>METHODS</b>   |              |
| 8.                          | Manipulating Proteins, DNA, and RNA  | 501          |
| 9.                          | Visualizing Cells  | 579          |
| <b>PART IV</b>              | <b>INTERNAL ORGANIZATION OF THE CELL</b>   |              |
| 10.                         | Membrane Structure   | 617          |
| 11.                         | Membrane Transport of Small Molecules and the Electrical Properties of Membranes | 651          |
| 12.                         | Intracellular Compartments and Protein Sorting                                   | 695          |
| 13.                         | Intracellular Vesicular Traffic  | 749          |
| 14.                         | Energy Conversion: Mitochondria and Chloroplasts                                 | 813          |
| 15.                         | Mechanisms of Cell Communication   | 879          |
| 16.                         | The Cytoskeleton   | 965          |
| 17.                         | The Cell Cycle   | 1053         |
| 18.                         | Apoptosis  | 1115         |
| <b>PART V</b>               | <b>CELLS IN THEIR SOCIAL CONTEXT</b>   |              |
| 19.                         | Cell Junctions, Cell Adhesion, and the Extracellular Matrix                      | 1131         |
| 20.                         | Cancer   | 1205         |
|                             | Chapters 21–25 available on Media DVD-ROM  |              |
| 21.                         | Sexual Reproduction: Meiosis, Germ Cells, and Fertilization                      | 1269         |
| 22.                         | Development of Multicellular Organisms   | 1305         |
| 23.                         | Specialized Tissues, Stem Cells, and Tissue Renewal                              | 1417         |
| 24.                         | Pathogens, Infection, and Innate Immunity  | 1485         |
| 25.                         | The Adaptive Immune System   | 1539         |
| <i>Glossary</i>             |  | <i>G–1</i>   |
| <i>Index</i>                |  | <i>I–1</i>   |
| <i>Tables</i>               | <i>The Genetic Code, Amino Acids</i>   | <i>T–1</i>   |

# Special Features

|            |  |               |
|------------|--|---------------|
| Table 1–1  | Some Genomes That Have Been Completely Sequenced   | p. 18         |
| Table 1–2  | The Numbers of Gene Families, Classified by Function, That Are Common to All Three Domains of the Living World | p. 24         |
| Table 2–1  | Covalent and Noncovalent Chemical Bonds  | p. 53         |
| Table 2–2  | The Types of Molecules That Form a Bacterial Cell  | p. 55         |
| Table 2–3  | Approximate Chemical Compositions of a Typical Bacterium and a Typical Mammalian Cell                          | p. 63         |
| Table 2–4  | Relationship Between the Standard Free-Energy Change, $\Delta G^\circ$ , and the Equilibrium Constant          | p. 77         |
| Panel 2–1  | Chemical Bonds and Groups Commonly Encountered in Biological Molecules   | pp. 106–107   |
| Panel 2–2  | Water and Its Influence on the Behavior of Biological Molecules  | pp. 108–109   |
| Panel 2–3  | The Principal Types of Weak Noncovalent Bonds that Hold Macromolecules Together                                | pp. 110–111   |
| Panel 2–4  | An Outline of Some of the Types of Sugars Commonly Found in Cells  | pp. 112–113   |
| Panel 2–5  | Fatty Acids and Other Lipids   | pp. 114–115   |
| Panel 2–6  | A Survey of the Nucleotides  | pp. 116–117   |
| Panel 2–7  | Free Energy and Biological Reactions   | pp. 118–119   |
| Panel 2–8  | Details of the 10 Steps of Glycolysis  | pp. 120–121   |
| Panel 2–9  | The Complete Citric Acid Cycle   | pp. 122–123   |
| Panel 3–1  | The 20 Amino Acids Found in Proteins   | pp. 128–129   |
| Panel 3–2  | Four Different Ways of Depicting a Small Protein, the SH2 Domain   | pp. 132–133   |
| Table 3–1  | Some Common Types of Enzymes   | p. 159        |
| Panel 3–3  | Some of the Methods Used to Study Enzymes  | pp. 162–163   |
| Table 4–1  | Some Vital Statistics for the Human Genome   | p. 206        |
| Table 5–3  | Three Major Classes of Transposable Elements   | p. 318        |
| Table 6–1  | Principal Types of RNAs Produced in Cells  | p. 336        |
| Panel 8–1  | Review of Classical Genetics   | pp. 554–555   |
| Table 10–1 | Approximate Lipid Compositions of Different Cell Membranes   | p. 624        |
| Table 11–1 | A Comparison of Ion Concentrations Inside and Outside a Typical Mammalian Cell                                 | p. 652        |
| Panel 11–2 | The Derivation of the Nernst Equation  | p. 670        |
| Panel 11–3 | Some Classical Experiments on the Squid Giant Axon   | p. 679        |
| Table 12–1 | Relative Volumes Occupied by the Major Intracellular Compartments in a Liver Cell (Hepatocyte)                 | p. 697        |
| Table 12–2 | Relative Amounts of Membrane Types in Two Kinds of Eucaryotic Cells  | p. 697        |
| Table 14–1 | Product Yields from the Oxidation of Sugars and Fats   | p. 824        |
| Panel 14–1 | Redox Potentials   | p. 830        |
| Table 15–5 | The Ras Superfamily of Monomeric GTPases   | p. 926        |
| Panel 16–2 | The Polymerization of Actin and Tubulin  | pp. 978–979   |
| Panel 16–3 | Accessory Proteins that Control the Assembly and Position of Cytoskeletal Filaments                            | pp. 994–995   |
| Table 17–2 | Summary of the Major Cell-Cycle Regulatory Proteins  | p. 1066       |
| Panel 17–1 | The Principle Stages of M Phase (Mitosis and Cytokinesis) in an Animal Cell                                    | pp. 1072–1073 |

# Detailed Contents

|   |          |  |           |
|---|----------|--|-----------|
| <b>Chapter 1 Cells and Genomes</b>  | <b>1</b> |  |           |
| THE UNIVERSAL FEATURES OF CELLS ON EARTH  | 1        | The World of Animal Cells Is Represented By a Worm, a Fly, a Mouse, and a Human                        | 36        |
| All Cells Store Their Hereditary Information in the Same Linear Chemical Code (DNA)                 | 2        | Studies in <i>Drosophila</i> Provide a Key to Vertebrate Development                                   | 37        |
| All Cells Replicate Their Hereditary Information by Templated Polymerization                        | 3        | The Vertebrate Genome Is a Product of Repeated Duplication   | 38        |
| All Cells Transcribe Portions of Their Hereditary Information into the Same Intermediary Form (RNA) | 4        | Genetic Redundancy Is a Problem for Geneticists, But It Creates Opportunities for Evolving Organisms   | 39        |
| All Cells Use Proteins as Catalysts   | 5        | The Mouse Serves as a Model for Mammals  | 39        |
| All Cells Translate RNA into Protein in the Same Way  | 6        | Humans Report on Their Own Peculiarities   | 40        |
| The Fragment of Genetic Information Corresponding to One Protein Is One Gene                        | 7        | We Are All Different in Detail   | 41        |
| Life Requires Free Energy   | 8        | Summary  | 42        |
| All Cells Function as Biochemical Factories Dealing with the Same Basic Molecular Building Blocks   | 8        | Problems   | 42        |
| All Cells Are Enclosed in a Plasma Membrane Across Which Nutrients and Waste Materials Must Pass    | 9        | References   | 44        |
| All Cells Are Enclosed in a Plasma Membrane Across Which Nutrients and Waste Materials Must Pass    | 9        |  |           |
| A Living Cell Can Exist with Fewer Than 500 Genes   | 10       | <b>Chapter 2 Cell Chemistry and Biosynthesis</b>   | <b>45</b> |
| Summary   | 11       | THE CHEMICAL COMPONENTS OF A CELL  | 45        |
| THE DIVERSITY OF GENOMES AND THE TREE OF LIFE   | 11       | Cells Are Made From a Few Types of Atoms   | 45        |
| Cells Can Be Powered by a Variety of Free Energy Sources  | 12       | The Outermost Electrons Determine How Atoms Interact   | 46        |
| Some Cells Fix Nitrogen and Carbon Dioxide for Others   | 13       | Covalent Bonds Form by the Sharing of Electrons  | 48        |
| The Greatest Biochemical Diversity Exists Among Prokaryotic Cells                                   | 14       | There Are Different Types of Covalent Bonds  | 50        |
| The Tree of Life Has Three Primary Branches: Bacteria, Archaea, and Eucaryotes                      | 14       | An Atom Often Behaves as if It Has a Fixed Radius  | 51        |
| Some Genes Evolve Rapidly; Others Are Highly Conserved  | 16       | Water Is the Most Abundant Substance in Cells  | 51        |
| Most Bacteria and Archaea Have 1000–6000 Genes  | 17       | Some Polar Molecules Are Acids and Bases   | 52        |
| New Genes Are Generated from Preexisting Genes  | 18       | Four Types of Noncovalent Attractions Help Bring Molecules Together in Cells                           | 53        |
| Gene Duplications Give Rise to Families of Related Genes Within a Single Cell                       | 19       | A Cell Is Formed from Carbon Compounds   | 54        |
| Genes Can Be Transferred Between Organisms, Both in the Laboratory and in Nature                    | 21       | Cells Contain Four Major Families of Small Organic Molecules   | 55        |
| Sex Results in Horizontal Exchanges of Genetic Information Within a Species                         | 22       | Sugars Provide an Energy Source for Cells and Are the Subunits of Polysaccharides                      | 55        |
| The Function of a Gene Can Often Be Deduced from Its Sequence                                       | 22       | Fatty Acids Are Components of Cell Membranes, as Well as a Source of Energy                            | 58        |
| More Than 200 Gene Families Are Common to All Three Primary Branches of the Tree of Life            | 23       | Amino Acids Are the Subunits of Proteins   | 59        |
| Mutations Reveal the Functions of Genes   | 23       | Nucleotides Are the Subunits of DNA and RNA  | 61        |
| Molecular Biologists Have Focused a Spotlight on <i>E. coli</i>                                     | 24       | The Chemistry of Cells Is Dominated by Macromolecules with Remarkable Properties                       | 62        |
| Summary   | 26       | Noncovalent Bonds Specify Both the Precise Shape of a Macromolecule and its Binding to Other Molecules | 63        |
| GENETIC INFORMATION IN EUKARYOTES   | 26       | Summary  | 65        |
| Eucaryotic Cells May Have Originated as Predators   | 26       | CATALYSIS AND THE USE OF ENERGY BY CELLS   | 65        |
| Modern Eucaryotic Cells Evolved from a Symbiosis  | 27       | Cell Metabolism Is Organized by Enzymes  | 66        |
| Eucaryotes Have Hybrid Genomes  | 30       | Biological Order Is Made Possible by the Release of Heat Energy from Cells                             | 66        |
| Eucaryotic Genomes Are Big  | 30       | Photosynthetic Organisms Use Sunlight to Synthesize Organic Molecules                                  | 68        |
| Eucaryotic Genomes Are Rich in Regulatory DNA   | 31       | Cells Obtain Energy by the Oxidation of Organic Molecules  | 70        |
| The Genome Defines the Program of Multicellular Development   | 31       | Oxidation and Reduction Involve Electron Transfers   | 71        |
| Many Eucaryotes Live as Solitary Cells: the Protists  | 32       | Enzymes Lower the Barriers That Block Chemical Reactions   | 72        |
| A Yeast Serves as a Minimal Model Eucaryote   | 33       | How Enzymes Find Their Substrates: The Enormous Rapidity of Molecular Motions                          | 74        |
| The Expression Levels of All The Genes of An Organism Can Be Monitored Simultaneously               | 34       | The Free-Energy Change for a Reaction Determines Whether It Can Occur                                  | 75        |
| To Make Sense of Cells, We Need Mathematics, Computers, and Quantitative Information                | 35       | The Concentration of Reactants Influences the Free-Energy Change and a Reaction's Direction            | 76        |
| <i>Arabidopsis</i> Has Been Chosen Out of 300,000 Species As a Model Plant                          | 36       | For Sequential Reactions, $\Delta G^\circ$ Values Are Additive   | 77        |
|   |          | Activated Carrier Molecules Are Essential for Biosynthesis   | 78        |
|   |          | The Formation of an Activated Carrier Is Coupled to an Energetically Favorable Reaction                | 79        |

|  |            |   |            |
|--|------------|---|------------|
| ATP Is the Most Widely Used Activated Carrier Molecule                                     | 80         | Molecular Tunnels Channel Substrates in Enzymes with Multiple Catalytic Sites   | 167        |
| Energy Stored in ATP Is Often Harnessed to Join Two Molecules Together                     | 81         | Multienzyme Complexes Help to Increase the Rate of Cell Metabolism  | 168        |
| NADH and NADPH Are Important Electron Carriers   | 82         | The Cell Regulates the Catalytic Activities of its Enzymes  | 169        |
| There Are Many Other Activated Carrier Molecules in Cells                                  | 83         | Allosteric Enzymes Have Two or More Binding Sites That Interact   | 171        |
| The Synthesis of Biological Polymers Is Driven by ATP Hydrolysis                           | 84         | Two Ligands Whose Binding Sites Are Coupled Must Reciprocally Affect Each Other's Binding   | 171        |
| <i>Summary</i>   | 87         | Symmetric Protein Assemblies Produce Cooperative Allosteric Transitions   | 172        |
| <b>HOW CELLS OBTAIN ENERGY FROM FOOD</b>   | 88         | The Allosteric Transition in Aspartate Transcarbamoylase Is Understood in Atomic Detail   | 173        |
| Glycolysis Is a Central ATP-Producing Pathway  | 88         | Many Changes in Proteins Are Driven by Protein Phosphorylation  | 175        |
| Fermentations Produce ATP in the Absence of Oxygen   | 89         | A Eucaryotic Cell Contains a Large Collection of Protein Kinases and Protein Phosphatases   | 176        |
| Glycolysis Illustrates How Enzymes Couple Oxidation to Energy Storage                      | 91         | The Regulation of Cdk and Src Protein Kinases Shows How a Protein Can Function as a Microchip   | 177        |
| Organisms Store Food Molecules in Special Reservoirs                                       | 91         | Proteins That Bind and Hydrolyze GTP Are Ubiquitous Cellular Regulators   | 178        |
| Most Animal Cells Derive Their Energy from Fatty Acids Between Meals                       | 95         | Regulatory Proteins Control the Activity of GTP-Binding Proteins by Determining Whether GTP or GDP Is Bound                             | 179        |
| Sugars and Fats Are Both Degraded to Acetyl CoA in Mitochondria                            | 96         | Large Protein Movements Can Be Generated From Small Ones  | 179        |
| The Citric Acid Cycle Generates NADH by Oxidizing Acetyl Groups to CO <sub>2</sub>         | 97         | Motor Proteins Produce Large Movements in Cells   | 181        |
| Electron Transport Drives the Synthesis of the Majority of the ATP in Most Cells           | 100        | Membrane-Bound Transporters Harness Energy to Pump Molecules Through Membranes  | 182        |
| Amino Acids and Nucleotides Are Part of the Nitrogen Cycle                                 | 100        | Proteins Often Form Large Complexes That Function as Protein Machines   | 184        |
| Metabolism Is Organized and Regulated  | 101        | Protein Machines with Interchangeable Parts Make Efficient Use of Genetic Information   | 184        |
| <i>Summary</i>   | 103        | The Activation of Protein Machines Often Involves Positioning Them at Specific Sites  | 185        |
| <i>Problems</i>  | 103        | Many Proteins Are Controlled by Multisite Covalent Modification   | 186        |
| <i>References</i>  | 124        | A Complex Network of Protein Interactions Underlies Cell Function   | 187        |
|  |            | <i>Summary</i>  | 190        |
|  |            | <i>Problems</i>   | 191        |
|  |            | <i>References</i>   | 193        |
| <b>Chapter 3 Proteins</b>  | <b>125</b> | <b>Chapter 4 DNA, Chromosomes, and Genomes</b>  | <b>195</b> |
| <b>THE SHAPE AND STRUCTURE OF PROTEINS</b>   | 125        | <b>THE STRUCTURE AND FUNCTION OF DNA</b>  | 197        |
| The Shape of a Protein Is Specified by Its Amino Acid Sequence                             | 125        | A DNA Molecule Consists of Two Complementary Chains of Nucleotides  | 197        |
| Proteins Fold into a Conformation of Lowest Energy   | 130        | The Structure of DNA Provides a Mechanism for Heredity  | 199        |
| The $\alpha$ Helix and the $\beta$ Sheet Are Common Folding Patterns                       | 131        | In Eucaryotes, DNA Is Enclosed in a Cell Nucleus  | 200        |
| Protein Domains Are Modular Units from which Larger Proteins Are Built                     | 135        | <i>Summary</i>  | 201        |
| Few of the Many Possible Polypeptide Chains Will Be Useful to Cells                        | 136        | <b>CHROMOSOMAL DNA AND ITS PACKAGING IN THE CHROMATIN FIBER</b>   | 202        |
| Proteins Can Be Classified into Many Families  | 137        | Eucaryotic DNA Is Packaged into a Set of Chromosomes  | 202        |
| Sequence Searches Can Identify Close Relatives   | 139        | Chromosomes Contain Long Strings of Genes   | 204        |
| Some Protein Domains Form Parts of Many Different Proteins                                 | 140        | The Nucleotide Sequence of the Human Genome Shows How Our Genes Are Arranged  | 205        |
| Certain Pairs of Domains Are Found Together in Many Proteins                               | 141        | Genome Comparisons Reveal Evolutionarily Conserved DNA Sequences  | 207        |
| The Human Genome Encodes a Complex Set of Proteins, Revealing Much That Remains Unknown    | 142        | Chromosomes Exist in Different States Throughout the Life of a Cell   | 208        |
| Larger Protein Molecules Often Contain More Than One Polypeptide Chain                     | 142        | Each DNA Molecule That Forms a Linear Chromosome Must Contain a Centromere, Two Telomeres, and Replication Origins                      | 209        |
| Some Proteins Form Long Helical Filaments  | 143        | DNA Molecules Are Highly Condensed in Chromosomes   | 210        |
| Many Protein Molecules Have Elongated, Fibrous Shapes                                      | 145        | Nucleosomes Are a Basic Unit of Eucaryotic Chromosome Structure   | 211        |
| Many Proteins Contain a Surprisingly Large Amount of Unstructured Polypeptide Chain        | 146        | The Structure of the Nucleosome Core Particle Reveals How DNA Is Packaged   | 212        |
| Covalent Cross-Linkages Often Stabilize Extracellular Proteins                             | 147        | Nucleosomes Have a Dynamic Structure, and Are Frequently Subjected to Changes Catalyzed by ATP-Dependent Chromatin-Remodeling Complexes | 215        |
| Protein Molecules Often Serve as Subunits for the Assembly of Large Structures             | 148        | Nucleosomes Are Usually Packed Together into a Compact Chromatin Fiber  | 216        |
| Many Structures in Cells Are Capable of Self-Assembly                                      | 149        | <i>Summary</i>  | 218        |
| Assembly Factors Often Aid the Formation of Complex Biological Structures                  | 151        | <b>THE REGULATION OF CHROMATIN STRUCTURE</b>  | 219        |
| <i>Summary</i>   | 152        | Some Early Mysteries Concerning Chromatin Structure   | 220        |
| <b>PROTEIN FUNCTION</b>  | 152        |   |            |
| All Proteins Bind to Other Molecules   | 153        |   |            |
| The Surface Conformation of a Protein Determines Its Chemistry                             | 154        |   |            |
| Sequence Comparisons Between Protein Family Members Highlight Crucial Ligand-Binding Sites | 155        |   |            |
| Proteins Bind to Other Proteins Through Several Types of Interfaces                        | 156        |   |            |
| Antibody Binding Sites Are Especially Versatile  | 156        |   |            |
| The Equilibrium Constant Measures Binding Strength   | 157        |   |            |
| Enzymes Are Powerful and Highly Specific Catalysts   | 158        |   |            |
| Substrate Binding Is the First Step in Enzyme Catalysis                                    | 159        |   |            |
| Enzymes Speed Reactions by Selectively Stabilizing Transition States                       | 160        |   |            |
| Enzymes Can Use Simultaneous Acid and Base Catalysis                                       | 160        |   |            |
| Lysozyme Illustrates How an Enzyme Works   | 161        |   |            |
| Tightly Bound Small Molecules Add Extra Functions to Proteins                              | 166        |   |            |

|  |            |   |     |
|--|------------|---|-----|
| Heterochromatin Is Highly Organized and Unusually Resistant to Gene Expression   | 220        | DNA REPLICATION MECHANISMS  | 266 |
| The Core Histones Are Covalently Modified at Many Different Sites  | 222        | Base-Pairing Underlies DNA Replication and DNA Repair   | 266 |
| Chromatin Acquires Additional Variety through the Site-Specific Insertion of a Small Set of Histone Variants                               | 224        | The DNA Replication Fork Is Asymmetrical  | 266 |
| The Covalent Modifications and the Histone Variants Act in Concert to Produce a "Histone Code" That Helps to Determine Biological Function | 224        | The High Fidelity of DNA Replication Requires Several Proofreading Mechanisms                                     | 268 |
| A Complex of Code-Reader and Code-Writer Proteins Can Spread Specific Chromatin Modifications for Long Distances Along a Chromosome        | 226        | Only DNA Replication in the 5'-to-3' Direction Allows Efficient Error Correction                                  | 271 |
| Barrier DNA Sequences Block the Spread of Reader-Writer Complexes and Thereby Separate Neighboring Chromatin Domains                       | 227        | A Special Nucleotide-Polymerizing Enzyme Synthesizes Short RNA Primer Molecules on the Lagging Strand             | 272 |
| The Chromatin in Centromeres Reveals How Histone Variants Can Create Special Structures  | 228        | Special Proteins Help to Open Up the DNA Double Helix in Front of the Replication Fork                            | 273 |
| Chromatin Structures Can Be Directly Inherited   | 230        | A Sliding Ring Holds a Moving DNA Polymerase onto the DNA   | 273 |
| Chromatin Structures Add Unique Features to Eucaryotic Chromosome Function   | 231        | The Proteins at a Replication Fork Cooperate to Form a Replication Machine  | 275 |
| <i>Summary</i>   | 233        | A Strand-Directed Mismatch Repair System Removes Replication Errors That Escape from the Replication Machine      | 276 |
| THE GLOBAL STRUCTURE OF CHROMOSOMES  | 233        | DNA Topoisomerases Prevent DNA Tangling During Replication  | 278 |
| Chromosomes Are Folded into Large Loops of Chromatin   | 234        | DNA Replication Is Fundamentally Similar in Eucaryotes and Bacteria   | 280 |
| Polytene Chromosomes Are Uniquely Useful for Visualizing Chromatin Structures  | 236        | <i>Summary</i>  | 281 |
| There Are Multiple Forms of Heterochromatin  | 238        | THE INITIATION AND COMPLETION OF DNA REPLICATION IN CHROMOSOMES   | 281 |
| Chromatin Loops Decondense When the Genes Within Them Are Expressed  | 239        | DNA Synthesis Begins at Replication Origins   | 281 |
| Chromatin Can Move to Specific Sites Within the Nucleus to Alter Their Gene Expression   | 239        | Bacterial Chromosomes Typically Have a Single Origin of DNA Replication   | 282 |
| Networks of Macromolecules Form a Set of Distinct Biochemical Environments inside the Nucleus  | 241        | Eucaryotic Chromosomes Contain Multiple Origins of Replication  | 282 |
| Mitotic Chromosomes Are Formed from Chromatin in Its Most Condensed State  | 243        | In Eucaryotes DNA Replication Takes Place During Only One Part of the Cell Cycle                                  | 284 |
| <i>Summary</i>   | 245        | Different Regions on the Same Chromosome Replicate at Distinct Times in S Phase                                   | 285 |
| HOW GENOMES EVOLVE   | 245        | Highly Condensed Chromatin Replicates Late, While Genes in Less Condensed Chromatin Tend to Replicate Early       | 285 |
| Genome Alterations Are Caused by Failures of the Normal Mechanisms for Copying and Maintaining DNA   | 246        | Well-Defined DNA Sequences Serve as Replication Origins in a Simple Eucaryote, the Budding Yeast                  | 286 |
| The Genome Sequences of Two Species Differ in Proportion to the Length of Time That They Have Separately Evolved                           | 247        | A Large Multisubunit Complex Binds to Eucaryotic Origins of Replication   | 287 |
| Phylogenetic Trees Constructed from a Comparison of DNA Sequences Trace the Relationships of All Organisms                                 | 248        | The Mammalian DNA Sequences That Specify the Initiation of Replication Have Been Difficult to Identify            | 288 |
| A Comparison of Human and Mouse Chromosomes Shows How the Structures of Genomes Diverge  | 249        | New Nucleosomes Are Assembled Behind the Replication Fork   | 289 |
| The Size of a Vertebrate Genome Reflects the Relative Rates of DNA Addition and DNA Loss in a Lineage                                      | 251        | The Mechanisms of Eucaryotic Chromosome Duplication Ensure That Patterns of Histone Modification Can Be Inherited | 290 |
| We Can Reconstruct the Sequence of Some Ancient Genomes  | 251        | Telomerase Replicates the Ends of Chromosomes   | 292 |
| Multispecies Sequence Comparisons Identify Important DNA Sequences of Unknown Function   | 252        | Telomere Length Is Regulated by Cells and Organisms   | 293 |
| Accelerated Changes in Previously Conserved Sequences Can Help Decipher Critical Steps in Human Evolution                                  | 253        | <i>Summary</i>  | 294 |
| Gene Duplication Provides an Important Source of Genetic Novelty During Evolution  | 253        | DNA REPAIR  | 295 |
| Duplicated Genes Diverge   | 254        | Without DNA Repair, Spontaneous DNA Damage Would Rapidly Change DNA Sequences                                     | 296 |
| The Evolution of the Globin Gene Family Shows How DNA Duplications Contribute to the Evolution of Organisms                                | 256        | The DNA Double Helix Is Readily Repaired  | 296 |
| Genes Encoding New Proteins Can Be Created by the Recombination of Exons   | 257        | DNA Damage Can Be Removed by More Than One Pathway  | 297 |
| Neutral Mutations Often Spread to Become Fixed in a Population, with a Probability that Depends on Population Size                         | 257        | Coupling DNA Repair to Transcription Ensures That the Cell's Most Important DNA Is Efficiently Repaired           | 299 |
| A Great Deal Can Be Learned from Analyses of the Variation Among Humans  | 258        | The Chemistry of the DNA Bases Facilitates Damage Detection   | 300 |
| <i>Summary</i>   | 260        | Special DNA Polymerases Are Used in Emergencies to Repair DNA   | 302 |
| <i>Problems</i>  | 260        | Double-Strand Breaks Are Efficiently Repaired   | 302 |
| <i>References</i>  | 262        | DNA Damage Delays Progression of the Cell Cycle   | 303 |
| <b>Chapter 5 DNA Replication, Repair, and Recombination</b>  | <b>263</b> | <i>Summary</i>  | 304 |
| THE MAINTENANCE OF DNA SEQUENCES   | 263        | HOMOLOGOUS RECOMBINATION  | 304 |
| Mutation Rates Are Extremely Low   | 263        | Homologous Recombination Has Many Uses in the Cell  | 304 |
| Low Mutation Rates Are Necessary for Life as We Know It  | 265        | Homologous Recombination Has Common Features in All Cells   | 305 |
| <i>Summary</i>   | 265        | DNA Base-Pairing Guides Homologous Recombination  | 305 |
|  |            | The RecA Protein and its Homologs Enable a DNA Single Strand to Pair with a Homologous Region of DNA Double Helix | 307 |
|  |            | Branch Migration Can Either Enlarge Hetroduplex Regions or Release Newly Synthesized DNA as a Single Strand       | 308 |
|  |            | Homologous Recombination Can Flawlessly Repair Double-Stranded Breaks in DNA                                      | 308 |
|  |            | Cells Carefully Regulate the Use of Homologous Recombination in DNA Repair  | 310 |
|  |            | Holliday Junctions Are Often Formed During Homologous Recombination Events  | 311 |

|  |            |   |            |
|--|------------|---|------------|
| Meiotic Recombination Begins with a Programmed Double-Strand Break                               | 312        | FROM RNA TO PROTEIN   | 366        |
| Homologous Recombination Often Results in Gene Conversion  | 314        | An mRNA Sequence Is Decoded in Sets of Three Nucleotide   | 367        |
| Mismatch Proofreading Prevents Promiscuous Recombination   |            | tRNA Molecules Match Amino Acids to Codons in mRNA  | 368        |
| Between Two Poorly Matched DNA Sequences   | 315        | tRNAs Are Covalently Modified Before They Exit from the Nucleus                                 | 369        |
| <i>Summary</i>   | 316        | Specific Enzymes Couple Each Amino Acid to Its Appropriate tRNA Molecule                        | 370        |
| <b>TRANSPOSITION AND CONSERVATIVE SITE-SPECIFIC RECOMBINATION</b>                                | 316        | Editing by RNA Synthetases Ensures Accuracy   | 371        |
| Through Transposition, Mobile Genetic Elements Can Insert Into Any DNA Sequence                  | 317        | Amino Acids Are Added to the C-terminal End of a Growing Polypeptide Chain                      | 373        |
| DNA-Only Transposons Move by Both Cut-and-Paste and Replicative Mechanisms                       | 317        | The RNA Message Is Decoded in Ribosomes   | 373        |
| Some Viruses Use a Transposition Mechanism to Move Themselves into Host Cell Chromosomes         | 319        | Elongation Factors Drive Translation Forward and Improve Its Accuracy                           | 377        |
| Retroviral-like Retrotransposons Resemble Retroviruses, but Lack a Protein Coat                  | 320        | The Ribosome Is a Ribozyme  | 378        |
| A Large Fraction of the Human Genome Is Composed of Nonretroviral Retrotransposons               | 321        | Nucleotide Sequences in mRNA Signal Where to Start Protein Synthesis                            | 379        |
| Different Transposable Elements Predominate in Different Organisms                               | 322        | Stop Codons Mark the End of Translation   | 381        |
| Genome Sequences Reveal the Approximate Times that Transposable Elements Have Moved              | 323        | Proteins Are Made on Polyribosomes  | 381        |
| Conservative Site-Specific Recombination Can Reversibly Rearrange DNA                            | 323        | There Are Minor Variations in the Standard Genetic Code   | 382        |
| Conservative Site-Specific Recombination Was Discovered in Bacteriophage $\lambda$               | 324        | Inhibitors of Prokaryotic Protein Synthesis Are Useful as Antibiotics                           | 383        |
| Conservative Site-Specific Recombination Can Be Used to Turn Genes On or Off                     | 324        | Accuracy in Translation Requires the Expenditure of Free Energy                                 | 385        |
| <i>Summary</i>   | 326        | Quality Control Mechanisms Act to Prevent Translation of Damaged mRNAs                          | 385        |
| <i>Problems</i>  | 327        | Some Proteins Begin to Fold While Still Being Synthesized                                       | 387        |
| <i>References</i>  | 328        | Molecular Chaperones Help Guide the Folding of Most Proteins                                    | 388        |
| <b>Chapter 6 How Cells Read the Genome: From DNA to Protein</b>                                  | <b>329</b> | Exposed Hydrophobic Regions Provide Critical Signals for Protein Quality Control                | 390        |
| FROM DNA TO RNA  | 331        | The Proteasome Is a Compartmentalized Protease with Sequestered Active Sites                    | 391        |
| Portions of DNA Sequence Are Transcribed into RNA  | 332        | An Elaborate Ubiquitin-Conjugating System Marks Proteins for Destruction                        | 393        |
| Transcription Produces RNA Complementary to One Strand of DNA                                    | 333        | Many Proteins Are Controlled by Regulated Destruction   | 395        |
| Cells Produce Several Types of RNA   | 335        | Abnormally Folded Proteins Can Aggregate to Cause Destructive Human Diseases                    | 396        |
| Signals Encoded in DNA Tell RNA Polymerase Where to Start and Stop                               | 336        | There Are Many Steps From DNA to Protein  | 399        |
| Transcription Start and Stop Signals Are Heterogeneous in Nucleotide Sequence                    | 338        | <i>Summary</i>  | 399        |
| Transcription Initiation in Eucaryotes Requires Many Proteins                                    | 339        | <b>THE RNA WORLD AND THE ORIGINS OF LIFE</b>  | 400        |
| RNA Polymerase II Requires General Transcription Factors   | 340        | Life Requires Stored Information  | 401        |
| Polymerase II Also Requires Activator, Mediator, and Chromatin-Modifying Proteins                | 342        | Polynucleotides Can Both Store Information and Catalyze Chemical Reactions                      | 401        |
| Transcription Elongation Produces Superhelical Tension in DNA                                    | 343        | A Pre-RNA World May Predate the RNA World   | 402        |
| Transcription Elongation in Eucaryotes Is Tightly Coupled to RNA Processing                      | 345        | Single-Stranded RNA Molecules Can Fold into Highly Elaborate Structures                         | 403        |
| RNA Capping Is the First Modification of Eucaryotic Pre-mRNAs                                    | 346        | Self-Replicating Molecules Undergo Natural Selection  | 404        |
| RNA Splicing Removes Intron Sequences from Newly Transcribed Pre-mRNAs                           | 347        | How Did Protein Synthesis Evolve?   | 407        |
| Nucleotide Sequences Signal Where Splicing Occurs  | 349        | All Present-Day Cells Use DNA as Their Hereditary Material                                      | 408        |
| RNA Splicing Is Performed by the Spliceosome   | 349        | <i>Summary</i>  | 408        |
| The Spliceosome Uses ATP Hydrolysis to Produce a Complex Series of RNA-RNA Rearrangements        | 351        | <i>Problems</i>   | 409        |
| Other Properties of Pre-mRNA and Its Synthesis Help to Explain the Choice of Proper Splice Sites | 352        | <i>References</i>   | 410        |
| A Second Set of snRNPs Splice a Small Fraction of Intron Sequences in Animals and Plants         | 353        | <b>Chapter 7 Control of Gene Expression</b>   | <b>411</b> |
| RNA Splicing Shows Remarkable Plasticity   | 355        | <b>AN OVERVIEW OF GENE CONTROL</b>  | 411        |
| Spliceosome-Catalyzed RNA Splicing Probably Evolved from Self-Splicing Mechanisms                | 355        | The Different Cell Types of a Multicellular Organism Contain the Same DNA                       | 411        |
| RNA-Processing Enzymes Generate the 3' End of Eucaryotic mRNAs                                   | 357        | Different Cell Types Synthesize Different Sets of Proteins                                      | 412        |
| Mature Eucaryotic mRNAs Are Selectively Exported from the Nucleus                                | 358        | External Signals Can Cause a Cell to Change the Expression of Its Genes                         | 413        |
| Many Noncoding RNAs Are Also Synthesized and Processed in the Nucleus                            | 360        | Gene Expression Can Be Regulated at Many of the Steps in the Pathway from DNA to RNA to Protein | 415        |
| The Nucleolus Is a Ribosome-Producing Factory  | 362        | <i>Summary</i>  | 415        |
| The Nucleus Contains a Variety of Subnuclear Structures  | 363        | <b>DNA-BINDING MOTIFS IN GENE REGULATORY PROTEINS</b>   | 416        |
| <i>Summary</i>   | 366        | Gene Regulatory Proteins Were Discovered Using Bacterial Genetics                               | 416        |
|  |            | The Outside of the DNA Helix Can Be Read by Proteins  | 416        |
|  |            | Short DNA Sequences Are Fundamental Components of Genetic Switches                              | 418        |
|  |            | Gene Regulatory Proteins Contain Structural Motifs That Can Read DNA Sequences                  | 418        |
|  |            | The Helix-Turn-Helix Motif Is One of the Simplest and Most Common DNA-Binding Motifs            | 419        |

|  |     |  |            |
|--|-----|--|------------|
| Homeodomain Proteins Constitute a Special Class of Helix–Turn–Helix Proteins   | 420 | Expression of a Critical Gene Regulatory Protein Can Trigger the Expression of a Whole Battery of Downstream Genes | 463        |
| There Are Several Types of DNA-Binding Zinc Finger Motifs  | 421 | Combinatorial Gene Control Creates Many Different Cell Types in Eucaryotes   | 464        |
| $\beta$ sheets Can Also Recognize DNA  | 422 | A Single Gene Regulatory Protein Can Trigger the Formation of an Entire Organ                                      | 465        |
| Some Proteins Use Loops That Enter the Major and Minor Groove to Recognize DNA   | 423 | The Pattern of DNA Methylation Can Be Inherited When Vertebrate Cells Divide                                       | 467        |
| The Leucine Zipper Motif Mediates Both DNA Binding and Protein Dimerization  | 423 | Genomic Imprinting Is Based on DNA Methylation   | 468        |
| Heterodimerization Expands the Repertoire of DNA Sequences That Gene Regulatory Proteins Can Recognize   | 424 | CG-Rich Islands Are Associated with Many Genes in Mammals  | 470        |
| The Helix–Loop–Helix Motif Also Mediates Dimerization and DNA Binding  | 425 | Epigenetic Mechanisms Ensure That Stable Patterns of Gene Expression Can Be Transmitted to Daughter Cells          | 471        |
| It Is Not Yet Possible to Predict the DNA Sequences Recognized by All Gene Regulatory Proteins   | 426 | Chromosome-Wide Alterations in Chromatin Structure Can Be Inherited  | 473        |
| A Gel-Mobility Shift Assay Readily Detects Sequence-Specific DNA-Binding Proteins  | 427 | The Control of Gene Expression is Intrinsically Noisy  | 476        |
| DNA Affinity Chromatography Facilitates the Purification of Sequence-Specific DNA-Binding Proteins   | 428 | <i>Summary</i>   | 477        |
| The DNA Sequence Recognized by a Gene Regulatory Protein Can Be Determined Experimentally  | 429 | POST-TRANSCRIPTIONAL CONTROLS  | 477        |
| Phylogenetic Footprinting Identifies DNA Regulatory Sequences Through Comparative Genomics   | 431 | Transcription Attenuation Causes the Premature Termination of Some RNA Molecules                                   | 477        |
| Chromatin Immunoprecipitation Identifies Many of the Sites That Gene Regulatory Proteins Occupy in Living Cells                                    | 431 | Riboswitches Might Represent Ancient Forms of Gene Control   | 478        |
| <i>Summary</i>   | 432 | Alternative RNA Splicing Can Produce Different Forms of a Protein from the Same Gene                               | 479        |
| HOW GENETIC SWITCHES WORK  | 432 | The Definition of a Gene Has Had to Be Modified Since the Discovery of Alternative RNA Splicing                    | 480        |
| The Tryptophan Repressor Is a Simple Switch That Turns Genes On and Off in Bacteria  | 433 | Sex Determination in <i>Drosophila</i> Depends on a Regulated Series of RNA Splicing Events                        | 481        |
| Transcriptional Activators Turn Genes On   | 435 | A Change in the Site of RNA Transcript Cleavage and Poly-A Addition Can Change the C-terminus of a Protein         | 482        |
| A Transcriptional Activator and a Transcriptional Repressor Control the <i>Lac</i> Operon  | 435 | RNA Editing Can Change the Meaning of the RNA Message  | 483        |
| DNA Looping Occurs During Bacterial Gene Regulation  | 437 | RNA Transport from the Nucleus Can Be Regulated  | 485        |
| Bacteria Use Interchangeable RNA Polymerase Subunits to Help Regulate Gene Transcription   | 438 | Some mRNAs Are Localized to Specific Regions of the Cytoplasm  | 486        |
| Complex Switches Have Evolved to Control Gene Transcription in Eucaryotes  | 439 | The 5' and 3' Untranslated Regions of mRNAs Control Their Translation  | 488        |
| A Eucaryotic Gene Control Region Consists of a Promoter Plus Regulatory DNA Sequences  | 440 | The Phosphorylation of an Initiation Factor Regulates Protein Synthesis Globally                                   | 488        |
| Eucaryotic Gene Activator Proteins Promote the Assembly of RNA Polymerase and the General Transcription Factors at the Startpoint of Transcription | 441 | Initiation at AUG Codons Upstream of the Translation Start Can Regulate Eucaryotic Translation Initiation          | 489        |
| Eucaryotic Gene Activator Proteins Also Modify Local Chromatin Structure   | 442 | Internal Ribosome Entry Sites Provide Opportunities for Translation Control  | 491        |
| Gene Activator Proteins Work Synergistically   | 444 | Changes in mRNA Stability Can Regulate Gene Expression   | 492        |
| Eucaryotic Gene Repressor Proteins Can Inhibit Transcription in Various Ways   | 445 | Cytoplasmic Poly-A Addition Can Regulate Translation   | 493        |
| Eucaryotic Gene Regulatory Proteins Often Bind DNA Cooperatively   | 445 | Small Noncoding RNA Transcripts Regulate Many Animal and Plant Genes   | 493        |
| Complex Genetic Switches That Regulate <i>Drosophila</i> Development Are Built Up from Smaller Modules   | 447 | RNA Interference Is a Cell Defense Mechanism   | 495        |
| The <i>Drosophila Eve</i> Gene Is Regulated by Combinatorial Controls  | 448 | RNA Interference Can Direct Heterochromatin Formation  | 496        |
| Complex Mammalian Gene Control Regions Are Also Constructed from Simple Regulatory Modules   | 450 | RNA Interference Has Become a Powerful Experimental Tool   | 497        |
| Insulators Are DNA Sequences That Prevent Eucaryotic Gene Regulatory Proteins from Influencing Distant Genes                                       | 452 | <i>Summary</i>   | 497        |
| Gene Switches Rapidly Evolve   | 453 | <i>Problems</i>  | 497        |
| <i>Summary</i>   | 453 | <i>References</i>  | 499        |
| THE MOLECULAR GENETIC MECHANISMS THAT CREATE SPECIALIZED CELL TYPES  | 454 | <b>Chapter 8 Manipulating Proteins, DNA, and RNA</b>   | <b>501</b> |
| DNA Rearrangements Mediate Phase Variation in Bacteria   | 454 | ISOLATING CELLS AND GROWING THEM IN CULTURE  | 501        |
| A Set of Gene Regulatory Proteins Determines Cell Type in a Budding Yeast  | 455 | Cells Can Be Isolated from Intact Tissues  | 502        |
| Two Proteins That Repress Each Other's Synthesis Determine the Heritable State of Bacteriophage Lambda   | 457 | Cells Can Be Grown in Culture  | 502        |
| Simple Gene Regulatory Circuits Can Be Used to Make Memory Devices   | 458 | Eucaryotic Cell Lines Are a Widely Used Source of Homogeneous Cells  | 505        |
| Transcriptional Circuits Allow the Cell to Carry Out Logic Operations  | 459 | Embryonic Stem Cells Could Revolutionize Medicine  | 505        |
| Synthetic Biology Creates New Devices from Existing Biological Parts   | 460 | Somatic Cell Nuclear Transplantation May Provide a Way to Generate Personalized Stem Cells                         | 507        |
| Circadian Clocks Are Based on Feedback Loops in Gene Regulation  | 460 | Hybridoma Cell Lines Are Factories That Produce Monoclonal Antibodies  | 508        |
| A Single Gene Regulatory Protein Can Coordinate the Expression of a Set of Genes   | 462 | <i>Summary</i>   | 510        |
|  |     | PURIFYING PROTEINS   | 510        |
|  |     | Cells Can Be Separated into Their Component Fractions  | 510        |
|  |     | Cell Extracts Provide Accessible Systems to Study Cell Functions   | 511        |
|  |     | Proteins Can Be Separated by Chromatography  | 512        |
|  |     | Affinity Chromatography Exploits Specific Binding Sites on Proteins  | 513        |
|  |     | Genetically-Engineered Tags Provide an Easy Way to Purify Proteins   | 514        |

|   |            |   |            |
|---|------------|---|------------|
| Purified Cell-Free Systems Are Required for the Precise Dissection of Molecular Functions                         | 516        | Large Collections of Tagged Knockouts Provide a Tool for Examining the Function of Every Gene in an Organism  | 569        |
| <i>Summary</i>  | 516        | RNA Interference Is a Simple and Rapid Way to Test Gene Function  | 571        |
| <b>ANALYZING PROTEINS</b>   | <b>517</b> | Reporter Genes and <i>In Situ</i> Hybridization Reveal When and Where a Gene Is Expressed                     | 572        |
| Proteins Can Be Separated by SDS Polyacrylamide-Gel Electrophoresis   | 517        | Expression of Individual Genes Can Be Measured Using Quantitative RT-PCR                                      | 573        |
| Specific Proteins Can Be Detected by Blotting with Antibodies   | 518        | Microarrays Monitor the Expression of Thousands of Genes at Once  | 574        |
| Mass Spectrometry Provides a Highly Sensitive Method for Identifying Unknown Proteins                             | 519        | Single-Cell Gene Expression Analysis Reveals Biological "Noise"   | 575        |
| Two-Dimensional Separation Methods are Especially Powerful  | 521        | <i>Summary</i>  | 576        |
| Hydrodynamic Measurements Reveal the Size and Shape of a Protein Complex  | 522        | <i>Problems</i>   | 576        |
| Sets of Interacting Proteins Can Be Identified by Biochemical Methods   | 523        | <i>References</i>   | 578        |
| Protein-Protein Interactions Can Also Be Identified by a Two-Hybrid Technique in Yeast                            | 523        | <b>Chapter 9 Visualizing Cells</b>  | <b>579</b> |
| Combining Data Derived from Different Techniques Produces Reliable Protein-Interaction Maps                       | 524        | <b>LOOKING AT CELLS IN THE LIGHT MICROSCOPE</b>   | <b>579</b> |
| Optical Methods Can Monitor Protein Interactions in Real Time   | 524        | The Light Microscope Can Resolve Details 0.2 $\mu\text{m}$ Apart  | 580        |
| Some Techniques Can Monitor Single Molecules  | 526        | Living Cells Are Seen Clearly in a Phase-Contrast or a Differential-Interference-Contrast Microscope          | 583        |
| Protein Function Can Be Selectively Disrupted with Small Molecules  | 527        | Images Can Be Enhanced and Analyzed by Digital Techniques   | 583        |
| Protein Structure Can Be Determined Using X-Ray Diffraction   | 527        | Intact Tissues Are Usually Fixed and Sectioned before Microscopy  | 585        |
| NMR Can Be Used to Determine Protein Structure in Solution  | 529        | Specific Molecules Can Be Located in Cells by Fluorescence Microscopy   | 586        |
| Protein Sequence and Structure Provide Clues About Protein Function   | 530        | Antibodies Can Be Used to Detect Specific Molecules   | 588        |
| <i>Summary</i>  | 531        | Imaging of Complex Three-Dimensional Objects Is Possible with the Optical Microscope                          | 589        |
| <b>ANALYZING AND MANIPULATING DNA</b>   | <b>532</b> | The Confocal Microscope Produces Optical Sections by Excluding Out-of-Focus Light                             | 590        |
| Restriction Nucleases Cut Large DNA Molecules into Fragments  | 532        | Fluorescent Proteins Can Be Used to Tag Individual Proteins in Living Cells and Organisms                     | 592        |
| Gel Electrophoresis Separates DNA Molecules of Different Sizes  | 534        | Protein Dynamics Can Be Followed in Living Cells  | 593        |
| Purified DNA Molecules Can Be Specifically Labeled with Radioisotopes or Chemical Markers <i>in vitro</i>         | 535        | Light-Emitting Indicators Can Measure Rapidly Changing Intracellular Ion Concentrations                       | 596        |
| Nucleic Acid Hybridization Reactions Provide a Sensitive Way of Detecting Specific Nucleotide Sequences           | 535        | Several Strategies Are Available by Which Membrane-Impermeant Substances Can Be Introduced into Cells         | 597        |
| Northern and Southern Blotting Facilitate Hybridization with Electrophoretically Separated Nucleic Acid Molecules | 538        | Light Can Be Used to Manipulate Microscopic Objects As Well As to Image Them                                  | 598        |
| Genes Can Be Cloned Using DNA Libraries   | 540        | Single Molecules Can Be Visualized by Using Total Internal Reflection Fluorescence Microscopy                 | 599        |
| Two Types of DNA Libraries Serve Different Purposes   | 541        | Individual Molecules Can Be Touched and Moved Using Atomic Force Microscopy                                   | 600        |
| cDNA Clones Contain Uninterrupted Coding Sequences  | 544        | Molecules Can Be Labeled with Radioisotopes   | 600        |
| Genes Can Be Selectively Amplified by PCR   | 544        | Radioisotopes Are Used to Trace Molecules in Cells and Organisms  | 602        |
| Cells Can Be Used As Factories to Produce Specific Proteins   | 546        | <i>Summary</i>  | 603        |
| Proteins and Nucleic Acids Can Be Synthesized Directly by Chemical Reactions                                      | 548        | <b>LOOKING AT CELLS AND MOLECULES IN THE ELECTRON MICROSCOPE</b>  | <b>604</b> |
| DNA Can Be Rapidly Sequenced  | 548        | The Electron Microscope Resolves the Fine Structure of the Cell   | 604        |
| Nucleotide Sequences Are Used to Predict the Amino Acid Sequences of Proteins                                     | 550        | Biological Specimens Require Special Preparation for the Electron Microscope                                  | 605        |
| The Genomes of Many Organisms Have Been Fully Sequenced   | 551        | Specific Macromolecules Can Be Localized by Immunogold Electron Microscopy                                    | 606        |
| <i>Summary</i>  | 552        | Images of Surfaces Can Be Obtained by Scanning Electron Microscopy  | 607        |
| <b>STUDYING GENE EXPRESSION AND FUNCTION</b>  | <b>553</b> | Metal Shadowing Allows Surface Features to Be Examined at High Resolution by Transmission Electron Microscopy | 608        |
| Classical Genetics Begins by Disrupting a Cell Process by Random Mutagenesis                                      | 553        | Negative Staining and Cryoelectron Microscopy Both Allow Macromolecules to Be Viewed at High Resolution       | 610        |
| Genetic Screens Identify Mutants with Specific Abnormalities  | 556        | Multiple Images Can Be Combined to Increase Resolution  | 610        |
| Mutations Can Cause Loss or Gain of Protein Function  | 557        | Different Views of a Single Object Can Be Combined to Give a Three-Dimensional Reconstruction                 | 612        |
| Complementation Tests Reveal Whether Two Mutations Are in the Same Gene or Different Genes                        | 558        | <i>Summary</i>  | 612        |
| Genes Can Be Ordered in Pathways by Epistasis Analysis  | 558        | <i>Problems</i>   | 614        |
| Genes Identified by Mutations Can Be Cloned   | 559        | <i>References</i>   | 615        |
| Human Genetics Presents Special Problems and Special Opportunities  | 560        | <b>Chapter 10 Membrane Structure</b>  | <b>617</b> |
| Human Genes Are Inherited in Haplotype Blocks, Which Can Aid in the Search for Mutations That Cause Disease       | 561        | <b>THE LIPID BILAYER</b>  | <b>617</b> |
| Complex Traits Are Influenced by Multiple Genes   | 563        | Phosphoglycerides, Sphingolipids, and Sterols Are the Major Lipids in Cell Membranes                          | 618        |
| Reverse Genetics Begins with a Known Gene and Determines Which Cell Processes Require Its Function                | 563        | Phospholipids Spontaneously Form Bilayers   | 620        |
| Genes Can Be Re-Engineered in Several Ways  | 564        |   |            |
| Engineered Genes Can Be Inserted into the Germ Line of Many Organisms   | 565        |   |            |
| Animals Can Be Genetically Altered  | 566        |   |            |
| Transgenic Plants Are Important for Both Cell Biology and Agriculture   | 568        |   |            |



|  |            |  |            |
|--|------------|--|------------|
| The Lipid Bilayer Is a Two-Dimensional Fluid   | 621        | Patch-Clamp Recording Indicates That Individual Gated Channels   | 680        |
| The Fluidity of a Lipid Bilayer Depends on Its Composition   | 622        | Open in an All-or-Nothing Fashion  |            |
| Despite Their Fluidity, Lipid Bilayers Can Form Domains of Different Compositions  | 624        | Voltage-Gated Cation Channels Are Evolutionarily and Structurally Related  | 682        |
| Lipid Droplets Are Surrounded by a Phospholipid Monolayer  | 625        | Transmitter-Gated Ion Channels Convert Chemical Signals into Electrical Ones at Chemical Synapses                          | 682        |
| The Asymmetry of the Lipid Bilayer Is Functionally Important   | 626        | Chemical Synapses Can Be Excitatory or Inhibitory  | 684        |
| Glycolipids Are Found on the Surface of All Plasma Membranes   | 628        | The Acetylcholine Receptors at the Neuromuscular Junction Are Transmitter-Gated Cation Channels                            | 684        |
| <i>Summary</i>   | 629        | Transmitter-Gated Ion Channels Are Major Targets for Psychoactive Drugs  | 686        |
| <b>MEMBRANE PROTEINS</b>   | 629        | Neuromuscular Transmission Involves the Sequential Activation of Five Different Sets of Ion Channels                       | 687        |
| Membrane Proteins Can Be Associated with the Lipid Bilayer in Various Ways   | 629        | Single Neurons Are Complex Computation Devices   | 688        |
| Lipid Anchors Control the Membrane Localization of Some Signaling Proteins   | 630        | Neuronal Computation Requires a Combination of at Least Three Kinds of K <sup>+</sup> Channels                             | 689        |
| In Most Transmembrane Proteins the Polypeptide Chain Crosses the Lipid Bilayer in an $\alpha$ -Helical Conformation                              | 631        | Long-Term Potentiation (LTP) in the Mammalian Hippocampus Depends on Ca <sup>2+</sup> Entry Through NMDA-Receptor Channels | 691        |
| Transmembrane $\alpha$ Helices Often Interact with One Another   | 632        | <i>Summary</i>   | 692        |
| Some $\beta$ Barrels Form Large Transmembrane Channels   | 634        | <i>Problems</i>  | 693        |
| Many Membrane Proteins Are Glycosylated  | 635        | <i>References</i>  | 694        |
| Membrane Proteins Can Be Solubilized and Purified in Detergents  | 636        |  |            |
| Bacteriorhodopsin Is a Light-Driven Proton Pump That Traverses the Lipid Bilayer as Seven $\alpha$ Helices                                       | 640        | <b>Chapter 12 Intracellular Compartments and Protein Sorting</b>   | <b>695</b> |
| Membrane Proteins Often Function as Large Complexes  | 642        |  |            |
| Many Membrane Proteins Diffuse in the Plane of the Membrane  | 642        | <b>THE COMPARTMENTALIZATION OF CELLS</b>   | 695        |
| Cells Can Confine Proteins and Lipids to Specific Domains Within a Membrane  | 645        | All Eucaryotic Cells Have the Same Basic Set of Membrane-Enclosed Organelles   | 695        |
| The Cortical Cytoskeleton Gives Membranes Mechanical Strength and Restrict Membrane Protein Diffusion  | 646        | Evolutionary Origins Explain the Topological Relationships of Organelles   | 697        |
| <i>Summary</i>   | 648        | Proteins Can Move Between Compartments in Different Ways   | 699        |
| <i>Problems</i>  | 648        | Signal Sequences Direct Proteins to the Correct Cell Address   | 701        |
| <i>References</i>  | 650        | Most Organelles Cannot Be Constructed <i>De Novo</i> : They Require Information in the Organelle Itself                    | 702        |
|  |            | <i>Summary</i>   | 704        |
| <b>Chapter 11 Membrane Transport of Small Molecules and the Electrical Properties of Membranes</b>   | <b>651</b> | <b>THE TRANSPORT OF MOLECULES BETWEEN THE NUCLEUS AND THE CYTOSOL</b>  | 704        |
|  |            | Nuclear Pore Complexes Perforate the Nuclear Envelope  | 705        |
| <b>PRINCIPLES OF MEMBRANE TRANSPORT</b>  | 651        | Nuclear Localization Signals Direct Nuclear Proteins to the Nucleus  | 705        |
| Protein-Free Lipid Bilayers Are Highly Impermeable to Ions   | 652        | Nuclear Import Receptors Bind to Both Nuclear Localization Signals and NPC proteins  | 707        |
| There Are Two Main Classes of Membrane Transport Proteins: Transporters and Channels   | 652        | Nuclear Export Works Like Nuclear Import, But in Reverse   | 708        |
| Active Transport Is Mediated by Transporters Coupled to an Energy Source   | 653        | The Ran GTPase Imposes Directionality on Transport Through NPCs  | 708        |
| <i>Summary</i>   | 654        | Transport Through NPCs Can Be Regulated by Controlling Access to the Transport Machinery                                   | 709        |
|  |            | During Mitosis the Nuclear Envelope Disassembles   | 710        |
| <b>TRANSPORTERS AND ACTIVE MEMBRANE TRANSPORT</b>  | 654        | <i>Summary</i>   | 712        |
| Active Transport Can Be Driven by Ion Gradients  | 656        | <b>THE TRANSPORT OF PROTEINS INTO MITOCHONDRIA AND CHLOROPLASTS</b>  | 713        |
| Transporters in the Plasma Membrane Regulate Cytosolic pH  | 657        | Translocation into Mitochondria Depends on Signal Sequences and Protein Translocators                                      | 713        |
| An Asymmetric Distribution of Transporters in Epithelial Cells Underlies the Transcellular Transport of Solutes                                  | 658        | Mitochondrial Precursor Proteins Are Imported as Unfolded Polypeptide Chains   | 715        |
| There Are Three Classes of ATP-Driven Pumps  | 659        | ATP Hydrolysis and a Membrane Potential Drive Protein Import into the Matrix Space   | 716        |
| The Ca <sup>2+</sup> Pump Is the Best-Understood P-type ATPase   | 660        | Bacteria and Mitochondria Use Similar Mechanisms to Insert Porins into their Outer Membrane                                | 717        |
| The Plasma Membrane P-type Na <sup>+</sup> -K <sup>+</sup> Pump Establishes the Na <sup>+</sup> Gradient Across the Plasma Membrane              | 661        | Transport into the Inner Mitochondrial Membrane and Intermembrane Space Occurs Via Several Routes                          | 717        |
| ABC Transporters Constitute the Largest Family of Membrane Transport Proteins  | 663        | Two Signal Sequences Direct Proteins to the Thylakoid Membrane in Chloroplasts   | 719        |
| <i>Summary</i>   | 667        | <i>Summary</i>   | 720        |
|  |            | <b>PEROXISOMES</b>   | 721        |
| <b>ION CHANNELS AND THE ELECTRICAL PROPERTIES OF MEMBRANES</b>   | 667        | Peroxisomes Use Molecular Oxygen and Hydrogen Peroxide to Perform Oxidative Reactions                                      | 721        |
| Ion Channels Are Ion-Selective and Fluctuate Between Open and Closed States  | 667        | A Short Signal Sequence Directs the Import of Proteins into Peroxisomes  | 722        |
| The Membrane Potential in Animal Cells Depends Mainly on K <sup>+</sup> Leak Channels and the K <sup>+</sup> Gradient Across the Plasma Membrane | 669        | <i>Summary</i>   | 723        |
| The Resting Potential Decays Only Slowly When the Na <sup>+</sup> -K <sup>+</sup> Pump Is Stopped  | 669        |  |            |
| The Three-Dimensional Structure of a Bacterial K <sup>+</sup> Channel Shows How an Ion Channel Can Work  | 671        |  |            |
| Aquaporins Are Permeable to Water But Impermeable to Ions  | 673        |  |            |
| The Function of a Neuron Depends on Its Elongated Structure  | 675        |  |            |
| Voltage-Gated Cation Channels Generate Action Potentials in Electrically Excitable Cells   | 676        |  |            |
| Myelination Increases the Speed and Efficiency of Action Potential Propagation in Nerve Cells  | 678        |  |            |

|  |            |   |            |
|--|------------|---|------------|
| <b>THE ENDOPLASMIC RETICULUM</b>   | 723        | <b>TRANSPORT FROM THE <i>TRANS</i> GOLGI NETWORK TO LYSOSOMES</b>   | 779        |
| The ER Is Structurally and Functionally Diverse  | 724        | Lysosomes Are the Principal Sites of Intracellular Digestion  | 779        |
| Signal Sequences Were First Discovered in Proteins Imported into the Rough ER  | 726        | Lysosomes Are Heterogeneous   | 780        |
| A Signal-Recognition Particle (SRP) Directs ER Signal Sequences to a Specific Receptor in the Rough ER Membrane                                | 727        | Plant and Fungal Vacuoles Are Remarkably Versatile Lysosomes  | 781        |
| The Polypeptide Chain Passes Through an Aqueous Pore in the Translocator   | 730        | Multiple Pathways Deliver Materials to Lysosomes  | 782        |
| Translocation Across the ER Membrane Does Not Always Require Ongoing Polypeptide Chain Elongation  | 731        | A Mannose 6-Phosphate Receptor Recognizes Lysosomal Proteins in the <i>Trans</i> Golgi Network                                  | 783        |
| In Single-Pass Transmembrane Proteins, a Single Internal ER Signal Sequence Remains in the Lipid Bilayer as a Membrane-Spanning $\alpha$ Helix | 732        | The M6P Receptor Shuttles Between Specific Membranes  | 784        |
| Combinations of Start-Transfer and Stop-Transfer Signals Determine the Topology of Multipass Transmembrane Proteins                            | 734        | A Signal Patch in the Hydrolase Polypeptide Chain Provides the Cue for M6P Addition   | 785        |
| Translocated Polypeptide Chains Fold and Assemble in the Lumen of the Rough ER   | 736        | Defects in the GlcNAc Phosphotransferase Cause a Lysosomal Storage Disease in Humans  | 785        |
| Most Proteins Synthesized in the Rough ER Are Glycosylated by the Addition of a Common <i>N</i> -Linked Oligosaccharide                        | 736        | Some Lysosomes Undergo Exocytosis   | 786        |
| Oligosaccharides Are Used as Tags to Mark the State of Protein Folding   | 738        | <i>Summary</i>  | 786        |
| Improperly Folded Proteins Are Exported from the ER and Degraded in the Cytosol  | 739        | <b>TRANSPORT INTO THE CELL FROM THE PLASMA MEMBRANE: ENDOCYTOSIS</b>  | 787        |
| Misfolded Proteins in the ER Activate an Unfolded Protein Response   | 740        | Specialized Phagocytic Cells Can Ingest Large Particles   | 787        |
| Some Membrane Proteins Acquire a Covalently Attached Glycosylphosphatidylinositol (GPI) Anchor   | 742        | Pinocytic Vesicles Form from Coated Pits in the Plasma Membrane   | 789        |
| The ER Assembles Most Lipid Bilayers   | 743        | Not All Pinocytic Vesicles Are Clathrin-Coated  | 790        |
| <i>Summary</i>   | 745        | Cells Use Receptor-Mediated Endocytosis to Import Selected Extracellular Macromolecules   | 791        |
| <i>Problems</i>  | 746        | Endocytosed Materials That Are Not Retrieved from Endosomes End Up in Lysosomes   | 792        |
| <i>References</i>  | 748        | Specific Proteins Are Retrieved from Early Endosomes and Returned to the Plasma Membrane  | 793        |
| <b>Chapter 13 Intracellular Vesicular Traffic</b>  | <b>749</b> | Multivesicular Bodies Form on the Pathway to Late Endosomes   | 795        |
| <b>THE MOLECULAR MECHANISMS OF MEMBRANE TRANSPORT AND THE MAINTENANCE OF COMPARTMENTAL DIVERSITY</b>   | 750        | Transcytosis Transfers Macromolecules Across Epithelial Cell Sheets   | 797        |
| There Are Various Types of Coated Vesicles   | 751        | Epithelial Cells Have Two Distinct Early Endosomal Compartments but a Common Late Endosomal Compartment                         | 798        |
| The Assembly of a Clathrin Coat Drives Vesicle Formation   | 754        | <i>Summary</i>  | 799        |
| Not All Coats Form Basket-like Structures  | 755        | <b>TRANSPORT FROM THE <i>TRANS</i> GOLGI NETWORK TO THE CELL EXTERIOR: EXOCYTOSIS</b>   | 799        |
| Phosphoinositides Mark Organelles and Membrane Domains   | 757        | Many Proteins and Lipids Seem to Be Carried Automatically from the Golgi Apparatus to the Cell Surface                          | 800        |
| Cytoplasmic Proteins Regulate the Pinching-Off and Uncoating of Coated Vesicles  | 757        | Secretory Vesicles Bud from the <i>Trans</i> Golgi Network  | 801        |
| Monomeric GTPases Control Coat Assembly  | 758        | Proteins Are Often Proteolytically Processed During the Formation of Secretory Vesicles   | 803        |
| Not All Transport Vesicles Are Spherical   | 760        | Secretory Vesicles Wait Near the Plasma Membrane Until Signaled to Release Their Contents                                       | 803        |
| Rab Proteins Guide Vesicle Targeting   | 760        | Regulated Exocytosis Can Be a Localized Response of the Plasma Membrane and Its Underlying Cytoplasm                            | 804        |
| SNAREs Mediate Membrane Fusion   | 762        | Secretory Vesicle Membrane Components Are Quickly Removed from the Plasma Membrane  | 805        |
| Interacting SNAREs Need to Be Pried Apart Before They Can Function Again   | 764        | Some Regulated Exocytosis Events Serve to Enlarge the Plasma Membrane   | 805        |
| Viral Fusion Proteins and SNAREs May Use Similar Fusion Mechanisms   | 764        | Polarized Cells Direct Proteins from the <i>Trans</i> Golgi Network to the Appropriate Domain of the Plasma Membrane            | 805        |
| <i>Summary</i>   | 766        | Different Strategies Guide Membrane Proteins and Lipids Selectively to the Correct Plasma Membrane Domains                      | 806        |
| <b>TRANSPORT FROM THE ER THROUGH THE GOLGI APPARATUS</b>   | 766        | Synaptic Vesicles Can Form Directly from Endocytic Vesicles   | 807        |
| Proteins Leave the ER in COPII-Coated Transport Vesicles   | 767        | <i>Summary</i>  | 809        |
| Only Proteins That Are Properly Folded and Assembled Can Leave the ER  | 767        | <i>Problems</i>   | 810        |
| Vesicular Tubular Clusters Mediate Transport from the ER to the Golgi Apparatus  | 768        | <i>References</i>   | 812        |
| The Retrieval Pathway to the ER Uses Sorting Signals   | 769        | <b>Chapter 14 Energy Conversion: Mitochondria and Chloroplasts</b>  | <b>813</b> |
| Many Proteins Are Selectively Retained in the Compartments in Which They Function  | 771        | <b>THE MITOCHONDRION</b>  | 815        |
| The Golgi Apparatus Consists of an Ordered Series of Compartments  | 771        | The Mitochondrion Contains an Outer Membrane, an Inner Membrane, and Two Internal Compartments                                  | 816        |
| Oligosaccharide Chains Are Processed in the Golgi Apparatus  | 773        | The Citric Acid Cycle Generates High-Energy Electrons   | 817        |
| Proteoglycans Are Assembled in the Golgi Apparatus   | 775        | A Chemiosmotic Process Converts Oxidation Energy into ATP   | 817        |
| What Is the Purpose of Glycosylation?  | 776        | NADH Transfers its Electrons to Oxygen Through Three Large Respiratory Enzyme Complexes   | 819        |
| Transport Through the Golgi Apparatus May Occur by Vesicular Transport or Cisternal Maturation   | 777        | As Electrons Move Along the Respiratory Chain, Energy Is Stored as an Electrochemical Proton Gradient Across the Inner Membrane | 820        |
| Golgi Matrix Proteins Help Organize the Stack  | 778        | The Proton Gradient Drives ATP Synthesis  | 821        |
| <i>Summary</i>   | 779        |   |            |

|   |     |  |            |
|---|-----|--|------------|
| The Proton Gradient Drives Coupled Transport Across the Inner Membrane  | 822 | Mitochondria and Chloroplasts Have Diverse Genomes   | 859        |
| Proton Gradients Produce Most of the Cell's ATP   | 822 | Mitochondria and Chloroplasts Probably Both Evolved from Endosymbiotic Bacteria  | 859        |
| Mitochondria Maintain a High ATP:ADP Ratio in Cells   | 823 | Mitochondria Have a Relaxed Codon Usage and Can Have a Variant Genetic Code  | 861        |
| A Large Negative Value of $\Delta G$ for ATP Hydrolysis Makes ATP Useful to the Cell  | 824 | Animal Mitochondria Contain the Simplest Genetic Systems Known   | 862        |
| ATP Synthase Can Function in Reverse to Hydrolyze ATP and Pump $H^+$  | 826 | Some Organelle Genes Contain Introns   | 863        |
| <i>Summary</i>  | 827 | The Chloroplast Genome of Higher Plants Contains About 120 Genes   | 863        |
| <b>ELECTRON-TRANSPORT CHAINS AND THEIR PROTON PUMPS</b>   | 827 | Mitochondrial Genes Are Inherited by a Non-Mendelian Mechanism   | 864        |
| Protons Are Unusually Easy to Move  | 827 | Organelle Genes Are Maternally Inherited in Many Organisms   | 866        |
| The Redox Potential Is a Measure of Electron Affinities   | 828 | Petite Mutants in Yeasts Demonstrate the Overwhelming Importance of the Cell Nucleus for Mitochondrial Biogenesis                    | 866        |
| Electron Transfers Release Large Amounts of Energy  | 829 | Mitochondria and Plastids Contain Tissue-Specific Proteins that Are Encoded in the Cell Nucleus                                      | 867        |
| Spectroscopic Methods Identified Many Electron Carriers in the Respiratory Chain  | 829 | Mitochondria Import Most of Their Lipids; Chloroplasts Make Most of Theirs   | 867        |
| The Respiratory Chain Includes Three Large Enzyme Complexes Embedded in the Inner Membrane                                  | 831 | Mitochondria May Contribute to the Aging of Cells and Organisms  | 868        |
| An Iron-Copper Center in Cytochrome Oxidase Catalyzes Efficient $O_2$ Reduction   | 832 | Why Do Mitochondria and Chloroplasts Have Their Own Genetic Systems?   | 868        |
| Electron Transfers in the Inner Mitochondrial Membrane Are Mediated by Electron Tunneling during Random Collisions          | 834 | <i>Summary</i>   | 870        |
| A Large Drop in Redox Potential Across Each of the Three Respiratory Enzyme Complexes Provides the Energy for $H^+$ Pumping | 835 | <b>THE EVOLUTION OF ELECTRON-TRANSPORT CHAINS</b>  | 870        |
| The $H^+$ Pumping Occurs by Distinct Mechanisms in the Three Major Enzyme Complexes   | 835 | The Earliest Cells Probably Used Fermentation to Produce ATP   | 870        |
| $H^+$ Ionophores Uncouple Electron Transport from ATP Synthesis   | 836 | Electron-Transport Chains Enabled Anaerobic Bacteria to Use Nonfermentable Molecules as Their Major Source of Energy                 | 871        |
| Respiratory Control Normally Restrains Electron Flow Through the Chain  | 837 | By Providing an Inexhaustible Source of Reducing Power, Photosynthetic Bacteria Overcame a Major Evolutionary Obstacle               | 872        |
| Natural Uncouplers Convert the Mitochondria in Brown Fat into Heat-Generating Machines                                      | 838 | The Photosynthetic Electron-Transport Chains of Cyanobacteria Produced Atmospheric Oxygen and Permitted New Life-Forms               | 873        |
| The Mitochondrion Plays Many Critical Roles in Cell Metabolism  | 838 | <i>Summary</i>   | 875        |
| Bacteria Also Exploit Chemiosmotic Mechanisms to Harness Energy   | 839 | <i>Problems</i>  | 877        |
| <i>Summary</i>  | 840 | <i>References</i>  | 878        |
| <b>CHLOROPLASTS AND PHOTOSYNTHESIS</b>  | 840 | <b>Chapter 15 Mechanisms of Cell Communication</b>   | <b>879</b> |
| The Chloroplast Is One Member of the Plastid Family of Organelles   | 841 | <b>GENERAL PRINCIPLES OF CELL COMMUNICATION</b>  | 879        |
| Chloroplasts Resemble Mitochondria But Have an Extra Compartment  | 842 | Extracellular Signal Molecules Bind to Specific Receptors  | 880        |
| Chloroplasts Capture Energy from Sunlight and Use It to Fix Carbon  | 843 | Extracellular Signal Molecules Can Act Over Either Short or Long Distances   | 881        |
| Carbon Fixation Is Catalyzed by Ribulose Bisphosphate Carboxylase   | 844 | Gap Junctions Allow Neighboring Cells to Share Signaling Information   | 884        |
| Each $CO_2$ Molecule That Is Fixed Consumes Three Molecules of ATP and Two Molecules of NADPH                               | 845 | Each Cell Is Programmed to Respond to Specific Combinations of Extracellular Signal Molecules  | 884        |
| Carbon Fixation in Some Plants Is Compartmentalized to Facilitate Growth at Low $CO_2$ Concentrations                       | 846 | Different Types of Cells Usually Respond Differently to the Same Extracellular Signal Molecule                                       | 885        |
| Photosynthesis Depends on the Photochemistry of Chlorophyll Molecules   | 847 | The Fate of Some Developing Cells Depends on Their Position in Morphogen Gradients   | 886        |
| A Photochemical Reaction Center Plus an Antenna Complex Form a Photosystem  | 848 | A Cell Can Alter the Concentration of an Intracellular Molecule Quickly Only If the Lifetime of the Molecule Is Short                | 886        |
| In a Reaction Center, Light Energy Captured by Chlorophyll Creates a Strong Electron Donor from a Weak One                  | 849 | Nitric Oxide Gas Signals by Directly Regulating the Activity of Specific Proteins Inside the Target Cell                             | 887        |
| Noncyclic Photophosphorylation Produces Both NADPH and ATP  | 850 | Nuclear Receptors Are Ligand-Modulated Gene Regulatory Proteins  | 889        |
| Chloroplasts Can Make ATP by Cyclic Photophosphorylation Without Making NADPH   | 853 | The Three Largest Classes of Cell-Surface Receptor Proteins Are Ion-Channel-Coupled, G-Protein-Coupled, and Enzyme-Coupled Receptors | 891        |
| Photosystems I and II Have Related Structures, and Also Resemble Bacterial Photosystems                                     | 853 | Most Activated Cell-Surface Receptors Relay Signals Via Small Molecules and a Network of Intracellular Signaling Proteins            | 893        |
| The Proton-Motive Force Is the Same in Mitochondria and Chloroplasts  | 853 | Many Intracellular Signaling Proteins Function as Molecular Switches That Are Activated by Phosphorylation or GTP Binding            | 895        |
| Carrier Proteins in the Chloroplast Inner Membrane Control Metabolite Exchange with the Cytosol                             | 854 | Intracellular Signaling Complexes Enhance the Speed, Efficiency, and Specificity of the Response                                     | 897        |
| Chloroplasts Also Perform Other Crucial Biosyntheses  | 855 | Modular Interaction Domains Mediate Interactions Between Intracellular Signaling Proteins  | 897        |
| <i>Summary</i>  | 855 | Cells Can Use Multiple Mechanisms to Respond Abruptly to a Gradually Increasing Concentration of an Extracellular Signal             | 899        |
| <b>THE GENETIC SYSTEMS OF MITOCHONDRIA AND PLASTIDS</b>   | 855 | Intracellular Signaling Networks Usually Make Use of Feedback Loops  | 901        |
| Mitochondria and Chloroplasts Contain Complete Genetic Systems  | 856 | Cells Can Adjust Their Sensitivity to a Signal   | 902        |
| Organelle Growth and Division Determine the Number of Mitochondria and Plastids in a Cell                                   | 857 | <i>Summary</i>   | 903        |

|  |     |  |            |
|--|-----|--|------------|
| SIGNALING THROUGH G-PROTEIN-COUPLED CELL-SURFACE RECEPTORS (GPCRS) AND SMALL INTRACELLULAR MEDIATORS   | 904 | Ethylene Blocks the Degradation of Specific Gene Regulatory Proteins in the Nucleus                          | 957        |
| Trimeric G Proteins Relay Signals from GPCRs   | 905 | Regulated Positioning of Auxin Transporters Patterns Plant Growth  | 959        |
| Some G Proteins Regulate the Production of Cyclic AMP  | 905 | Phytochromes Detect Red Light, and Cryptochromes Detect Blue Light   | 960        |
| Cyclic-AMP-Dependent Protein Kinase (PKA) Mediates Most of the Effects of Cyclic AMP   | 908 | <i>Summary</i>   | 961        |
| Some G Proteins Activate An Inositol Phospholipid Signaling Pathway by Activating Phospholipase C- $\beta$                                     | 909 | <i>Problems</i>  | 962        |
| Ca <sup>2+</sup> Functions as a Ubiquitous Intracellular Mediator  | 912 | <i>References</i>  | 964        |
| The Frequency of Ca <sup>2+</sup> Oscillations Influences a Cell's Response  | 912 | <b>Chapter 16 The Cytoskeleton</b>   | <b>965</b> |
| Ca <sup>2+</sup> /Calmodulin-Dependent Protein Kinases (CaM-Kinases) Mediate Many of the Responses to Ca <sup>2+</sup> Signals in Animal Cells | 914 | THE SELF-ASSEMBLY AND DYNAMIC STRUCTURE OF CYTOSKELETAL FILAMENTS  | 965        |
| Some G Proteins Directly Regulate Ion Channels   | 916 | Cytoskeletal Filaments Are Dynamic and Adaptable   | 966        |
| Smell and Vision Depend on GPCRs That Regulate Cyclic-Nucleotide-Gated Ion Channels  | 917 | The Cytoskeleton Can Also Form Stable Structures   | 969        |
| Intracellular Mediators and Enzymatic Cascades Amplify Extracellular Signals   | 919 | Each Type of Cytoskeletal Filament Is Constructed from Smaller Protein Subunits                              | 970        |
| GPCR Desensitization Depends on Receptor Phosphorylation   | 920 | Filaments Formed from Multiple Protofilaments Have Advantageous Properties                                   | 971        |
| <i>Summary</i>   | 921 | Nucleation Is the Rate-Limiting Step in the Formation of a Cytoskeletal Polymer                              | 973        |
| SIGNALING THROUGH ENZYME-COUPLED CELL-SURFACE RECEPTORS  | 921 | The Tubulin and Actin Subunits Assemble Head-to-Tail to Create Polar Filaments                               | 973        |
| Activated Receptor Tyrosine Kinases (RTKs) Phosphorylate Themselves  | 922 | Microtubules and Actin Filaments Have Two Distinct Ends That Grow at Different Rates                         | 975        |
| Phosphorylated Tyrosines on RTKs Serve as Docking Sites for Intracellular Signaling Proteins   | 923 | Filament Treadmilling and Dynamic Instability Are Consequences of Nucleotide Hydrolysis by Tubulin and Actin | 976        |
| Proteins with SH2 Domains Bind to Phosphorylated Tyrosines   | 924 | Treadmilling and Dynamic Instability Aid Rapid Cytoskeletal Rearrangement                                    | 980        |
| Ras Belongs to a Large Superfamily of Monomeric GTPases  | 926 | Tubulin and Actin Have Been Highly Conserved During Eucaryotic Evolution                                     | 982        |
| RTKs Activate Ras Via Adaptors and GEFs: Evidence from the Developing <i>Drosophila</i> Eye  | 927 | Intermediate Filament Structure Depends on The Lateral Bundling and Twisting of Coiled Coils                 | 983        |
| Ras Activates a MAP Kinase Signaling Module  | 928 | Intermediate Filaments Impart Mechanical Stability to Animal Cells   | 985        |
| Scaffold Proteins Help Prevent Cross-Talk Between Parallel MAP Kinase Modules  | 930 | Drugs Can Alter Filament Polymerization  | 987        |
| Rho Family GTPases Functionally Couple Cell-Surface Receptors to the Cytoskeleton  | 931 | Bacterial Cell Organization and Cell Division Depend on Homologs of the Eucaryotic Cytoskeleton              | 989        |
| PI 3-Kinase Produces Lipid Docking Sites in the Plasma Membrane  | 932 | <i>Summary</i>   | 991        |
| The PI-3-Kinase-Akt Signaling Pathway Stimulates Animal Cells to Survive and Grow  | 934 | HOW CELLS REGULATE THEIR CYTOSKELETAL FILAMENTS  | 992        |
| The Downstream Signaling Pathways Activated By RTKs and GPCRs Overlap  | 935 | A Protein Complex Containing $\gamma$ -Tubulin Nucleates Microtubules  | 992        |
| Tyrosine-Kinase-Associated Receptors Depend on Cytoplasmic Tyrosine Kinases  | 935 | Microtubules Emanate from the Centrosome in Animal Cells   | 992        |
| Cytokine Receptors Activate the JAK-STAT Signaling Pathway, Providing a Fast Track to the Nucleus  | 937 | Actin Filaments Are Often Nucleated at the Plasma Membrane   | 996        |
| Protein Tyrosine Phosphatases Reverse Tyrosine Phosphorylations  | 938 | The Mechanism of Nucleation Influences Large-Scale Filament Organization                                     | 998        |
| Signal Proteins of the TGF $\beta$ Superfamily Act Through Receptor Serine/Threonine Kinases and Smads   | 939 | Proteins That Bind to the Free Subunits Modify Filament Elongation   | 999        |
| Serine/Threonine and Tyrosine Protein Kinases Are Structurally Related   | 941 | Severing Proteins Regulate the Length and Kinetic Behavior of Actin Filaments and Microtubules               | 1000       |
| Bacterial Chemotaxis Depends on a Two-Component Signaling Pathway Activated by Histidine-Kinase-Associated Receptors                           | 941 | Proteins That Bind Along the Sides of Filaments Can Either Stabilize or Destabilize Them                     | 1001       |
| Receptor Methylation Is Responsible for Adaptation in Bacterial Chemotaxis   | 943 | Proteins That Interact with Filament Ends Can Dramatically Change Filament Dynamics                          | 1002       |
| <i>Summary</i>   | 944 | Different Kinds of Proteins Alter the Properties of Rapidly Growing Microtubule Ends                         | 1003       |
| SIGNALING PATHWAYS DEPENDENT ON REGULATED PROTEOLYSIS OF LATENT GENE REGULATORY PROTEINS   | 946 | Filaments Are Organized into Higher-Order Structures in Cells  | 1005       |
| The Receptor Protein Notch Is a Latent Gene Regulatory Protein   | 946 | Intermediate Filaments Are Cross-Linked and Bundled Into Strong Arrays                                       | 1005       |
| Wnt Proteins Bind to Frizzled Receptors and Inhibit the Degradation of $\beta$ -Catenin  | 948 | Cross-Linking Proteins with Distinct Properties Organize Different Assemblies of Actin Filaments             | 1006       |
| Hedgehog Proteins Bind to Patched Relieving Its Inhibition of Smoothened   | 950 | Filamin and Spectrin Form Actin Filament Webs  | 1008       |
| Many Stressful and Inflammatory Stimuli Act Through an NF $\kappa$ B-Dependent Signaling Pathway   | 952 | Cytoskeletal Elements Make Many Attachments to Membrane  | 1009       |
| <i>Summary</i>   | 954 | <i>Summary</i>   | 1010       |
| SIGNALING IN PLANTS  | 955 | MOLECULAR MOTORS   | 1010       |
| Multicellularity and Cell Communication Evolved Independently in Plants and Animals  | 955 | Actin-Based Motor Proteins Are Members of the Myosin Superfamily   | 1011       |
| Receptor Serine/Threonine Kinases Are the Largest Class of Cell-Surface Receptors in Plants  | 956 | There Are Two Types of Microtubule Motor Proteins: Kinesins and Dyneins                                      | 1014       |
|  |     | The Structural Similarity of Myosin and Kinesin Indicates a Common Evolutionary Origin                       | 1015       |
|  |     | Motor Proteins Generate Force by Coupling ATP Hydrolysis to Conformational Changes                           | 1016       |

|  |             |   |             |
|--|-------------|---|-------------|
| Motor Protein Kinetics Are Adapted to Cell Functions   | 1020        | Microtubule-Dependent Motor Proteins Govern Spindle Assembly and Function                   | 1077        |
| Motor Proteins Mediate the Intracellular Transport of Membrane-Enclosed Organelles                               | 1021        | Two Mechanisms Collaborate in the Assembly of a Bipolar Mitotic Spindle                     | 1077        |
| The Cytoskeleton Localizes Specific RNA Molecules  | 1022        | Centrosome Duplication Occurs Early in the Cell Cycle                                       | 1078        |
| Cells Regulate Motor Protein Function  | 1023        | M-Cdk Initiates Spindle Assembly in Prophase  | 1078        |
| <i>Summary</i>   | 1025        | The Completion of Spindle Assembly in Animal Cells Requires Nuclear Envelope Breakdown      | 1079        |
| <b>THE CYTOSKELETON AND CELL BEHAVIOR</b>  | 1025        | Microtubule Instability Increases Greatly in Mitosis  | 1080        |
| Sliding of Myosin II and Actin Filaments Causes Muscles to Contract  | 1026        | Mitotic Chromosomes Promote Bipolar Spindle Assembly  | 1081        |
| A Sudden Rise in Cytosolic Ca <sup>2+</sup> Concentration Initiates Muscle Contraction                           | 1028        | Kinetochores Attach Sister Chromatids to the Spindle  | 1082        |
| Heart Muscle Is a Precisely Engineered Machine   | 1031        | Bi-Orientation Is Achieved by Trial and Error   | 1083        |
| Cilia and Flagella Are Motile Structures Built from Microtubules and Dyneins                                     | 1031        | Multiple Forces Move Chromosomes on the Spindle   | 1085        |
| Construction of the Mitotic Spindle Requires Microtubule Dynamics and the Interactions of Many Motor Proteins    | 1034        | The APC/C Triggers Sister-Chromatid Separation and the Completion of Mitosis                | 1087        |
| Many Cells Can Crawl Across A Solid Substratum   | 1036        | Unattached Chromosomes Block Sister-Chromatid Separation: The Spindle Assembly Checkpoint   | 1088        |
| Actin Polymerization Drives Plasma Membrane Protrusion   | 1037        | Chromosomes Segregate in Anaphase A and B   | 1089        |
| Cell Adhesion and Traction Allow Cells to Pull Themselves Forward  | 1040        | Segregated Chromosomes Are Packaged in Daughter Nuclei at Telophase                         | 1090        |
| Members of the Rho Protein Family Cause Major Rearrangements of the Actin Cytoskeleton                           | 1041        | Meiosis Is a Special Form of Nuclear Division Involved in Sexual Reproduction               | 1090        |
| Extracellular Signals Can Activate the Three Rho Protein Family Members  | 1043        | <i>Summary</i>  | 1092        |
| External Signals Can Dictate the Direction of Cell Migration   | 1045        | <b>CYTOKINESIS</b>  | 1092        |
| Communication Between the Microtubule and Actin Cytoskeletons Coordinates Whole-Cell Polarization and Locomotion | 1046        | Actin and Myosin II in the Contractile Ring Generate the Force for Cytokinesis              | 1093        |
| The Complex Morphological Specialization of Neurons Depends on the Cytoskeleton                                  | 1047        | Local Activation of RhoA Triggers Assembly and Contraction of the Contractile Ring          | 1094        |
| <i>Summary</i>   | 1050        | The Microtubules of the Mitotic Spindle Determine the Plane of Animal Cell Division         | 1095        |
| <i>Problems</i>  | 1050        | The Phragmoplast Guides Cytokinesis in Higher Plants  | 1097        |
| <i>References</i>  | 1052        | Membrane-Enclosed Organelles Must Be Distributed to Daughter Cells During Cytokinesis       | 1098        |
| <b>Chapter 17 The Cell Cycle</b>   | <b>1053</b> | Some Cells Reposition Their Spindle to Divide Asymmetrically                                | 1099        |
| <b>OVERVIEW OF THE CELL CYCLE</b>  | 1054        | Mitosis Can Occur Without Cytokinesis   | 1099        |
| The Eucaryotic Cell Cycle Is Divided into Four Phases  | 1054        | The G <sub>1</sub> Phase Is a Stable State of Cdk Inactivity                                | 1100        |
| Cell-Cycle Control Is Similar in All Eucaryotes  | 1056        | <i>Summary</i>  | 1101        |
| Cell-Cycle Control Can Be Dissected Genetically by Analysis of Yeast Mutants                                     | 1056        | <b>CONTROL OF CELL DIVISION AND CELL GROWTH</b>   | 1101        |
| Cell-Cycle Control Can Be Analyzed Biochemically in Animal Embryos   | 1057        | Mitogens Stimulate Cell Division  | 1102        |
| Cell-Cycle Control Can Be Studied in Cultured Mammalian Cells  | 1059        | Cells Can Delay Division by Entering a Specialized Nondividing State                        | 1103        |
| Cell-Cycle Progression Can Be Studied in Various Ways  | 1059        | Mitogens Stimulate G <sub>1</sub> -Cdk and G <sub>1</sub> /S-Cdk Activities                 | 1103        |
| <i>Summary</i>   | 1060        | DNA Damage Blocks Cell Division: The DNA Damage Response                                    | 1105        |
| <b>THE CELL-CYCLE CONTROL SYSTEM</b>   | 1060        | Many Human Cells Have a Built-In Limitation on the Number of Times They Can Divide          | 1007        |
| The Cell-Cycle Control System Triggers the Major Events of the Cell Cycle  | 1060        | Abnormal Proliferation Signals Cause Cell-Cycle Arrest or Apoptosis, Except in Cancer Cells | 1107        |
| The Cell-Cycle Control System Depends on Cyclically Activated Cyclin-Dependent Protein Kinases (Cdks)            | 1062        | Organism and Organ Growth Depend on Cell Growth   | 1108        |
| Inhibitory Phosphorylation and Cdk Inhibitory Proteins (CKIs) Can Suppress Cdk Activity                          | 1063        | Proliferating Cells Usually Coordinate Their Growth and Division                            | 1108        |
| The Cell-Cycle Control System Depends on Cyclical Proteolysis  | 1064        | Neighboring Cells Compete for Extracellular Signal Proteins                                 | 1110        |
| Cell-Cycle Control Also Depends on Transcriptional Regulation  | 1065        | Animals Control Total Cell Mass by Unknown Mechanisms                                       | 1111        |
| The Cell-Cycle Control System Functions as a Network of Biochemical Switches                                     | 1065        | <i>Summary</i>  | 1112        |
| <i>Summary</i>   | 1067        | <i>Problems</i>   | 1112        |
| <b>S PHASE</b>   | 1067        | <i>References</i>   | 1113        |
| S-Cdk Initiates DNA Replication Once Per Cycle   | 1067        | <b>Chapter 18 Apoptosis</b>   | <b>1115</b> |
| Chromosome Duplication Requires Duplication of Chromatin Structure   | 1069        | Programmed Cell Death Eliminates Unwanted Cells   | 1115        |
| Cohesins Help Hold Sister Chromatids Together  | 1070        | Apoptotic Cells Are Biochemically Recognizable  | 1117        |
| <i>Summary</i>   | 1071        | Apoptosis Depends on an Intracellular Proteolytic Cascade That Is Mediated by Caspases      | 1118        |
| <b>MITOSIS</b>   | 1071        | Cell-Surface Death Receptors Activate the Extrinsic Pathway of Apoptosis                    | 1120        |
| M-Cdk Drives Entry Into Mitosis  | 1071        | The Intrinsic Pathway of Apoptosis Depends on Mitochondria                                  | 1121        |
| Dephosphorylation Activates M-Cdk at the Onset of Mitosis  | 1074        | Bcl2 Proteins Regulate the Intrinsic Pathway of Apoptosis                                   | 1121        |
| Condensin Helps Configure Duplicated Chromosomes for Separation  | 1075        | IAPs Inhibit Caspases   | 1124        |
| The Mitotic Spindle Is a Microtubule-Based Machine   | 1075        | Extracellular Survival Factors Inhibit Apoptosis in Various Ways                            | 1126        |
|  |             | Either Excessive or Insufficient Apoptosis Can Contribute to Disease                        | 1127        |
|  |             | <i>Summary</i>  | 1128        |
|  |             | <i>Problems</i>   | 1128        |
|  |             | <i>References</i>   | 1129        |

## Chapter 19 Cell Junctions, Cell Adhesion, and the Extracellular Matrix

|  |             |
|--|-------------|
| <b>CADHERINS AND CELL-CELL ADHESION</b>  | <b>1133</b> |
| Cadherins Mediate Ca <sup>2+</sup> -Dependent Cell-Cell Adhesion in All Animals  | 1135        |
| The Cadherin Superfamily in Vertebrates Includes Hundreds of Different Proteins, Including Many with Signaling Functions | 1136        |
| Cadherins Mediate Homophilic Adhesion  | 1137        |
| Selective Cell-Cell Adhesion Enables Dissociated Vertebrate Cells to Reassemble into Organized Tissues                   | 1139        |
| Cadherins Control the Selective Assortment of Cells  | 1140        |
| Twist Regulates Epithelial-Mesenchymal Transitions   | 1141        |
| Catenins Link Classical Cadherins to the Actin Cytoskeleton  | 1142        |
| Adherens Junctions Coordinate the Actin-Based Motility of Adjacent Cells   | 1142        |
| Desmosome Junctions Give Epithelia Mechanical Strength   | 1143        |
| Cell-Cell Junctions Send Signals into the Cell Interior  | 1145        |
| Selectins Mediate Transient Cell-Cell Adhesions in the Bloodstream   | 1145        |
| Members of the Immunoglobulin Superfamily of Proteins Mediate Ca <sup>2+</sup> -Independent Cell-Cell Adhesion           | 1146        |
| Many Types of Cell Adhesion Molecules Act in Parallel to Create a Synapse  | 1147        |
| Scaffold Proteins Organize Junctional Complexes  | 1148        |
| <i>Summary</i>   | 1149        |
| <b>TIGHT JUNCTIONS AND THE ORGANIZATION OF EPITHELIA</b>   | <b>1150</b> |
| Tight Junctions Form a Seal Between Cells and a Fence Between Membrane Domains   | 1150        |
| Scaffold Proteins in Junctional Complexes Play a Key Part in the Control of Cell Proliferation                           | 1153        |
| Cell-Cell Junctions and the Basal Lamina Govern Apico-Basal Polarity in Epithelia  | 1155        |
| A Separate Signaling System Controls Planar Cell Polarity  | 1157        |
| <i>Summary</i>   | 1158        |
| <b>PASSAGEWAYS FROM CELL TO CELL: GAP JUNCTIONS AND PLASMODESMATA</b>  | <b>1158</b> |
| Gap Junctions Couple Cells Both Electrically and Metabolically   | 1158        |
| A Gap-Junction Connexon Is Made Up of Six Transmembrane Connexin Subunits  | 1159        |
| Gap Junctions Have Diverse Functions   | 1161        |
| Cells Can Regulate the Permeability of Their Gap Junctions   | 1161        |
| In Plants, Plasmodesmata Perform Many of the Same Functions as Gap Junctions   | 1162        |
| <i>Summary</i>   | 1163        |
| <b>THE BASAL LAMINA</b>  | <b>1164</b> |
| Basal Laminae Underlie All Epithelia and Surround Some Nonepithelial Cell Types  | 1164        |
| Laminin Is a Primary Component of the Basal Lamina   | 1165        |
| Type IV Collagen Gives the Basal Lamina Tensile Strength   | 1166        |
| Basal Laminae Have Diverse Functions   | 1167        |
| <i>Summary</i>   | 1169        |
| <b>INTEGRINS AND CELL-MATRIX ADHESION</b>  | <b>1169</b> |
| Integrins Are Transmembrane Heterodimers That Link to the Cytoskeleton   | 1170        |
| Integrins Can Switch Between an Active and an Inactive Conformation  | 1170        |
| Integrin Defects Are Responsible for Many Different Genetic Diseases   | 1172        |
| Integrins Cluster to Form Strong Adhesions   | 1174        |
| Extracellular Matrix Attachments Act Through Integrins to Control Cell Proliferation and Survival                        | 1175        |
| Integrins Recruit Intracellular Signaling Proteins at Sites of Cell-Substratum Adhesion                                  | 1176        |
| Integrins Can Produce Localized Intracellular Effects  | 1177        |
| <i>Summary</i>   | 1178        |

|  |             |
|--|-------------|
| <b>THE EXTRACELLULAR MATRIX OF ANIMAL CONNECTIVE TISSUES</b>   | <b>1178</b> |
| The Extracellular Matrix Is Made and Oriented by the Cells Within It   | 1179        |
| Glycosaminoglycan (GAG) Chains Occupy Large Amounts of Space and Form Hydrated Gels                          | 1179        |
| Hyaluronan Acts as a Space Filler and a Facilitator of Cell Migration During Tissue Morphogenesis and Repair | 1180        |
| Proteoglycans Are Composed of GAG Chains Covalently Linked to a Core Protein                                 | 1181        |
| Proteoglycans Can Regulate the Activities of Secreted Proteins   | 1182        |
| Cell-Surface Proteoglycans Act as Co-Receptors   | 1183        |
| Collagens Are the Major Proteins of the Extracellular Matrix   | 1184        |
| Collagen Chains Undergo a Series of Post-Translational Modifications   | 1186        |
| Propeptides Are Clipped Off Procollagen After Its Secretion to Allow Assembly of Fibrils                     | 1187        |
| Secreted Fibril-Associated Collagens Help Organize the Fibrils   | 1187        |
| Cells Help Organize the Collagen Fibrils They Secrete by Exerting Tension on the Matrix                      | 1189        |
| Elastin Gives Tissues Their Elasticity   | 1189        |
| Fibronectin Is an Extracellular Protein That Helps Cells Attach to the Matrix                                | 1191        |
| Tension Exerted by Cells Regulates Assembly of Fibronectin Fibrils   | 1191        |
| Fibronectin Binds to Integrins Through an RGD Motif  | 1193        |
| Cells Have to Be Able to Degrade Matrix, as Well as Make it  | 1193        |
| Matrix Degradation Is Localized to the Vicinity of Cells   | 1194        |
| <i>Summary</i>   | 1195        |

|  |             |
|--|-------------|
| <b>THE PLANT CELL WALL</b>   | <b>1195</b> |
| The Composition of the Cell Wall Depends on the Cell Type  | 1195        |
| The Tensile Strength of the Cell Wall Allows Plant Cells to Develop Turgor Pressure                            | 1197        |
| The Primary Cell Wall Is Built from Cellulose Microfibrils Interwoven with a Network of Pectic Polysaccharides | 1197        |
| Oriented Cell-Wall Deposition Controls Plant Cell Growth   | 1199        |
| Microtubules Orient Cell-Wall Deposition   | 1200        |
| <i>Summary</i>   | 1202        |
| <i>Problems</i>  | 1202        |
| <i>References</i>  | 1204        |

## Chapter 20 Cancer 1205

|   |             |
|---|-------------|
| <b>CANCER AS A MICROEVOLUTIONARY PROCESS</b>  | <b>1205</b> |
| Cancer Cells Reproduce Without Restraint and Colonize Other Tissues   | 1206        |
| Most Cancers Derive from a Single Abnormal Cell   | 1207        |
| Cancer Cells Contain Somatic Mutations  | 1208        |
| A Single Mutation Is Not Enough to Cause Cancer   | 1209        |
| Cancers Develop Gradually from Increasingly Aberrant Cells  | 1210        |
| Cervical Cancers Are Prevented by Early Detection   | 1211        |
| Tumor Progression Involves Successive Rounds of Random Inherited Change Followed by Natural Selection             | 1212        |
| The Epigenetic Changes That Accumulate in Cancer Cells Involve Inherited Chromatin Structures and DNA Methylation | 1213        |
| Human Cancer Cells Are Genetically Unstable   | 1214        |
| Cancerous Growth Often Depends on Defective Control of Cell Death, Cell Differentiation, or Both                  | 1215        |
| Cancer Cells Are Usually Altered in Their Responses to DNA Damage and Other Forms of Stress                       | 1216        |
| Human Cancer Cells Escape a Built-In Limit to Cell Proliferation  | 1217        |
| A Small Population of Cancer Stem Cells Maintains Many Tumors   | 1217        |
| How Do Cancer Stem Cells Arise?   | 1218        |
| To Metastasize, Malignant Cancer Cells Must Survive and Proliferate in a Foreign Environment                      | 1220        |
| Tumors Induce Angiogenesis  | 1220        |
| The Tumor Microenvironment Influences Cancer Development  | 1222        |
| Many Properties Typically Contribute to Cancerous Growth  | 1223        |
| <i>Summary</i>  | 1223        |

|   |      |  |             |
|---|------|--|-------------|
| THE PREVENTABLE CAUSES OF CANCER  | 1224 | There Is Still Much More to Do   | 1264        |
| Many, But Not All, Cancer-Causing Agents Damage DNA   | 1225 | <i>Summary</i>   | 1265        |
| Tumor Initiators Damage DNA; Tumor Promoters Do Not   | 1226 | <i>Problems</i>  | 1265        |
| Viruses and Other Infections Contribute to a Significant Proportion of Human Cancers  | 1227 | <i>References</i>  | 1267        |
| Identification of Carcinogens Reveals Ways to Avoid Cancer  | 1229 | <b>Chapters 21–25 available on Media DVD-ROM</b>                                 |             |
| <i>Summary</i>  | 1230 | <b>Chapter 21 Sexual Reproduction: Meiosis, Germ Cells, and Fertilization</b>    | <b>1269</b> |
| FINDING THE CANCER-CRITICAL GENES   | 1230 | OVERVIEW OF SEXUAL REPRODUCTION  | 1269        |
| The Identification of Gain-of-Function and Loss-of-Function Mutations Requires Different Methods                                | 1231 | The Haploid Phase in Higher Eucaryotes Is Brief                                  | 1269        |
| Retroviruses Can Act as Vectors for Oncogenes That Alter Cell Behavior  | 1232 | Meiosis Creates Genetic Diversity  | 1271        |
| Different Searches for Oncogenes Have Converged on the Same Gene—Ras  | 1233 | Sexual Reproduction Gives Organisms a Competitive Advantage                      | 1271        |
| Studies of Rare Hereditary Cancer Syndromes First Identified Tumor Suppressor Genes   | 1234 | <i>Summary</i>   | 1272        |
| Tumor Suppressor Genes Can Also Be Identified from Studies of Tumors  | 1235 | MEIOSIS  | 1272        |
| Both Genetic and Epigenetic Mechanisms Can Inactivate Tumor Suppressor Genes  | 1235 | Gametes Are Produced by Two Meiotic Cell Divisions                               | 1272        |
| Genes Mutated in Cancer Can Be Made Overactive in Many Ways   | 1237 | Duplicated Homologs (and Sex Chromosomes) Pair During Early Prophase I           | 1274        |
| The Hunt for Cancer-Critical Genes Continues  | 1239 | Homolog Pairing Culminates in the Formation of a Synaptonemal Complex            | 1275        |
| <i>Summary</i>  | 1240 | Homolog Segregation Depends on Meiosis-Specific, Kinetochore-Associated Proteins | 1276        |
| THE MOLECULAR BASIS OF CANCER-CELL BEHAVIOR   | 1240 | Meiosis Frequently Goes Wrong  | 1278        |
| Studies of Both Developing Embryos and Genetically Engineered Mice Have Helped to Uncover the Function of Cancer-Critical Genes | 1241 | Crossing-Over Enhances Genetic Reassortment                                      | 1279        |
| Many Cancer-Critical Genes Regulate Cell Proliferation  | 1242 | Crossing-Over Is Highly Regulated  | 1280        |
| Distinct Pathways May Mediate the Disregulation of Cell-Cycle Progression and the Disregulation of Cell Growth in Cancer Cells  | 1244 | Meiosis Is Regulated Differently in Male and Female Mammals                      | 1280        |
| Mutations in Genes That Regulate Apoptosis Allow Cancer Cells to Survive When They Should Not                                   | 1245 | <i>Summary</i>   | 1281        |
| Mutations in the <i>p53</i> Gene Allow Many Cancer Cells to Survive and Proliferate Despite DNA Damage                          | 1246 | PRIMORDIAL GERM CELLS AND SEX DETERMINATION IN MAMMALS                           | 1282        |
| DNA Tumor Viruses Block the Action of Key Tumor Suppressor Proteins   | 1247 | Signals from Neighbors Specify PGCs in Mammalian Embryos                         | 1282        |
| The Changes in Tumor Cells That Lead to Metastasis Are Still Largely a Mystery  | 1249 | PGCs Migrate into the Developing Gonads  | 1283        |
| Colorectal Cancers Evolve Slowly Via a Succession of Visible Changes  | 1250 | The <i>Sry</i> Gene Directs the Developing Mammalian Gonad to Become a Testis    | 1283        |
| A Few Key Genetic Lesions Are Common to a Large Fraction of Colorectal Cancers  | 1251 | Many Aspects of Sexual Reproduction Vary Greatly between Animal Species          | 1285        |
| Some Colorectal Cancers Have Defects in DNA Mismatch Repair   | 1254 | <i>Summary</i>   | 1286        |
| The Steps of Tumor Progression Can Often Be Correlated with Specific Mutations  | 1254 | EGGS   | 1287        |
| Each Case of Cancer Is Characterized by Its Own Array of Genetic Lesions  | 1256 | An Egg Is Highly Specialized for Independent Development                         | 1287        |
| <i>Summary</i>  | 1256 | Eggs Develop in Stages   | 1288        |
| CANCER TREATMENT: PRESENT AND FUTURE  | 1256 | Oocytes Use Special Mechanisms to Grow to Their Large Size                       | 1290        |
| The Search for Cancer Cures Is Difficult but Not Hopeless   | 1257 | Most Human Oocytes Die Without Maturing  | 1291        |
| Traditional Therapies Exploit the Genetic Instability and Loss of Cell-Cycle Checkpoint Responses in Cancer Cells               | 1257 | <i>Summary</i>   | 1292        |
| New Drugs Can Exploit the Specific Cause of a Tumor's Genetic Instability   | 1257 | SPERM  | 1292        |
| Genetic Instability Helps Cancers Become Progressively More Resistant to Therapies  | 1259 | Sperm Are Highly Adapted for Delivering Their DNA to an Egg                      | 1292        |
| New Therapies Are Emerging from Our Knowledge of Cancer Biology   | 1260 | Sperm Are Produced Continuously in the Mammalian Testis                          | 1293        |
| Small Molecules Can Be Designed to Inhibit Specific Oncogenic Proteins  | 1260 | Sperm Develop as a Syncytium   | 1294        |
| Tumor Blood Vessels Are Logical Targets for Cancer Therapy  | 1262 | <i>Summary</i>   | 1296        |
| Many Cancers May Be Treatable by Enhancing the Immune Response Against a Specific Tumor   | 1262 | FERTILIZATION  | 1297        |
| Treating Patients with Several Drugs Simultaneously Has Potential Advantages for Cancer Therapy                                 | 1263 | Ejaculated Sperm Become Capacitated in the Female Genital Tract                  | 1297        |
| Gene Expression Profiling Can Help Classify Cancers into Clinically Meaningful Subgroups  | 1264 | Capacitated Sperm Bind to the Zona Pellucida and Undergo an Acrosome Reaction    | 1298        |
|   |      | The Mechanism of Sperm–Egg Fusion Is Still Unknown                               | 1298        |
|   |      | Sperm Fusion Activates the Egg by Increasing Ca <sup>2+</sup> in the Cytosol     | 1299        |
|   |      | The Cortical Reaction Helps Ensure That Only One Sperm Fertilizes the Egg        | 1300        |
|   |      | The Sperm Provides Centrioles as Well as Its Genome to the Zygote                | 1301        |
|   |      | IVF and ICSI Have Revolutionized the Treatment of Human Infertility              | 1301        |
|   |      | <i>Summary</i>   | 1303        |
|   |      | <i>References</i>  | 1304        |
|   |      | <b>Chapter 22 Development of Multicellular Organisms</b>                         | <b>1305</b> |
|   |      | UNIVERSAL MECHANISMS OF ANIMAL DEVELOPMENT                                       | 1305        |
|   |      | Animals Share Some Basic Anatomical Features                                     | 1307        |

|  |      |   |      |
|--|------|---|------|
| Multicellular Animals Are Enriched in Proteins Mediating Cell Interactions and Gene Regulation                                   | 1308 | Egg-Polarity, Gap, and Pair-Rule Genes Create a Transient Pattern That Is Remembered by Other Genes           | 1340 |
| Regulatory DNA Defines the Program of Development  | 1309 | <i>Summary</i>  | 1341 |
| Manipulation of the Embryo Reveals the Interactions Between Its Cells  | 1310 | <b>HOMEOTIC SELECTOR GENES AND THE PATTERNING OF THE ANTEROPOSTERIOR AXIS</b>                                 | 1341 |
| Studies of Mutant Animals Identify the Genes That Control Developmental Processes  | 1311 | The Hox Code Specifies Anterior-Posterior Differences   | 1342 |
| A Cell Makes Developmental Decisions Long Before It Shows a Visible Change   | 1311 | Homeotic Selector Genes Code for DNA-Binding Proteins That Interact with Other Gene Regulatory Proteins       | 1342 |
| Cells Have Remembered Positional Values That Reflect Their Location in the Body  | 1312 | The Homeotic Selector Genes Are Expressed Sequentially According to Their Order in the Hox Complex            | 1343 |
| Inductive Signals Can Create Orderly Differences Between Initially Identical Cells   | 1313 | The Hox Complex Carries a Permanent Record of Positional Information  | 1344 |
| Sister Cells Can Be Born Different by an Asymmetric Cell Division  | 1313 | The Anteroposterior Axis Is Controlled by <i>Hox</i> Selector Genes in Vertebrates Also                       | 1344 |
| Positive Feedback Can Create Asymmetry Where There Was None Before   | 1314 | <i>Summary</i>  | 1347 |
| Positive Feedback Generates Patterns, Creates All-or-None Outcomes, and Provides Memory  | 1315 | <b>ORGANOGENESIS AND THE PATTERNING OF APPENDAGES</b>   | 1347 |
| A Small Set of Signaling Pathways, Used Repeatedly, Controls Developmental Patterning  | 1316 | Conditional and Induced Somatic Mutations Make it Possible to Analyze Gene Functions Late in Development      | 1348 |
| Morphogens Are Long-Range Inducers That Exert Graded Effects   | 1316 | Body Parts of the Adult Fly Develop From Imaginal Discs   | 1349 |
| Extracellular Inhibitors of Signal Molecules Shape the Response to the Inducer   | 1317 | Homeotic Selector Genes Are Essential for the Memory of Positional Information in Imaginal Disc Cells         | 1351 |
| Developmental Signals Can Spread Through Tissue in Several Different Ways  | 1318 | Specific Regulatory Genes Define the Cells That Will Form an Appendage  | 1351 |
| Programs That Are Intrinsic to a Cell Often Define the Time-Course of its Development  | 1319 | The Insect Wing Disc Is Divided into Compartments   | 1352 |
| Initial Patterns Are Established in Small Fields of Cells and Refined by Sequential Induction as the Embryo Grows                | 1319 | Four Familiar Signaling Pathways Combine to Pattern the Wing Disc: Wingless, Hedgehog, Dpp, and Notch         | 1353 |
| <i>Summary</i>   | 1320 | The Size of Each Compartment Is Regulated by Interactions Among Its Cells                                     | 1353 |
| <b>CAENORHABDITIS ELEGANS: DEVELOPMENT FROM THE PERSPECTIVE OF THE INDIVIDUAL CELL</b>   | 1321 | Similar Mechanisms Pattern the Limbs of Vertebrates   | 1355 |
| <i>Caenorhabditis elegans</i> Is Anatomically Simple   | 1321 | Localized Expression of Specific Classes of Gene Regulatory Proteins Foreshadows Cell Differentiation         | 1356 |
| Cell Fates in the Developing Nematode Are Almost Perfectly Predictable   | 1322 | Lateral Inhibition Singles Out Sensory Mother Cells Within Proneural Clusters                                 | 1357 |
| Products of Maternal-Effect Genes Organize the Asymmetric Division of the Egg  | 1323 | Lateral Inhibition Drives the Progeny of the Sensory Mother Cell Toward Different Final Fates                 | 1357 |
| Progressively More Complex Patterns Are Created by Cell-Cell Interactions  | 1324 | Planar Polarity of Asymmetric Divisions is Controlled by Signaling via the Receptor Frizzled                  | 1358 |
| Microsurgery and Genetics Reveal the Logic of Developmental Control; Gene Cloning and Sequencing Reveal Its Molecular Mechanisms | 1325 | Asymmetric Stem-Cell Divisions Generate Additional Neurons in the Central Nervous System                      | 1359 |
| Cells Change Over Time in Their Responsiveness to Developmental Signals  | 1325 | Asymmetric Neuroblast Divisions Segregate an Inhibitor of Cell Division into Just One of the Daughter Cells   | 1361 |
| Heterochronic Genes Control the Timing of Development  | 1326 | Notch Signaling Regulates the Fine-Grained Pattern of Differentiated Cell Types in Many Different Tissues     | 1362 |
| Cells Do Not Count Cell Divisions in Timing Their Internal Programs  | 1327 | Some Key Regulatory Genes Define a Cell Type; Others Can Activate the Program for Creation of an Entire Organ | 1362 |
| Selected Cells Die by Apoptosis as Part of the Program of Development  | 1327 | <i>Summary</i>  | 1363 |
| <i>Summary</i>   | 1328 | <b>CELL MOVEMENTS AND THE SHAPING OF THE VERTEBRATE BODY</b>  | 1363 |
| <b>DROSOPHILA AND THE MOLECULAR GENETICS OF PATTERN FORMATION: GENESIS OF THE BODY PLAN</b>                                      | 1328 | The Polarity of the Amphibian Embryo Depends on the Polarity of the Egg                                       | 1364 |
| The Insect Body Is Constructed as a Series of Segmental Units  | 1329 | Cleavage Produces Many Cells from One   | 1365 |
| <i>Drosophila</i> Begins Its Development as a Syncytium  | 1330 | Gastrulation Transforms a Hollow Ball of Cells into a Three-Layered Structure with a Primitive Gut            | 1365 |
| Genetic Screens Define Groups of Genes Required for Specific Aspects of Early Patterning   | 1332 | The Movements of Gastrulation Are Precisely Predictable   | 1366 |
| Interactions of the Oocyte With Its Surroundings Define the Axes of the Embryo: the Role of the Egg-Polarity Genes               | 1333 | Chemical Signals Trigger the Mechanical Processes   | 1367 |
| The Dorsovernal Signaling Genes Create a Gradient of a Nuclear Gene Regulatory Protein   | 1334 | Active Changes of Cell Packing Provide a Driving Force for Gastrulation                                       | 1368 |
| Dpp and Sog Set Up a Secondary Morphogen Gradient to Refine the Pattern of the Dorsal Part of the Embryo                         | 1336 | Changing Patterns of Cell Adhesion Molecules Force Cells Into New Arrangements                                | 1369 |
| The Insect Dorsovernal Axis Corresponds to the Vertebrate Ventrodorsal Axis  | 1336 | The Notochord Elongates, While the Neural Plate Rolls Up to Form the Neural Tube                              | 1370 |
| Three Classes of Segmentation Genes Refine the Anterior-Posterior Maternal Pattern and Subdivide the Embryo                      | 1336 | A Gene-Expression Oscillator Controls Segmentation of the Mesoderm Into Somites                               | 1371 |
| The Localized Expression of Segmentation Genes Is Regulated by a Hierarchy of Positional Signals                                 | 1337 | Delayed Negative Feedback May Generate the Oscillations of the Segmentation Clock                             | 1373 |
| The Modular Nature of Regulatory DNA Allows Genes to Have Multiple Independently Controlled Functions                            | 1339 | Embryonic Tissues Are Invaded in a Strictly Controlled Fashion by Migratory Cells                             | 1373 |
|  |      | The Distribution of Migrant Cells Depends on Survival Factors as Well as Guidance Cues                        | 1375 |



|   |             |   |      |
|---|-------------|---|------|
| Left–Right Asymmetry of the Vertebrate Body Derives From Molecular Asymmetry in the Early Embryo                  | 1376        | The Basal Layer Contains Both Stem Cells and Transit Amplifying Cells   | 1422 |
| <i>Summary</i>  | 1377        | Transit amplifying Divisions Are Part of the Strategy of Growth Control   | 1423 |
| <b>THE MOUSE</b>  | 1378        | Stem Cells of Some Tissues Selectively Retain Original DNA Strands  | 1424 |
| Mammalian Development Begins With a Specialized Preamble  | 1378        | The Rate of Stem-Cell Division Can Increase Dramatically When New Cells Are Needed Urgently                     | 1425 |
| The Early Mammalian Embryo Is Highly Regulative   | 1380        | Many Interacting Signals Govern Epidermal Renewal   | 1426 |
| Totipotent Embryonic Stem Cells Can Be Obtained From a Mammalian Embryo   | 1380        | The Mammary Gland Undergoes Cycles of Development and Regression  | 1426 |
| Interactions Between Epithelium and Mesenchyme Generate Branching Tubular Structures                              | 1381        | <i>Summary</i>  | 1428 |
| <i>Summary</i>  | 1382        | <b>SENSORY EPITHELIA</b>  | 1429 |
| <b>NEURAL DEVELOPMENT</b>   | 1383        | Olfactory Sensory Neurons Are Continually Replaced  | 1429 |
| Neurons Are Assigned Different Characters According to the Time and Place Where They Are Born                     | 1383        | Auditory Hair Cells Have to Last a Lifetime   | 1430 |
| The Character Assigned to a Neuron at Its Birth Governs the Connections It Will Form                              | 1385        | Most Permanent Cells Renew Their Parts: the Photoreceptor Cells of the Retina                                   | 1432 |
| Each Axon or Dendrite Extends by Means of a Growth Cone at Its Tip  | 1386        | <i>Summary</i>  | 1433 |
| The Growth Cone Pilots the Developing Neurite Along a Precisely Defined Path <i>In Vivo</i>                       | 1387        | <b>THE AIRWAYS AND THE GUT</b>  | 1434 |
| Growth Cones Can Change Their Sensibilities as They Travel  | 1389        | Adjacent Cell Types Collaborate in the Alveoli of the Lungs   | 1434 |
| Target Tissues Release Neurotrophic Factors That Control Nerve Cell Growth and Survival                           | 1389        | Goblet Cells, Ciliated Cells, and Macrophages Collaborate to Keep the Airways Clean                             | 1434 |
| Neuronal Specificity Guides the Formation of Orderly Neural Maps  | 1391        | The Lining of the Small Intestine Renews Itself Faster Than Any Other Tissue                                    | 1436 |
| Axons From Different Regions of the Retina Respond Differently to a Gradient of Repulsive Molecules in the Tectum | 1392        | Wnt Signaling Maintains the Gut Stem-Cell Compartment   | 1438 |
| Diffuse Patterns of Synaptic Connections Are Sharpened by Activity-Dependent Remodeling                           | 1393        | Notch Signaling Controls Gut Cell Diversification   | 1439 |
| Experience Molds the Pattern of Synaptic Connections in the Brain   | 1395        | Ephrin–Eph Signaling Controls the Migrations of Gut Epithelial Cells  | 1440 |
| Adult Memory and Developmental Synapse Remodeling May Depend on Similar Mechanisms                                | 1396        | Wnt, Hedgehog, PDGF, and BMP Signaling Pathways Combine to Delimit the Stem-Cell Niche                          | 1441 |
| <i>Summary</i>  | 1397        | The Liver Functions as an Interface Between the Digestive Tract and the Blood                                   | 1442 |
| <b>PLANT DEVELOPMENT</b>  | 1398        | Liver Cell Loss Stimulates Liver Cell Proliferation   | 1443 |
| <i>Arabidopsis</i> Serves as a Model Organism for Plant Molecular Genetics  | 1398        | Tissue Renewal Does Not Have to Depend on Stem Cells: Insulin-Secreting Cells in the Pancreas                   | 1444 |
| The <i>Arabidopsis</i> Genome Is Rich in Developmental Control Genes  | 1399        | <i>Summary</i>  | 1445 |
| Embryonic Development Starts by Establishing a Root–Shoot Axis and Then Halts Inside the Seed                     | 1400        | <b>BLOOD VESSELS, LYMPHATICS, AND ENDOTHELIAL CELLS</b>   | 1445 |
| The Parts of a Plant Are Generated Sequentially by Meristems  | 1403        | Endothelial Cells Line All Blood Vessels and Lymphatics   | 1445 |
| Development of the Seedling Depends on Environmental Signals  | 1403        | Endothelial Tip Cells Pioneer Angiogenesis  | 1446 |
| Long-Range Hormonal Signals Coordinate Developmental Events in Separate Parts of the Plant                        | 1403        | Different Types of Endothelial Cells Form Different Types of Vessel   | 1447 |
| The Shaping of Each New Structure Depends on Oriented Cell Division and Expansion                                 | 1406        | Tissues Requiring a Blood Supply Release VEGF; Notch Signaling Between Endothelial Cells Regulates the Response | 1448 |
| Each Plant Module Grows From a Microscopic Set of Primordia in a Meristem   | 1407        | Signals from Endothelial Cells Control Recruitment of Pericytes and Smooth Muscle Cells to Form the Vessel Wall | 1450 |
| Polarized Auxin Transport Controls the Pattern of Primordia in the Meristem                                       | 1408        | <i>Summary</i>  | 1450 |
| Cell Signaling Maintains the Meristem   | 1409        | <b>RENEWAL BY MULTIPOTENT STEM CELLS: BLOOD CELL FORMATION</b>  | 1450 |
| Regulatory Mutations Can Transform Plant Topology by Altering Cell Behavior in the Meristem                       | 1410        | The Three Main Categories of White Blood Cells Are Granulocytes, Monocytes, and Lymphocytes                     | 1451 |
| The Switch to Flowering Depends on Past and Present Environmental Cues  | 1412        | The Production of Each Type of Blood Cell in the Bone Marrow Is Individually Controlled                         | 1453 |
| Homeotic Selector Genes Specify the Parts of a Flower   | 1413        | Bone Marrow Contains Hemopoietic Stem Cells   | 1454 |
| <i>Summary</i>  | 1415        | A Multipotent Stem Cell Gives Rise to All Classes of Blood Cells  | 1456 |
| <i>References</i>   | 1415        | Commitment Is a Stepwise Process  | 1456 |
| <b>Chapter 23 Specialized Tissues, Stem Cells, and Tissue Renewal</b>   | <b>1417</b> | Divisions of Committed Progenitor Cells Amplify the Number of Specialized Blood Cells                           | 1457 |
| <b>EPIDERMIS AND ITS RENEWAL BY STEM CELLS</b>  | 1417        | Stem Cells Depend on Contact Signals From Stromal Cells   | 1458 |
| Epidermal Cells Form a Multilayered Waterproof Barrier  | 1419        | Factors That Regulate Hemopoiesis Can Be Analyzed in Culture  | 1459 |
| Differentiating Epidermal Cells Express a Sequence of Different Genes as They Mature                              | 1420        | Erythropoiesis Depends on the Hormone Erythropoietin  | 1459 |
| Stem Cells in the Basal Layer Provide for Renewal of the Epidermis  | 1420        | Multiple CSFs Influence Neutrophil and Macrophage Production  | 1460 |
| The Two Daughters of a Stem Cell Do Not Always Have to Become Different   | 1421        | The Behavior of a Hemopoietic Cell Depends Partly on Chance   | 1461 |
|   |             | Regulation of Cell Survival Is as Important as Regulation of Cell Proliferation                                 | 1462 |
|   |             | <i>Summary</i>  | 1462 |
|   |             | <b>GENESIS, MODULATION, AND REGENERATION OF SKELETAL MUSCLE</b>   | 1463 |
|   |             | Myoblasts Fuse to Form New Skeletal Muscle Fibers   | 1464 |

|  |      |  |      |
|--|------|--|------|
| Muscle Cells Can Vary Their Properties by Changing the Protein Isoforms They Contain                                 | 1465 | Pathogens Evolve Rapidly   | 1518 |
| Skeletal Muscle Fibers Secrete Myostatin to Limit Their Own Growth   | 1465 | Antigenic Variation in Pathogens Occurs by Multiple Mechanisms   | 1519 |
| Some Myoblasts Persist as Quiescent Stem Cells in the Adult  | 1466 | Error-Prone Replication Dominates Viral Evolution  | 1520 |
| <i>Summary</i>   | 1467 | Drug-Resistant Pathogens Are a Growing Problem   | 1521 |
| <b>FIBROBLASTS AND THEIR TRANSFORMATIONS: THE CONNECTIVE-TISSUE CELL FAMILY</b>                                      | 1467 | <i>Summary</i>   | 1524 |
| Fibroblasts Change Their Character in Response to Chemical Signals   | 1467 | <b>BARRIERS TO INFECTION AND THE INNATE IMMUNE SYSTEM</b>  | 1524 |
| The Extracellular Matrix May Influence Connective-Tissue Cell Differentiation by Affecting Cell Shape and Attachment | 1468 | Epithelial Surfaces and Defensins Help Prevent Infection   | 1525 |
| Osteoblasts Make Bone Matrix   | 1469 | Human Cells Recognize Conserved Features of Pathogens  | 1526 |
| Most Bones Are Built Around Cartilage Models   | 1470 | Complement Activation Targets Pathogens for Phagocytosis or Lysis  | 1528 |
| Bone Is Continually Remodeled by the Cells Within It   | 1472 | Toll-like Proteins and NOD Proteins Are an Ancient Family of Pattern Recognition Receptors                               | 1530 |
| Osteoclasts Are Controlled by Signals From Osteoblasts   | 1473 | Phagocytic Cells Seek, Engulf, and Destroy Pathogens   | 1531 |
| Fat Cells Can Develop From Fibroblasts   | 1474 | Activated Macrophages Contribute to the Inflammatory Response at Sites of Infection                                      | 1533 |
| Leptin Secreted by Fat Cells Provides Feedback to Regulate Eating  | 1475 | Virus-Infected Cells Take Drastic Measures to Prevent Viral Replication  | 1534 |
| <i>Summary</i>   | 1476 | Natural Killer Cells Induce Virus-Infected Cells to Kill Themselves  | 1535 |
| <b>STEM-CELL ENGINEERING</b>   | 1476 | Dendritic Cells Provide the Link Between the Innate and Adaptive Immune Systems  | 1536 |
| Hemopoietic Stem Cells Can Be Used to Replace Diseased Blood Cells with Healthy Ones                                 | 1477 | <i>Summary</i>   | 1537 |
| Epidermal Stem Cell Populations Can Be Expanded in Culture for Tissue Repair   | 1477 | <i>References</i>  | 1537 |
| Neural Stem Cells Can Be Manipulated in Culture  | 1478 | <b>Chapter 25 The Adaptive Immune System</b>   | 1539 |
| Neural Stem Cells Can Repopulate the Central Nervous System  | 1478 | <b>LYMPHOCYTES AND THE CELLULAR BASIS OF ADAPTIVE IMMUNITY</b>   | 1540 |
| Stem Cells in the Adult Body Are Tissue-Specific   | 1479 | Lymphocytes Are Required for Adaptive Immunity   | 1540 |
| ES Cells Can Make Any Part of the Body   | 1480 | The Innate and Adaptive Immune Systems Work Together   | 1541 |
| Patient-Specific ES Cells Could Solve the Problem of Immune Rejection  | 1481 | B Lymphocytes Develop in the Bone Marrow; T Lymphocytes Develop in the Thymus  | 1543 |
| ES Cells Are Useful for Drug Discovery and Analysis of Disease   | 1482 | The Adaptive Immune System Works by Clonal Selection   | 1544 |
| <i>Summary</i>   | 1482 | Most Antigens Activate Many Different Lymphocyte Clones  | 1545 |
| <i>References</i>  | 1483 | Immunological Memory Involves Both Clonal Expansion and Lymphocyte Differentiation                                       | 1545 |
| <b>Chapter 24 Pathogens, Infection, and Innate Immunity</b>  | 1485 | Immunological Tolerance Ensures That Self Antigens Are Not Normally Attacked   | 1547 |
| <b>INTRODUCTION TO PATHOGENS</b>   | 1486 | Lymphocytes Continuously Circulate Through Peripheral Lymphoid Organs  | 1549 |
| Pathogens Have Evolved Specific Mechanisms for Interacting with Their Hosts  | 1486 | <i>Summary</i>   | 1551 |
| The Signs and Symptoms of Infection May Be Caused by the Pathogen or by the Host's Responses                         | 1487 | <b>B CELLS AND ANTIBODIES</b>  | 1551 |
| Pathogens Are Phylogenetically Diverse   | 1488 | B Cells Make Antibodies as Both Cell-Surface Antigen Receptors and Secreted Proteins                                     | 1552 |
| Bacterial Pathogens Carry Specialized Virulence Genes  | 1489 | A Typical Antibody Has Two Identical Antigen-Binding Sites   | 1552 |
| Fungal and Protozoan Parasites Have Complex Life Cycles with Multiple Forms  | 1494 | An Antibody Molecule Is Composed of Heavy and Light Chains   | 1552 |
| All Aspects of Viral Propagation Depend on Host Cell Machinery   | 1496 | There Are Five Classes of Antibody Heavy Chains, Each with Different Biological Properties                               | 1553 |
| Prions Are Infectious Proteins   | 1498 | The Strength of an Antibody–Antigen Interaction Depends on Both the Number and the Affinity of the Antigen-Binding Sites | 1557 |
| Infectious Disease Agents Are Linked To Cancer, Heart Disease, and Other Chronic Illnesses                           | 1499 | Antibody Light and Heavy Chains Consist of Constant and Variable Regions   | 1558 |
| <i>Summary</i>   | 1501 | The Light and Heavy Chains Are Composed of Repeating Ig Domains  | 1559 |
| <b>CELL BIOLOGY OF INFECTION</b>   | 1501 | An Antigen-Binding Site Is Constructed from Hypervariable Loops  | 1560 |
| Pathogens Cross Protective Barriers to Colonize the Host   | 1501 | <i>Summary</i>   | 1561 |
| Pathogens That Colonize Epithelia Must Avoid Clearance by the Host   | 1502 | <b>THE GENERATION OF ANTIBODY DIVERSITY</b>  | 1562 |
| Intracellular Pathogens Have Mechanisms for Both Entering and Leaving Host Cells                                     | 1504 | Antibody Genes Are Assembled From Separate Gene Segments During B Cell Development                                       | 1562 |
| Virus Particles Bind to Molecules Displayed on the Host Cell Surface   | 1505 | Imprecise Joining of Gene Segments Greatly Increases the Diversity of V Regions  | 1564 |
| Virions Enter Host Cells by Membrane Fusion, Pore Formation, or Membrane Disruption                                  | 1506 | The Control of V(D)J Recombination Ensures That B Cells Are Monospecific   | 1565 |
| Bacteria Enter Host Cells by Phagocytosis  | 1507 | Antigen-Driven Somatic Hypermutation Fine-Tunes Antibody Responses   | 1566 |
| Intracellular Eucaryotic Parasites Actively Invade Host Cells  | 1508 | B Cells Can Switch the Class of Antibody They Make   | 1567 |
| Many Pathogens Alter Membrane Traffic in the Host Cell   | 1511 | <i>Summary</i>   | 1568 |
| Viruses and Bacteria Use the Host Cell Cytoskeleton for Intracellular Movement                                       | 1514 |  |      |
| Viral Infections Take Over the Metabolism of the Host Cell   | 1517 |  |      |
| Pathogens Can Alter the Behavior of the Host Organism to Facilitate the Spread of the Pathogen                       | 1518 |  |      |

|  |      |   |      |
|--|------|---|------|
| T CELLS AND MHC PROTEINS   | 1569 | Most Developing Cytotoxic and Helper T Cells That Could Be Activated by Self-Peptide–MHC Complexes Are Eliminated in the Thymus | 1586 |
| T Cell Receptors (TCRs) Are Antibodylike Heterodimers  | 1570 | Some Organ-Specific Proteins Are Ectopically Expressed in the Thymus Medulla  | 1587 |
| Antigen Presentation by Dendritic Cells Can Either Activate or Tolerize T Cells                              | 1571 | The Function of MHC Proteins Helps Explain Their Polymorphism   | 1588 |
| Effector Cytotoxic T Cells Induce Infected Target Cells to Kill Themselves                                   | 1572 | <i>Summary</i>  | 1588 |
| Effector Helper T Cells Help Activate Other Cells of the Innate and Adaptive Immune Systems                  | 1573 | HELPER T CELLS AND LYMPHOCYTE ACTIVATION  | 1589 |
| Regulatory T Cells Suppress the Activity of Other T Cells  | 1574 | Activated Dendritic Cells Use Multiple Mechanisms to Activate T Cells   | 1590 |
| T Cells Recognize Foreign Peptides Bound to MHC Proteins   | 1575 | The Activation of T Cells Is Controlled by Negative Feedback  | 1591 |
| MHC Proteins Were Identified in Transplantation Reactions Before Their Functions Were Known                  | 1575 | The Subclass of Effector Helper T Cell Determines the Nature of the Adaptive Immune Response                                    | 1592 |
| Class I and Class II MHC Proteins Are Structurally Similar Heterodimers                                      | 1576 | T <sub>H</sub> 1 Cells Activate Infected Macrophages and Stimulate An Inflammatory Response                                     | 1594 |
| An MHC Protein Binds a Peptide and Interacts with a T Cell Receptor  | 1577 | Antigen Binding to B Cell Receptors (BCRs) Is Only One Step in B Cell Activation  | 1595 |
| MHC Proteins Help Direct T Cells to Their Appropriate Targets  | 1579 | Antigen-Specific Helper T Cells Are Essential for Activating Most B Cells   | 1597 |
| CD4 and CD8 Co-Receptors Bind to Invariant Parts of MHC Proteins   | 1580 | A Special Class of B Cells Recognize T-Cell-Independent Antigens  | 1598 |
| Cytotoxic T Cells Recognize Fragments of Foreign Cytosolic Proteins in Association with Class I MHC Proteins | 1581 | Immune Recognition Molecules Belong to the Ancient Ig Superfamily   | 1599 |
| Helper T Cells Respond to Fragments of Endocytosed Foreign Protein Associated with Class II MHC Proteins     | 1583 | <i>Summary</i>  | 1600 |
| Potentially Useful T Cells Are Positively Selected in the Thymus   | 1585 | <i>References</i>   | 1600 |