Contents

| Lists of Figures, Tables and Abbreviations Preface | | viii xii |
|---|--|-------------|
| 1 | Introduction | I |
| 2 | The Use of Statistical Methods in Social Research | 7 |
| 3 | Cases and Variables | 21 |
| 4 | The Logic of Sampling | 34 |
| 5 | Estimating and Measuring One Important Thing | 51 |
| 6 | Studying the Relationship between Two Variables | 71 |
| 7 | Linear Regression Models and Their Generalizations | 91 |
| 8 | Time Matters | 113 |
| 9 | Statistical Case-oriented Methods | 132 |
| 10 | Methods for Analysing Latent Variables | 153 |
| 11 | Causal Analysis | 173 |
| Bibliography Index | | 190 195 |

Lists of Figures, Tables and Abbreviations

Figures

| Figure 4.1 | Distinguishing cluster sampling from stratified sampling | 47 |
|-------------|--|-----|
| Figure 5.1 | Distribution of respondents across religions in the UK, 2006 | 59 |
| Figure 5.2 | Religiousness in the UK, 2006 | 60 |
| Figure 5.3 | Statistic for sample as parameter for population | 62 |
| Figure 6.1 | Relationship between r and scatter plot | 73 |
| Figure 6.2 | Correlation in four sections of a two-dimensional space | 74 |
| Figure 6.3 | Anscombe's Quartet: Same regression line, different datasets | 85 |
| Figure 6.4 | Difference between males and females in terms of | |
| | the number of hours chosen to work, UK, 2006 | 87 |
| Figure 7.1 | Ideal situation for a multiple linear regression model | 95 |
| Figure 7.2 | Realistic situation for a multiple linear regression model | 95 |
| Figure 7.3 | Effect of correlated error terms on classical linear | |
| | regression model | 106 |
| Figure 7.4 | Effect of ignoring grouping on statistical significance | 106 |
| Figure 8.1 | Different types of cases in an event history study | 124 |
| Figure 9.1 | An illustrated dendrogram on social trust | 138 |
| Figure 9.2 | Mapping out the relative positions of three cases | |
| | in one dimension | 146 |
| Figure 9.3 | A biplot of access to the Internet and opinions | |
| | on homosexuality | 151 |
| Figure 10.1 | An illustration of variances of four observed variables | 157 |
| Figure 10.2 | An illustration of scree plot | 159 |
| Figure 10.3 | A confirmatory factor model with four observed variables | 163 |
| Figure 10.4 | A two-factor measurement model | 163 |
| Figure 10.5 | A random intercept and random slope latent | |
| | growth curve model | 170 |
| Figure 11.1 | Ideal situation for inferring causal effect | 183 |
| Figure 11.7 | Observation available for treatment but not for control | 104 |

| Figure 11.3 | Observation available for control but not for treatment | 184 |
|-------------|---|-----|
| Figure 11.4 | Association of the error term with assignment | 187 |
| Figure 11.5 | Use of an instrumental variable | 188 |

Tables

| Table 3.1 | Case-by-variable matrix | 22 |
|------------|--|-----|
| Table 3.2 | Levels of measurement and their relations | 29 |
| Table 4.1 | US population structure by age, gender and the number | |
| | of races, 2000 | 39 |
| Table 4.2 | Scenarios of bias and precision in probability sampling | 42 |
| Table 5.1 | Effect of <i>k</i> on <i>D</i> and <i>I</i> Q <i>V</i> | 61 |
| Table 6.1 | A generic 2 × 2 table | 76 |
| Table 6.2 | British identity and ethnicity | 79 |
| Table 6.3 | A generic 3×3 table of two ordinal variables | 81 |
| Table 6.4 | Components of the F-test | 87 |
| Table 7.1 | Regression output for the supervisor performance example | 96 |
| Table 7.2 | Data matrix without grouping | 104 |
| Table 7.3 | Data matrix with grouping | 104 |
| Table 7.4 | Lower level data matrix | 105 |
| Table 7.5 | Higher level data matrix | 105 |
| Table 7.6 | Scenarios of random intercept and coefficients in | |
| | multilevel modelling | 107 |
| Table 8.1 | An example of the long form data matrix in a | |
| | longitudinal study | 118 |
| Table 8.2 | Wide form of event history data | 125 |
| Table 8.3 | A protocol of the life table | 126 |
| Table 8.4 | Long form of event history data matrix | 126 |
| Table 9.1 | A classification matrix for two groups | 142 |
| Table 9.2 | An illustration of similarity matrix with four objects | 147 |
| Table 9.3 | Personal use of Internet/email/WWW and opinions | |
| | on homosexuality | 150 |
| Table 10.1 | A hypothetical correlation matrix | 156 |
| Table 10.2 | Re-organization of Table 10.1 | 156 |
| Table 10.3 | An ideal factor loadings table | 160 |
| Table 10.4 | Correlations of the indicators measured with | |
| | a four-point scale | 162 |

| Table 10.5 | Correlations of the indicators measured with | |
|------------|--|-----|
| | an 11-point scale | 162 |
| Table 10.6 | A generic covariance matrix of five variables | 167 |
| Table 10.7 | A generic correlation matrix of five variables | 167 |
| Table 11.1 | Scenarios of cause and outcome | 175 |
| Table 11.2 | Four scenarios of an instrumental variable | |
| | and assignment | 189 |

Abbreviations

| ANOVA | analysis of variance |
|---------|--|
| BHPS | British Household Panel Study |
| CFA | confirmatory factor analysis |
| CFI | comparative fit index |
| CLA | cluster analysis |
| COA | correspondence analysis |
| CPI | consumer price index |
| DA | discriminant analysis |
| DE | design effect |
| df | degrees of freedom |
| EFA | exploratory factor analysis |
| ESS | European Social Survey |
| GSS | General Social Survey |
| HLM | hierarchical linear models (also the name of software) |
| ICC | intra-class correlation coefficient |
| IQR | inter-quartile range |
| IQV | index of qualitative variation |
| LATE | local average treatment effect |
| LGCM | latent growth curve models |
| MANCOVA | multiple analysis of covariance |
| MANOVA | multiple analysis of variance |
| MDS | multidimensional scaling |
| PCA | principal component analysis |
| PSID | Panel Study of Income Dynamics |
| PSU | primary sampling unit |
| QCA | qualitative comparative analysis |
| RMSEA | root mean square error of approximation |
| RMSR | root means square residual |
| s.d. | standard deviation |
| s.e. | standard error |
| SEM | structural equation models |
| SES | socio-economic status |
| | |

X