

CONTENTS

Preface v

Part I THE MOLECULAR DESIGN OF LIFE

Chapter 1 Biochemistry: An Evolving Science 1

1.1 Biochemical Unity Underlies Biological Diversity 1

1.2 DNA Illustrates the Interplay Between Form and Function 4

DNA is constructed from four building blocks 4

Two single strands of DNA combine to form a double helix 5

DNA structure explains heredity and the storage of information 5

1.3 Concepts from Chemistry Explain the Properties of Biological Molecules 6

The double helix can form from its component strands 6

Covalent and noncovalent bonds are important for the structure and stability of biological molecules 7

The double helix is an expression of the rules of chemistry 10

The laws of thermodynamics govern the behavior of biochemical systems 11

Heat is released in the formation of the double helix 12

Acid–base reactions are central in many biochemical processes 13

Acid–base reactions can disrupt the double helix 14

Buffers regulate pH in organisms and in the laboratory 15

1.4 The Genomic Revolution Is Transforming Biochemistry and Medicine 17

The sequencing of the human genome is a landmark in human history 17

Genome sequences encode proteins and patterns of expression 18

Individuality depends on the interplay between genes and environment 19

APPENDIX: Visualizing Molecular Structures I: Small Molecules 21

Chapter 2 Protein Composition and Structure 25

2.1 Proteins Are Built from a Repertoire of 20 Amino Acids 27

2.2 Primary Structure: Amino Acids Are Linked by Peptide Bonds to Form Polypeptide Chains 33

Proteins have unique amino acid sequences specified by genes 35

Polypeptide chains are flexible yet conformationally restricted 36

2.3 Secondary Structure: Polypeptide Chains Can Fold into Regular Structures Such As the Alpha Helix, the Beta Sheet, and Turns and Loops	38	3.2 Amino Acid Sequences of Proteins Can Be Determined Experimentally	81
The alpha helix is a coiled structure stabilized by intrachain hydrogen bonds	38	Peptide sequences can be determined by automated Edman degradation	82
Beta sheets are stabilized by hydrogen bonding between polypeptide strands	40	Proteins can be specifically cleaved into small peptides to facilitate analysis	84
Polypeptide chains can change direction by making reverse turns and loops	42	Genomic and proteomic methods are complementary	86
Fibrous proteins provide structural support for cells and tissues	43	3.3 Immunology Provides Important Techniques with Which to Investigate Proteins	86
2.4 Tertiary Structure: Water-Soluble Proteins Fold into Compact Structures with Nonpolar Cores	45	Antibodies to specific proteins can be generated	86
2.5 Quaternary Structure: Polypeptide Chains Can Assemble into Multisubunit Structures	48	Monoclonal antibodies with virtually any desired specificity can be readily prepared	88
2.6 The Amino Acid Sequence of a Protein Determines Its Three-Dimensional Structure	49	Proteins can be detected and quantified by using an enzyme-linked immunosorbent assay	90
Amino acids have different propensities for forming alpha helices, beta sheets, and beta turns	50	Western blotting permits the detection of proteins separated by gel electrophoresis	91
Protein folding is a highly cooperative process	52	Fluorescent markers make the visualization of proteins in the cell possible	92
Proteins fold by progressive stabilization of intermediates rather than by random search	52	3.4 Mass Spectrometry Is a Powerful Technique for the Identification of Peptides and Proteins	93
Prediction of three-dimensional structure from sequence remains a great challenge	54	The mass of a protein can be precisely determined by mass spectrometry	93
Some proteins are inherently unstructured and can exist in multiple conformations	54	Peptides can be sequenced by mass spectrometry	95
Protein misfolding and aggregation are associated with some neurological diseases	55	Individual proteins can be identified by mass spectrometry	96
Protein modification and cleavage confer new capabilities	57	3.5 Peptides Can Be Synthesized by Automated Solid-Phase Methods	97
APPENDIX: Visualizing Molecular Structures II: Proteins	60	3.6 Three-Dimensional Protein Structure Can Be Determined by X-ray Crystallography and NMR Spectroscopy	100
Chapter 3 Exploring Proteins and Proteomes	67	X-ray crystallography reveals three-dimensional structure in atomic detail	100
The proteome is the functional representation of the genome	68	Nuclear magnetic resonance spectroscopy can reveal the structures of proteins in solution	103
3.1 The Purification of Proteins Is an Essential First Step in Understanding Their Function	68	Chapter 4 DNA, RNA, and the Flow of Information	113
The assay: How do we recognize the protein that we are looking for?	69	4.1 A Nucleic Acid Consists of Four Kinds of Bases Linked to a Sugar–Phosphate Backbone	114
Proteins must be released from the cell to be purified	69	RNA and DNA differ in the sugar component and one of the bases	114
Proteins can be purified according to solubility, size, charge, and binding affinity	70	Nucleotides are the monomeric units of nucleic acids	115
Proteins can be separated by gel electrophoresis and displayed	73	DNA molecules are very long	117
A protein purification scheme can be quantitatively evaluated	77	4.2 A Pair of Nucleic Acid Chains with Complementary Sequences Can Form a Double-Helical Structure	117
Ultracentrifugation is valuable for separating biomolecules and determining their masses	78	The double helix is stabilized by hydrogen bonds and van der Waals interactions	117
Protein purification can be made easier with the use of recombinant DNA technology	80	DNA can assume a variety of structural forms	119
		Z-DNA is a left-handed double helix in which backbone phosphates zigzag	120

Some DNA molecules are circular and supercoiled	121	5.2 Recombinant DNA Technology Has Revolutionized All Aspects of Biology	154
Single-stranded nucleic acids can adopt elaborate structures	121	Restriction enzymes and DNA ligase are key tools in forming recombinant DNA molecules	154
4.3 The Double Helix Facilitates the Accurate Transmission of Hereditary Information	122	Plasmids and lambda phage are choice vectors for DNA cloning in bacteria	155
Differences in DNA density established the validity of the semiconservative-replication hypothesis	123	Bacterial and yeast artificial chromosomes	157
The double helix can be reversibly melted	124	Specific genes can be cloned from digests of genomic DNA	157
4.4 DNA Is Replicated by Polymerases That Take Instructions from Templates	125	Complementary DNA prepared from mRNA can be expressed in host cells	160
DNA polymerase catalyzes phosphodiester-bridge formation	125	Proteins with new functions can be created through directed changes in DNA	162
The genes of some viruses are made of RNA	126	Recombinant methods enable the exploration of the functional effects of disease-causing mutations	163
4.5 Gene Expression Is the Transformation of DNA Information into Functional Molecules	127	5.3 Complete Genomes Have Been Sequenced and Analyzed	163
Several kinds of RNA play key roles in gene expression	127	The genomes of organisms ranging from bacteria to multicellular eukaryotes have been sequenced	164
All cellular RNA is synthesized by RNA polymerases	128	The sequencing of the human genome has been finished	165
RNA polymerases take instructions from DNA templates	130	Next-generation sequencing methods enable the rapid determination of a whole genome sequence	166
Transcription begins near promoter sites and ends at terminator sites	130	Comparative genomics has become a powerful research tool	166
Transfer RNAs are the adaptor molecules in protein synthesis	131	5.4 Eukaryotic Genes Can Be Quantitated and Manipulated with Considerable Precision	167
4.6 Amino Acids Are Encoded by Groups of Three Bases Starting from a Fixed Point	132	Gene-expression levels can be comprehensively examined	167
Major features of the genetic code	133	New genes inserted into eukaryotic cells can be efficiently expressed	169
Messenger RNA contains start and stop signals for protein synthesis	134	Transgenic animals harbor and express genes introduced into their germ lines	170
The genetic code is nearly universal	135	Gene disruption provides clues to gene function	170
4.7 Most Eukaryotic Genes Are Mosaics of Introns and Exons	135	RNA interference provides an additional tool for disrupting gene expression	171
RNA processing generates mature RNA	136	Tumor-inducing plasmids can be used to introduce new genes into plant cells	172
Many exons encode protein domains	137	Human gene therapy holds great promise for medicine	173
<hr/> Chapter 5 Exploring Genes and Genomes	145	<hr/> Chapter 6 Exploring Evolution and Bioinformatics	181
5.1 The Exploration of Genes Relies on Key Tools	146	6.1 Homologs Are Descended from a Common Ancestor	182
Restriction enzymes split DNA into specific fragments	147	6.2 Statistical Analysis of Sequence Alignments Can Detect Homology	183
Restriction fragments can be separated by gel electrophoresis and visualized	147	The statistical significance of alignments can be estimated by shuffling	185
DNA can be sequenced by controlled termination of replication	149	Distant evolutionary relationships can be detected through the use of substitution matrices	186
DNA probes and genes can be synthesized by automated solid-phase methods	150	Databases can be searched to identify homologous sequences	189
Selected DNA sequences can be greatly amplified by the polymerase chain reaction	151		
PCR is a powerful technique in medical diagnostics, forensics, and studies of molecular evolution	152		
The tools for recombinant DNA technology have been used to identify disease-causing mutations	153		

6.3 Examination of Three-Dimensional Structure Enhances Our Understanding of Evolutionary Relationships 190

Tertiary structure is more conserved than primary structure	191
Knowledge of three-dimensional structures can aid in the evaluation of sequence alignments	192
Repeated motifs can be detected by aligning sequences with themselves	192
Convergent evolution illustrates common solutions to biochemical challenges	193
Comparison of RNA sequences can be a source of insight into RNA secondary structures	194

6.4 Evolutionary Trees Can Be Constructed on the Basis of Sequence Information 195

6.5 Modern Techniques Make the Experimental Exploration of Evolution Possible 196

Ancient DNA can sometimes be amplified and sequenced	196
Molecular evolution can be examined experimentally	197

Chapter 7 Hemoglobin: Portrait of a Protein in Action 203

7.1 Myoglobin and Hemoglobin Bind Oxygen at Iron Atoms in Heme 204

Changes in heme electronic structure upon oxygen binding are the basis for functional imaging studies	205
The structure of myoglobin prevents the release of reactive oxygen species	206
Human hemoglobin is an assembly of four myoglobin-like subunits	207

7.2 Hemoglobin Binds Oxygen Cooperatively 207

Oxygen binding markedly changes the quaternary structure of hemoglobin	209
Hemoglobin cooperativity can be potentially explained by several models	210
Structural changes at the heme groups are transmitted to the $\alpha_1\beta_1$ - $\alpha_2\beta_2$ interface	212
2,3-Bisphosphoglycerate in red cells is crucial in determining the oxygen affinity of hemoglobin	212
Carbon monoxide can disrupt oxygen transport by hemoglobin	213

7.3 Hydrogen Ions and Carbon Dioxide Promote the Release of Oxygen: The Bohr Effect 214

7.4 Mutations in Genes Encoding Hemoglobin Subunits Can Result in Disease 216

Sickle-cell anemia results from the aggregation of mutated deoxyhemoglobin molecules	217
Thalassemia is caused by an imbalanced production of hemoglobin chains	218
The accumulation of free alpha-hemoglobin chains is prevented	219

Additional globins are encoded in the human genome	219
APPENDIX: Binding Models Can Be Formulated in Quantitative Terms: the Hill Plot and the Concerted Model	221

Chapter 8 Enzymes: Basic Concepts and Kinetics 227

8.1 Enzymes Are Powerful and Highly Specific Catalysts 228

Many enzymes require cofactors for activity	229
Enzymes can transform energy from one form into another	229

8.2 Free Energy Is a Useful Thermodynamic Function for Understanding Enzymes 230

The free-energy change provides information about the spontaneity but not the rate of a reaction	230
The standard free-energy change of a reaction is related to the equilibrium constant	231
Enzymes alter only the reaction rate and not the reaction equilibrium	232

8.3 Enzymes Accelerate Reactions by Facilitating the Formation of the Transition State 233

The formation of an enzyme-substrate complex is the first step in enzymatic catalysis	234
The active sites of enzymes have some common features	235
The binding energy between enzyme and substrate is important for catalysis	237

8.4 The Michaelis-Menten Equation Describes the Kinetic Properties of Many Enzymes 237

Kinetics is the study of reaction rates	237
The steady-state assumption facilitates a description of enzyme kinetics	238
Variations in K_M can have physiological consequences	240
K_M and V_{max} values can be determined by several means	240
K_M and V_{max} values are important enzyme characteristics	241
k_{cat}/K_M is a measure of catalytic efficiency	242
Most biochemical reactions include multiple substrates	243
Allosteric enzymes do not obey Michaelis-Menten kinetics	245

8.5 Enzymes Can Be Inhibited by Specific Molecules 246

Reversible inhibitors are kinetically distinguishable	247
Irreversible inhibitors can be used to map the active site	249
Transition-state analogs are potent inhibitors of enzymes	251
Catalytic antibodies demonstrate the importance of selective binding of the transition state to enzymatic activity	251
Penicillin irreversibly inactivates a key enzyme in bacterial cell-wall synthesis	252

8.6 Enzymes Can Be Studied One Molecule at a Time 254

APPENDIX: Enzymes are Classified on the Basis of the Types of Reactions That They Catalyze 256

Chapter 9 Catalytic Strategies 261

A few basic catalytic principles are used by many enzymes 262

9.1 Proteases Facilitate a Fundamentally Difficult Reaction 263

Chymotrypsin possesses a highly reactive serine residue 263

Chymotrypsin action proceeds in two steps linked by a covalently bound intermediate 264

Serine is part of a catalytic triad that also includes histidine and aspartate 265

Catalytic triads are found in other hydrolytic enzymes 268

The catalytic triad has been dissected by site-directed mutagenesis 270

Cysteine, aspartyl, and metalloproteases are other major classes of peptide-cleaving enzymes 271

Protease inhibitors are important drugs 272

9.2 Carbonic Anhydrases Make a Fast Reaction Faster 274

Carbonic anhydrase contains a bound zinc ion essential for catalytic activity 275

Catalysis entails zinc activation of a water molecule 276

A proton shuttle facilitates rapid regeneration of the active form of the enzyme 277

Convergent evolution has generated zinc-based active sites in different carbonic anhydrases 279

9.3 Restriction Enzymes Catalyze Highly Specific DNA-Cleavage Reactions 279

Cleavage is by in-line displacement of 3'-oxygen from phosphorus by magnesium-activated water 280

Restriction enzymes require magnesium for catalytic activity 282

The complete catalytic apparatus is assembled only within complexes of cognate DNA molecules, ensuring specificity 283

Host-cell DNA is protected by the addition of methyl groups to specific bases 285

Type II restriction enzymes have a catalytic core in common and are probably related by horizontal gene transfer 286

9.4 Myosins Harness Changes in Enzyme Conformation to Couple ATP Hydrolysis to Mechanical Work 287

ATP hydrolysis proceeds by the attack of water on the gamma-phosphoryl group 287

Formation of the transition state for ATP hydrolysis is associated with a substantial conformational change 288

The altered conformation of myosin persists for a substantial period of time 290

Myosins are a family of enzymes containing P-loop structures 291

Chapter 10 Regulatory Strategies 299

10.1 Aspartate Transcarbamoylase Is Allosterically Inhibited by the End Product of Its Pathway 300

Allosterically regulated enzymes do not follow Michaelis–Menten kinetics 301

ATCase consists of separable catalytic and regulatory subunits 301

Allosteric interactions in ATCase are mediated by large changes in quaternary structure 302

Allosteric regulators modulate the T-to-R equilibrium 305

10.2 Isozymes Provide a Means of Regulation Specific to Distinct Tissues and Developmental Stages 306

10.3 Covalent Modification Is a Means of Regulating Enzyme Activity 307

Kinases and phosphatases control the extent of protein phosphorylation 308

Phosphorylation is a highly effective means of regulating the activities of target proteins 310

Cyclic AMP activates protein kinase A by altering the quaternary structure 311

ATP and the target protein bind to a deep cleft in the catalytic subunit of protein kinase A 312

10.4 Many Enzymes Are Activated by Specific Proteolytic Cleavage 312

Chymotrypsinogen is activated by specific cleavage of a single peptide bond 313

Proteolytic activation of chymotrypsinogen leads to the formation of a substrate-binding site 314

The generation of trypsin from trypsinogen leads to the activation of other zymogens 315

Some proteolytic enzymes have specific inhibitors 316

Blood clotting is accomplished by a cascade of zymogen activations 317

Fibrinogen is converted by thrombin into a fibrin clot 318

Prothrombin is readied for activation by a vitamin K-dependent modification 320

Hemophilia revealed an early step in clotting 321

The clotting process must be precisely regulated 321

Chapter 11 Carbohydrates 329

11.1 Monosaccharides Are the Simplest Carbohydrates 330

Many common sugars exist in cyclic forms 332

Pyranose and furanose rings can assume different conformations 334

Glucose is a reducing sugar	335	A membrane lipid is an amphipathic molecule containing a hydrophilic and a hydrophobic moiety	363
Monosaccharides are joined to alcohols and amines through glycosidic bonds	336	12.3 Phospholipids and Glycolipids Readily Form Bimolecular Sheets in Aqueous Media	364
Phosphorylated sugars are key intermediates in energy generation and biosyntheses	336	Lipid vesicles can be formed from phospholipids	365
11.2 Monosaccharides Are Linked to Form Complex Carbohydrates	337	Lipid bilayers are highly impermeable to ions and most polar molecules	366
Sucrose, lactose, and maltose are the common disaccharides	337	12.4 Proteins Carry Out Most Membrane Processes	367
Glycogen and starch are storage forms of glucose	338	Proteins associate with the lipid bilayer in a variety of ways	367
Cellulose, a structural component of plants, is made of chains of glucose	338	Proteins interact with membranes in a variety of ways	368
11.3 Carbohydrates Can Be Linked to Proteins to Form Glycoproteins	339	Some proteins associate with membranes through covalently attached hydrophobic groups	371
Carbohydrates can be linked to proteins through asparagine (<i>N</i> -linked) or through serine or threonine (<i>O</i> -linked) residues	340	Transmembrane helices can be accurately predicted from amino acid sequences	371
The glycoprotein erythropoietin is a vital hormone	340	12.5 Lipids and Many Membrane Proteins Diffuse Rapidly in the Plane of the Membrane	373
Proteoglycans, composed of polysaccharides and protein, have important structural roles	341	The fluid mosaic model allows lateral movement but not rotation through the membrane	374
Proteoglycans are important components of cartilage	342	Membrane fluidity is controlled by fatty acid composition and cholesterol content	374
Mucins are glycoprotein components of mucus	343	Lipid rafts are highly dynamic complexes formed between cholesterol and specific lipids	375
Protein glycosylation takes place in the lumen of the endoplasmic reticulum and in the Golgi complex	343	All biological membranes are asymmetric	375
Specific enzymes are responsible for oligosaccharide assembly	345	12.6 Eukaryotic Cells Contain Compartments Bounded by Internal Membranes	376
Blood groups are based on protein glycosylation patterns	345		
Errors in glycosylation can result in pathological conditions	346	Chapter 13 Membrane Channels and Pumps	383
Oligosaccharides can be “sequenced”	346	The expression of transporters largely defines the metabolic activities of a given cell type	384
11.4 Lectins Are Specific Carbohydrate-Binding Proteins	347	13.1 The Transport of Molecules Across a Membrane May Be Active or Passive	384
Lectins promote interactions between cells	348	Many molecules require protein transporters to cross membranes	384
Lectins are organized into different classes	348	Free energy stored in concentration gradients can be quantified	385
Influenza virus binds to sialic acid residues	349	13.2 Two Families of Membrane Proteins Use ATP Hydrolysis to Pump Ions and Molecules Across Membranes	386
Chapter 12 Lipids and Cell Membranes	357	P-type ATPases couple phosphorylation and conformational changes to pump calcium ions across membranes	386
Many common features underlie the diversity of biological membranes	358	Digitalis specifically inhibits the Na ⁺ –K ⁺ pump by blocking its dephosphorylation	389
12.1 Fatty Acids Are Key Constituents of Lipids	358	P-type ATPases are evolutionarily conserved and play a wide range of roles	390
Fatty acid names are based on their parent hydrocarbons	358	Multidrug resistance highlights a family of membrane pumps with ATP-binding cassette domains	390
Fatty acids vary in chain length and degree of unsaturation	359		
12.2 There Are Three Common Types of Membrane Lipids	360		
Phospholipids are the major class of membrane lipids	360		
Membrane lipids can include carbohydrate moieties	361		
Cholesterol is a lipid based on a steroid nucleus	362		
Archaeal membranes are built from ether lipids with branched chains	362		

13.3 Lactose Permease Is an Archetype of Secondary Transporters That Use One Concentration Gradient to Power the Formation of Another 392

13.4 Specific Channels Can Rapidly Transport Ions Across Membranes 394

Action potentials are mediated by transient changes in Na⁺ and K⁺ permeability 394

Patch-clamp conductance measurements reveal the activities of single channels 395

The structure of a potassium ion channel is an archetype for many ion-channel structures 395

The structure of the potassium ion channel reveals the basis of ion specificity 396

The structure of the potassium ion channel explains its rapid rate of transport 399

Voltage gating requires substantial conformational changes in specific ion-channel domains 399

A channel can be activated by occlusion of the pore: the ball-and-chain model 400

The acetylcholine receptor is an archetype for ligand-gated ion channels 401

Action potentials integrate the activities of several ion channels working in concert 402

Disruption of ion channels by mutations or chemicals can be potentially life threatening 404

13.5 Gap Junctions Allow Ions and Small Molecules to Flow Between Communicating Cells 405

13.6 Specific Channels Increase the Permeability of Some Membranes to Water 406

Chapter 14 Signal-Transduction Pathways 415

Signal transduction depends on molecular circuits 416

14.1 Heterotrimeric G Proteins Transmit Signals and Reset Themselves 417

Ligand binding to 7TM receptors leads to the activation of heterotrimeric G proteins 419

Activated G proteins transmit signals by binding to other proteins 420

Cyclic AMP stimulates the phosphorylation of many target proteins by activating protein kinase A 420

G proteins spontaneously reset themselves through GTP hydrolysis 421

Some 7TM receptors activate the phosphoinositide cascade 422

Calcium ion is a widely used second messenger 423

Calcium ion often activates the regulatory protein calmodulin 424

14.2 Insulin Signaling: Phosphorylation Cascades Are Central to Many Signal-Transduction Processes 425

The insulin receptor is a dimer that closes around a bound insulin molecule 426

Insulin binding results in the cross-phosphorylation and activation of the insulin receptor 426

The activated insulin-receptor kinase initiates a kinase cascade 426

Insulin signaling is terminated by the action of phosphatases 429

14.3 EGF Signaling: Signal-Transduction Pathways Are Poised to Respond 429

EGF binding results in the dimerization of the EGF receptor 429

The EGF receptor undergoes phosphorylation of its carboxyl-terminal tail 431

EGF signaling leads to the activation of Ras, a small G protein 431

Activated Ras initiates a protein kinase cascade 432

EGF signaling is terminated by protein phosphatases and the intrinsic GTPase activity of Ras 432

14.4 Many Elements Recur with Variation in Different Signal-Transduction Pathways 433

14.5 Defects in Signal-Transduction Pathways Can Lead to Cancer and Other Diseases 434

Monoclonal antibodies can be used to inhibit signal-transduction pathways activated in tumors 434

Protein kinase inhibitors can be effective anticancer drugs 435

Cholera and whooping cough are due to altered G-protein activity 435

Part II TRANSDUCING AND STORING ENERGY

Chapter 15 Metabolism: Basic Concepts and Design 443

15.1 Metabolism Is Composed of Many Coupled, Interconnecting Reactions 444

Metabolism consists of energy-yielding and energy-requiring reactions 444

A thermodynamically unfavorable reaction can be driven by a favorable reaction 445

15.2 ATP Is the Universal Currency of Free Energy in Biological Systems 446

ATP hydrolysis is exergonic 446

ATP hydrolysis drives metabolism by shifting the equilibrium of coupled reactions 447

The high phosphoryl potential of ATP results from structural differences between ATP and its hydrolysis products 449

Phosphoryl-transfer potential is an important form of cellular energy transformation 450

15.3 The Oxidation of Carbon Fuels Is an Important Source of Cellular Energy	451	Many adults are intolerant of milk because they are deficient in lactase	487
Compounds with high phosphoryl-transfer potential can couple carbon oxidation to ATP synthesis	452	Galactose is highly toxic if the transferase is missing	488
Ion gradients across membranes provide an important form of cellular energy that can be coupled to ATP synthesis	453	16.2 The Glycolytic Pathway Is Tightly Controlled	488
Energy from foodstuffs is extracted in three stages	453	Glycolysis in muscle is regulated to meet the need for ATP	489
15.4 Metabolic Pathways Contain Many Recurring Motifs	454	The regulation of glycolysis in the liver illustrates the biochemical versatility of the liver	490
Activated carriers exemplify the modular design and economy of metabolism	454	A family of transporters enables glucose to enter and leave animal cells	493
Many activated carriers are derived from vitamins	457	Cancer and exercise training affect glycolysis in a similar fashion	494
Key reactions are reiterated throughout metabolism	459	16.3 Glucose Can Be Synthesized from Noncarbohydrate Precursors	495
Metabolic processes are regulated in three principal ways	461	Gluconeogenesis is not a reversal of glycolysis	497
Aspects of metabolism may have evolved from an RNA world	463	The conversion of pyruvate into phosphoenolpyruvate begins with the formation of oxaloacetate	498
<hr/> Chapter 16 Glycolysis and Gluconeogenesis	469	Oxaloacetate is shuttled into the cytoplasm and converted into phosphoenolpyruvate	499
Glucose is generated from dietary carbohydrates	470	The conversion of fructose 1,6-bisphosphate into fructose 6-phosphate and orthophosphate is an irreversible step	500
Glucose is an important fuel for most organisms	471	The generation of free glucose is an important control point	500
16.1 Glycolysis Is an Energy-Conversion Pathway in Many Organisms	471	Six high-transfer-potential phosphoryl groups are spent in synthesizing glucose from pyruvate	501
Hexokinase traps glucose in the cell and begins glycolysis	471	16.4 Gluconeogenesis and Glycolysis Are Reciprocally Regulated	502
Fructose 1,6-bisphosphate is generated from glucose 6-phosphate	473	Energy charge determines whether glycolysis or gluconeogenesis will be most active	502
The six-carbon sugar is cleaved into two three-carbon fragments	474	The balance between glycolysis and gluconeogenesis in the liver is sensitive to blood-glucose concentration	503
Mechanism: Triose phosphate isomerase salvages a three-carbon fragment	475	Substrate cycles amplify metabolic signals and produce heat	505
The oxidation of an aldehyde to an acid powers the formation of a compound with high phosphoryl-transfer potential	476	Lactate and alanine formed by contracting muscle are used by other organs	505
Mechanism: Phosphorylation is coupled to the oxidation of glyceraldehyde 3-phosphate by a thioester intermediate	478	Glycolysis and gluconeogenesis are evolutionarily intertwined	507
ATP is formed by phosphoryl transfer from 1,3-bisphosphoglycerate	479	<hr/> Chapter 17 The Citric Acid Cycle	515
Additional ATP is generated with the formation of pyruvate	480	The citric acid cycle harvests high-energy electrons	516
Two ATP molecules are formed in the conversion of glucose into pyruvate	481	17.1 Pyruvate Dehydrogenase Links Glycolysis to the Citric Acid Cycle	517
NAD ⁺ is regenerated from the metabolism of pyruvate	482	Mechanism: The synthesis of acetyl coenzyme A from pyruvate requires three enzymes and five coenzymes	518
Fermentations provide usable energy in the absence of oxygen	484	Flexible linkages allow lipoamide to move between different active sites	520
The binding site for NAD ⁺ is similar in many dehydrogenases	485	17.2 The Citric Acid Cycle Oxidizes Two-Carbon Units	521
Fructose and galactose are converted into glycolytic intermediates	485	Citrate synthase forms citrate from oxaloacetate and acetyl coenzyme A	522

Mechanism: The mechanism of citrate synthase prevents undesirable reactions	522	Ubiquinol is the entry point for electrons from FADH ₂ of flavoproteins	553
Citrate is isomerized into isocitrate	524	Electrons flow from ubiquinol to cytochrome <i>c</i> through Q-cytochrome <i>c</i> oxidoreductase	553
Isocitrate is oxidized and decarboxylated to alpha-ketoglutarate	524	The Q cycle funnels electrons from a two-electron carrier to a one-electron carrier and pumps protons	554
Succinyl coenzyme A is formed by the oxidative decarboxylation of alpha-ketoglutarate	525	Cytochrome <i>c</i> oxidase catalyzes the reduction of molecular oxygen to water	555
A compound with high phosphoryl-transfer potential is generated from succinyl coenzyme A	525	Toxic derivatives of molecular oxygen such as superoxide radical are scavenged by protective enzymes	558
Mechanism: Succinyl coenzyme A synthetase transforms types of biochemical energy	526	Electrons can be transferred between groups that are not in contact	560
Oxaloacetate is regenerated by the oxidation of succinate	527	The conformation of cytochrome <i>c</i> has remained essentially constant for more than a billion years	561
The citric acid cycle produces high-transfer-potential electrons, ATP, and CO ₂	528	18.4 A Proton Gradient Powers the Synthesis of ATP	561
17.3 Entry to the Citric Acid Cycle and Metabolism Through It Are Controlled	530	ATP synthase is composed of a proton-conducting unit and a catalytic unit	563
The pyruvate dehydrogenase complex is regulated allosterically and by reversible phosphorylation	531	Proton flow through ATP synthase leads to the release of tightly bound ATP: The binding-change mechanism	564
The citric acid cycle is controlled at several points	532	Rotational catalysis is the world's smallest molecular motor	565
Defects in the citric acid cycle contribute to the development of cancer	533	Proton flow around the <i>c</i> ring powers ATP synthesis	566
17.4 The Citric Acid Cycle Is a Source of Biosynthetic Precursors	534	ATP synthase and G proteins have several common features	568
The citric acid cycle must be capable of being rapidly replenished	534	18.5 Many Shuttles Allow Movement Across Mitochondrial Membranes	568
The disruption of pyruvate metabolism is the cause of beriberi and poisoning by mercury and arsenic	535	Electrons from cytoplasmic NADH enter mitochondria by shuttles	569
The citric acid cycle may have evolved from preexisting pathways	536	The entry of ADP into mitochondria is coupled to the exit of ATP by ATP-ADP translocase	570
17.5 The Glyoxylate Cycle Enables Plants and Bacteria to Grow on Acetate	536	Mitochondrial transporters for metabolites have a common tripartite structure	571
Chapter 18 Oxidative Phosphorylation	543	18.6 The Regulation of Cellular Respiration Is Governed Primarily by the Need for ATP	572
18.1 Eukaryotic Oxidative Phosphorylation Takes Place in Mitochondria	544	The complete oxidation of glucose yields about 30 molecules of ATP	572
Mitochondria are bounded by a double membrane	544	The rate of oxidative phosphorylation is determined by the need for ATP	573
Mitochondria are the result of an endosymbiotic event	545	Regulated uncoupling leads to the generation of heat	574
18.2 Oxidative Phosphorylation Depends on Electron Transfer	546	Oxidative phosphorylation can be inhibited at many stages	576
The electron-transfer potential of an electron is measured as redox potential	546	Mitochondrial diseases are being discovered	576
A 1.14-volt potential difference between NADH and molecular oxygen drives electron transport through the chain and favors the formation of a proton gradient	548	Mitochondria play a key role in apoptosis	577
18.3 The Respiratory Chain Consists of Four Complexes: Three Proton Pumps and a Physical Link to the Citric Acid Cycle	549	Power transmission by proton gradients is a central motif of bioenergetics	577
The high-potential electrons of NADH enter the respiratory chain at NADH-Q oxidoreductase	551	Chapter 19 The Light Reactions of Photosynthesis	585
		Photosynthesis converts light energy into chemical energy	586
		19.1 Photosynthesis Takes Place in Chloroplasts	587
		The primary events of photosynthesis take place in thylakoid membranes	587
		Chloroplasts arose from an endosymbiotic event	588

19.2 Light Absorption by Chlorophyll Induces Electron Transfer	588	20.2 The Activity of the Calvin Cycle Depends on Environmental Conditions	617
A special pair of chlorophylls initiate charge separation	589	Rubisco is activated by light-driven changes in proton and magnesium ion concentrations	618
Cyclic electron flow reduces the cytochrome of the reaction center	592	Thioredoxin plays a key role in regulating the Calvin cycle	618
19.3 Two Photosystems Generate a Proton Gradient and NADPH in Oxygenic Photosynthesis	592	The C ₄ pathway of tropical plants accelerates photosynthesis by concentrating carbon dioxide	619
Photosystem II transfers electrons from water to plastoquinone and generates a proton gradient	592	Crassulacean acid metabolism permits growth in arid ecosystems	620
Cytochrome <i>bf</i> links photosystem II to photosystem I	595	20.3 The Pentose Phosphate Pathway Generates NADPH and Synthesizes Five-Carbon Sugars	621
Photosystem I uses light energy to generate reduced ferredoxin, a powerful reductant	595	Two molecules of NADPH are generated in the conversion of glucose 6-phosphate into ribulose 5-phosphate	621
Ferredoxin–NADP ⁺ reductase converts NADP ⁺ into NADPH	596	The pentose phosphate pathway and glycolysis are linked by transketolase and transaldolase	621
19.4 A Proton Gradient Across the Thylakoid Membrane Drives ATP Synthesis	597	Mechanism: Transketolase and transaldolase stabilize carbanionic intermediates by different mechanisms	624
The ATP synthase of chloroplasts closely resembles those of mitochondria and prokaryotes	598	20.4 The Metabolism of Glucose 6-phosphate by the Pentose Phosphate Pathway Is Coordinated with Glycolysis	626
Cyclic electron flow through photosystem I leads to the production of ATP instead of NADPH	599	The rate of the pentose phosphate pathway is controlled by the level of NADP ⁺	626
The absorption of eight photons yields one O ₂ , two NADPH, and three ATP molecules	600	The flow of glucose 6-phosphate depends on the need for NADPH, ribose 5-phosphate, and ATP	627
19.5 Accessory Pigments Funnel Energy into Reaction Centers	601	Through the looking-glass: The Calvin cycle and the pentose phosphate pathway are mirror images	629
Resonance energy transfer allows energy to move from the site of initial absorbance to the reaction center	601	20.5 Glucose 6-phosphate Dehydrogenase Plays a Key Role in Protection Against Reactive Oxygen Species	629
Light-harvesting complexes contain additional chlorophylls and carotenoids	602	Glucose 6-phosphate dehydrogenase deficiency causes a drug-induced hemolytic anemia	629
The components of photosynthesis are highly organized	603	A deficiency of glucose 6-phosphate dehydrogenase confers an evolutionary advantage in some circumstances	631
Many herbicides inhibit the light reactions of photosynthesis	604	Chapter 20 The Calvin Cycle and Pentose Phosphate Pathway	609
19.6 The Ability to Convert Light into Chemical Energy Is Ancient	604	20.1 The Calvin Cycle Synthesizes Hexoses from Carbon Dioxide and Water	610
		Carbon dioxide reacts with ribulose 1,5-bisphosphate to form two molecules of 3-phosphoglycerate	611
		Rubisco activity depends on magnesium and carbamate	612
		Rubisco also catalyzes a wasteful oxygenase reaction: Catalytic imperfection	613
		Hexose phosphates are made from phosphoglycerate, and ribulose 1,5-bisphosphate is regenerated	614
		Three ATP and two NADPH molecules are used to bring carbon dioxide to the level of a hexose	617
		Starch and sucrose are the major carbohydrate stores in plants	617
		Chapter 21 Glycogen Metabolism	637
		Glycogen metabolism is the regulated release and storage of glucose	638
		21.1 Glycogen Breakdown Requires the Interplay of Several Enzymes	639
		Phosphorylase catalyzes the phosphorolytic cleavage of glycogen to release glucose 1-phosphate	639
		Mechanism: Pyridoxal phosphate participates in the phosphorolytic cleavage of glycogen	640
		A debranching enzyme also is needed for the breakdown of glycogen	641
		Phosphoglucomutase converts glucose 1-phosphate into glucose 6-phosphate	642
		The liver contains glucose 6-phosphatase, a hydrolytic enzyme absent from muscle	643

21.2 Phosphorylase Is Regulated by Allosteric Interactions and Reversible Phosphorylation 643

Muscle phosphorylase is regulated by the intracellular energy charge 643

Liver phosphorylase produces glucose for use by other tissues 645

Phosphorylase kinase is activated by phosphorylation and calcium ions 645

21.3 Epinephrine and Glucagon Signal the Need for Glycogen Breakdown 646

G proteins transmit the signal for the initiation of glycogen breakdown 646

Glycogen breakdown must be rapidly turned off when necessary 648

The regulation of glycogen phosphorylase became more sophisticated as the enzyme evolved 649

21.4 Glycogen Is Synthesized and Degraded by Different Pathways 649

UDP-glucose is an activated form of glucose 649

Glycogen synthase catalyzes the transfer of glucose from UDP-glucose to a growing chain 650

A branching enzyme forms α -1,6 linkages 651

Glycogen synthase is the key regulatory enzyme in glycogen synthesis 651

Glycogen is an efficient storage form of glucose 651

21.5 Glycogen Breakdown and Synthesis Are Reciprocally Regulated 652

Protein phosphatase 1 reverses the regulatory effects of kinases on glycogen metabolism 653

Insulin stimulates glycogen synthesis by inactivating glycogen synthase kinase 654

Glycogen metabolism in the liver regulates the blood-glucose level 655

A biochemical understanding of glycogen-storage diseases is possible 656

Chapter 22 Fatty Acid Metabolism 663

Fatty acid degradation and synthesis mirror each other in their chemical reactions 664

22.1 Triacylglycerols Are Highly Concentrated Energy Stores 665

Dietary lipids are digested by pancreatic lipases 665

Dietary lipids are transported in chylomicrons 666

22.2 The Use of Fatty Acids As Fuel Requires Three Stages of Processing 667

Triacylglycerols are hydrolyzed by hormone-stimulated lipases 667

Fatty acids are linked to coenzyme A before they are oxidized 668

Carnitine carries long-chain activated fatty acids into the mitochondrial matrix 669

Acetyl CoA, NADH, and FADH₂ are generated in each round of fatty acid oxidation 670

The complete oxidation of palmitate yields 106 molecules of ATP 671

22.3 Unsaturated and Odd-Chain Fatty Acids Require Additional Steps for Degradation 672

An isomerase and a reductase are required for the oxidation of unsaturated fatty acids 672

Odd-chain fatty acids yield propionyl CoA in the final thiolysis step 673

Vitamin B₁₂ contains a corrin ring and a cobalt atom 674

Mechanism: Methylmalonyl CoA mutase catalyzes a rearrangement to form succinyl CoA 675

Fatty acids are also oxidized in peroxisomes 676

Ketone bodies are formed from acetyl CoA when fat breakdown predominates 677

Ketone bodies are a major fuel in some tissues 678

Animals cannot convert fatty acids into glucose 680

22.4 Fatty Acids Are Synthesized by Fatty Acid Synthase 680

Fatty acids are synthesized and degraded by different pathways 680

The formation of malonyl CoA is the committed step in fatty acid synthesis 681

Intermediates in fatty acid synthesis are attached to an acyl carrier protein 681

Fatty acid synthesis consists of a series of condensation, reduction, dehydration, and reduction reactions 682

Fatty acids are synthesized by a multifunctional enzyme complex in animals 683

The synthesis of palmitate requires 8 molecules of acetyl CoA, 14 molecules of NADPH, and 7 molecules of ATP 685

Citrate carries acetyl groups from mitochondria to the cytoplasm for fatty acid synthesis 686

Several sources supply NADPH for fatty acid synthesis 686

Fatty acid synthase inhibitors may be useful drugs 687

22.5 The Elongation and Unsaturation of Fatty Acids Are Accomplished by Accessory Enzyme Systems 687

Membrane-bound enzymes generate unsaturated fatty acids 688

Eicosanoid hormones are derived from polyunsaturated fatty acids 688

22.6 Acetyl CoA Carboxylase Plays a Key Role in Controlling Fatty Acid Metabolism 690

Acetyl CoA carboxylase is regulated by conditions in the cell 690

Acetyl CoA carboxylase is regulated by a variety of hormones 690

Chapter 23 Protein Turnover and Amino Acid Catabolism 697

23.1 Proteins Are Degraded to Amino Acids 698

The digestion of dietary proteins begins in the stomach and is completed in the intestine 698

Cellular proteins are degraded at different rates 699

23.2 Protein Turnover Is Tightly Regulated	699
Ubiquitin tags proteins for destruction	699
The proteasome digests the ubiquitin-tagged proteins	701
The ubiquitin pathway and the proteasome have prokaryotic counterparts	701
Protein degradation can be used to regulate biological function	702
23.3 The First Step in Amino Acid Degradation Is the Removal of Nitrogen	704
Alpha-amino groups are converted into ammonium ions by the oxidative deamination of glutamate	704
Mechanism: Pyridoxal phosphate forms Schiff-base intermediates in aminotransferases	705
Aspartate aminotransferase is an archetypal pyridoxal-dependent transaminase	706
Pyridoxal phosphate enzymes catalyze a wide array of reactions	707
Serine and threonine can be directly deaminated	708
Peripheral tissues transport nitrogen to the liver	708
23.4 Ammonium Ion Is Converted into Urea in Most Terrestrial Vertebrates	709
The urea cycle begins with the formation of carbamoyl phosphate	709
The urea cycle is linked to gluconeogenesis	711
Urea-cycle enzymes are evolutionarily related to enzymes in other metabolic pathways	712
Inherited defects of the urea cycle cause hyperammonemia and can lead to brain damage	712
Urea is not the only means of disposing of excess nitrogen	713
23.5 Carbon Atoms of Degraded Amino Acids Emerge As Major Metabolic Intermediates	714
Pyruvate is an entry point into metabolism for a number of amino acids	715
Oxaloacetate is an entry point into metabolism for aspartate and asparagine	716
Alpha-ketoglutarate is an entry point into metabolism for five-carbon amino acids	716
Succinyl coenzyme A is a point of entry for several nonpolar amino acids	717
Methionine degradation requires the formation of a key methyl donor, S-adenosylmethionine	717
The branched-chain amino acids yield acetyl CoA, acetoacetate, or propionyl CoA	717
Oxygenases are required for the degradation of aromatic amino acids	719
23.6 Inborn Errors of Metabolism Can Disrupt Amino Acid Degradation	721

Part III SYNTHESIZING THE MOLECULES OF LIFE

Chapter 24 The Biosynthesis of Amino Acids	729
Amino acid synthesis requires solutions to three key biochemical problems	730
24.1 Nitrogen Fixation: Microorganisms Use ATP and a Powerful Reductant to Reduce Atmospheric Nitrogen to Ammonia	730
The iron–molybdenum cofactor of nitrogenase binds and reduces atmospheric nitrogen	731
Ammonium ion is assimilated into an amino acid through glutamate and glutamine	733
24.2 Amino Acids Are Made from Intermediates of the Citric Acid Cycle and Other Major Pathways	735
Human beings can synthesize some amino acids but must obtain others from the diet	735
Aspartate, alanine, and glutamate are formed by the addition of an amino group to an alpha-ketoacid	736
A common step determines the chirality of all amino acids	737
The formation of asparagine from aspartate requires an adenylated intermediate	737
Glutamate is the precursor of glutamine, proline, and arginine	738
3-Phosphoglycerate is the precursor of serine, cysteine, and glycine	738
Tetrahydrofolate carries activated one-carbon units at several oxidation levels	739
S-Adenosylmethionine is the major donor of methyl groups	740
Cysteine is synthesized from serine and homocysteine	742
High homocysteine levels correlate with vascular disease	743
Shikimate and chorismate are intermediates in the biosynthesis of aromatic amino acids	743
Tryptophan synthase illustrates substrate channeling in enzymatic catalysis	746
24.3 Feedback Inhibition Regulates Amino Acid Biosynthesis	747
Branched pathways require sophisticated regulation	747
An enzymatic cascade modulates the activity of glutamine synthetase	749
24.4 Amino Acids Are Precursors of Many Biomolecules	750
Glutathione, a gamma-glutamyl peptide, serves as a sulfhydryl buffer and an antioxidant	751
Nitric oxide, a short-lived signal molecule, is formed from arginine	751

Porphyrins are synthesized from glycine and succinyl coenzyme A	752	The synthesis of purine nucleotides is controlled by feedback inhibition at several sites	777
Porphyrins accumulate in some inherited disorders of porphyrin metabolism	754	The synthesis of deoxyribonucleotides is controlled by the regulation of ribonucleotide reductase	778
Chapter 25 Nucleotide Biosynthesis	761	25.5 Disruptions in Nucleotide Metabolism Can Cause Pathological Conditions	778
Nucleotides can be synthesized by de novo or salvage pathways	762	The loss of adenosine deaminase activity results in severe combined immunodeficiency	778
25.1 The Pyrimidine Ring Is Assembled de Novo or Recovered by Salvage Pathways	763	Gout is induced by high serum levels of urate	779
Bicarbonate and other oxygenated carbon compounds are activated by phosphorylation	763	Lesch–Nyhan syndrome is a dramatic consequence of mutations in a salvage-pathway enzyme	780
The side chain of glutamine can be hydrolyzed to generate ammonia	763	Folic acid deficiency promotes birth defects such as spina bifida	781
Intermediates can move between active sites by channeling	763		
Orotate acquires a ribose ring from PRPP to form a pyrimidine nucleotide and is converted into uridylate	764	Chapter 26 The Biosynthesis of Membrane Lipids and Steroids	787
Nucleotide mono-, di-, and triphosphates are interconvertible	765	26.1 Phosphatidate Is a Common Intermediate in the Synthesis of Phospholipids and Triacylglycerols	788
CTP is formed by amination of UTP	765	The synthesis of phospholipids requires an activated intermediate	789
Salvage pathways recycle pyrimidine bases	766	Sphingolipids are synthesized from ceramide	791
25.2 Purine Bases Can Be Synthesized de Novo or Recycled by Salvage Pathways	766	Gangliosides are carbohydrate-rich sphingolipids that contain acidic sugars	792
The purine ring system is assembled on ribose phosphate	766	Sphingolipids confer diversity on lipid structure and function	793
The purine ring is assembled by successive steps of activation by phosphorylation followed by displacement	767	Respiratory distress syndrome and Tay–Sachs disease result from the disruption of lipid metabolism	793
AMP and GMP are formed from IMP	769	Phosphatidic acid phosphatase is a key regulatory enzyme in lipid metabolism	794
Enzymes of the purine synthesis pathway associate with one another in vivo	770	26.2 Cholesterol Is Synthesized from Acetyl Coenzyme A in Three Stages	795
Salvage pathways economize intracellular energy expenditure	770	The synthesis of mevalonate, which is activated as isopentenyl pyrophosphate, initiates the synthesis of cholesterol	795
25.3 Deoxyribonucleotides Are Synthesized by the Reduction of Ribonucleotides Through a Radical Mechanism	771	Squalene (C ₃₀) is synthesized from six molecules of isopentenyl pyrophosphate (C ₅)	796
Mechanism: A tyrosyl radical is critical to the action of ribonucleotide reductase	771	Squalene cyclizes to form cholesterol	797
Stable radicals other than tyrosyl radical are employed by other ribonucleotide reductases	773	26.3 The Complex Regulation of Cholesterol Biosynthesis Takes Place at Several Levels	798
Thymidylate is formed by the methylation of deoxyuridylate	774	Lipoproteins transport cholesterol and triacylglycerols throughout the organism	801
Dihydrofolate reductase catalyzes the regeneration of tetrahydrofolate, a one-carbon carrier	775	The blood levels of certain lipoproteins can serve diagnostic purposes	802
Several valuable anticancer drugs block the synthesis of thymidylate	775	Low-density lipoproteins play a central role in cholesterol metabolism	803
25.4 Key Steps in Nucleotide Biosynthesis Are Regulated by Feedback Inhibition	776	The absence of the LDL receptor leads to hypercholesterolemia and atherosclerosis	804
Pyrimidine biosynthesis is regulated by aspartate transcarbamoylase	777	Mutations in the LDL receptor prevent LDL release and result in receptor destruction	805

HDL appears to protect against arteriosclerosis	806	Metabolic adaptations in prolonged starvation minimize protein degradation	838
The clinical management of cholesterol levels can be understood at a biochemical level	807	27.6 Ethanol Alters Energy Metabolism in the Liver	840
26.4 Important Derivatives of Cholesterol Include Bile Salts and Steroid Hormones	807	Ethanol metabolism leads to an excess of NADH	840
Letters identify the steroid rings and numbers identify the carbon atoms	809	Excess ethanol consumption disrupts vitamin metabolism	842
Steroids are hydroxylated by cytochrome P450 monooxygenases that use NADPH and O ₂	809	Chapter 28 DNA Replication, Repair, and Recombination	849
The cytochrome P450 system is widespread and performs a protective function	810	28.1 DNA Replication Proceeds by the Polymerization of Deoxyribonucleoside Triphosphates Along a Template	850
Pregnenolone, a precursor of many other steroids, is formed from cholesterol by cleavage of its side chain	811	DNA polymerases require a template and a primer	850
Pregesterone and corticosteroids are synthesized from pregnenolone	811	All DNA polymerases have structural features in common	851
Androgens and estrogens are synthesized from pregnenolone	812	Two bound metal ions participate in the polymerase reaction	851
Vitamin D is derived from cholesterol by the ring-splitting activity of light	813	The specificity of replication is dictated by complementarity of shape between bases	852
Chapter 27 The Integration of Metabolism	821	An RNA primer synthesized by primase enables DNA synthesis to begin	853
27.1 Caloric Homeostasis Is a Means of Regulating Body Weight	822	One strand of DNA is made continuously, whereas the other strand is synthesized in fragments	853
27.2 The Brain Plays a Key Role in Caloric Homeostasis	824	DNA ligase joins ends of DNA in duplex regions	854
Signals from the gastrointestinal tract induce feelings of satiety	824	The separation of DNA strands requires specific helicases and ATP hydrolysis	854
Leptin and insulin regulate long-term control over caloric homeostasis	825	28.2 DNA Unwinding and Supercoiling Are Controlled by Topoisomerases	855
Leptin is one of several hormones secreted by adipose tissue	826	The linking number of DNA, a topological property, determines the degree of supercoiling	856
Leptin resistance may be a contributing factor to obesity	827	Topoisomerases prepare the double helix for unwinding	858
Dieting is used to combat obesity	827	Type I topoisomerases relax supercoiled structures	858
27.3 Diabetes Is a Common Metabolic Disease Often Resulting from Obesity	828	Type II topoisomerases can introduce negative supercoils through coupling to ATP hydrolysis	859
Insulin initiates a complex signal-transduction pathway in muscle	828	28.3 DNA Replication Is Highly Coordinated	861
Metabolic syndrome often precedes type 2 diabetes	830	DNA replication requires highly processive polymerases	861
Excess fatty acids in muscle modify metabolism	830	The leading and lagging strands are synthesized in a coordinated fashion	862
Insulin resistance in muscle facilitates pancreatic failure	831	DNA replication in <i>Escherichia coli</i> begins at a unique site	864
Metabolic derangements in type 1 diabetes result from insulin insufficiency and glucagon excess	832	DNA synthesis in eukaryotes is initiated at multiple sites	865
27.4 Exercise Beneficially Alters the Biochemistry of Cells	833	Telomeres are unique structures at the ends of linear chromosomes	866
Mitochondrial biogenesis is stimulated by muscular activity	834	Telomeres are replicated by telomerase, a specialized polymerase that carries its own RNA template	867
Fuel choice during exercise is determined by the intensity and duration of activity	835	28.4 Many Types of DNA Damage Can Be Repaired	867
27.5 Food Intake and Starvation Induce Metabolic Changes	836	Errors can arise in DNA replication	867
The starved–fed cycle is the physiological response to a fast	837	Bases can be damaged by oxidizing agents, alkylating agents, and light	868

DNA damage can be detected and repaired by a variety of systems	869	Enhancer sequences can stimulate transcription at start sites thousands of bases away	900
The presence of thymine instead of uracil in DNA permits the repair of deaminated cytosine	871	29.3 The Transcription Products of Eukaryotic Polymerases Are Processed	901
Some genetic diseases are caused by the expansion of repeats of three nucleotides	872	RNA polymerase I produces three ribosomal RNAs	901
Many cancers are caused by the defective repair of DNA	872	RNA polymerase III produces transfer RNA	902
Many potential carcinogens can be detected by their mutagenic action on bacteria	873	The product of RNA polymerase II, the pre-mRNA transcript, acquires a 5' cap and a 3' poly(A) tail	902
28.5 DNA Recombination Plays Important Roles in Replication, Repair, and Other Processes	874	Small regulatory RNAs are cleaved from larger precursors	904
RecA can initiate recombination by promoting strand invasion	874	RNA editing changes the proteins encoded by mRNA	904
Some recombination reactions proceed through Holliday-junction intermediates	875	Sequences at the ends of introns specify splice sites in mRNA precursors	905
<hr/> Chapter 29 RNA Synthesis and Processing	883	Splicing consists of two sequential transesterification reactions	906
RNA synthesis comprises three stages: Initiation, elongation, and termination	884	Small nuclear RNAs in spliceosomes catalyze the splicing of mRNA precursors	907
29.1 RNA Polymerases Catalyze Transcription	885	Transcription and processing of mRNA are coupled	909
RNA chains are formed de novo and grow in the 5'-to-3' direction	886	Mutations that affect pre-mRNA splicing cause disease	909
RNA polymerases backtrack and correct errors	888	Most human pre-mRNAs can be spliced in alternative ways to yield different proteins	910
RNA polymerase binds to promoter sites on the DNA template to initiate transcription	888	29.4 The Discovery of Catalytic RNA Was Revealing in Regard to Both Mechanism and Evolution	911
Sigma subunits of RNA polymerase recognize promoter sites	889	<hr/> Chapter 30 Protein Synthesis	921
RNA polymerases must unwind the template double helix for transcription to take place	890	30.1 Protein Synthesis Requires the Translation of Nucleotide Sequences into Amino Acid Sequences	922
Elongation takes place at transcription bubbles that move along the DNA template	890	The synthesis of long proteins requires a low error frequency	922
Sequences within the newly transcribed RNA signal termination	891	Transfer RNA molecules have a common design	923
Some messenger RNAs directly sense metabolite concentrations	892	Some transfer RNA molecules recognize more than one codon because of wobble in base-pairing	925
The <i>rho</i> protein helps to terminate the transcription of some genes	892	30.2 Aminoacyl Transfer RNA Synthetases Read the Genetic Code	927
Some antibiotics inhibit transcription	893	Amino acids are first activated by adenylation	927
Precursors of transfer and ribosomal RNA are cleaved and chemically modified after transcription in prokaryotes	895	Aminoacyl-tRNA synthetases have highly discriminating amino acid activation sites	928
29.2 Transcription in Eukaryotes Is Highly Regulated	896	Proofreading by aminoacyl-tRNA synthetases increases the fidelity of protein synthesis	929
Three types of RNA polymerase synthesize RNA in eukaryotic cells	897	Synthetases recognize various features of transfer RNA molecules	930
Three common elements can be found in the RNA polymerase II promoter region	898	Aminoacyl-tRNA synthetases can be divided into two classes	931
The TFIID protein complex initiates the assembly of the active transcription complex	899	30.3 The Ribosome Is the Site of Protein Synthesis	931
Multiple transcription factors interact with eukaryotic promoters	900	Ribosomal RNAs (5S, 16S, and 23S rRNA) play a central role in protein synthesis	932
		Ribosomes have three tRNA-binding sites that bridge the 30s and 50s subunits	934

The start signal is usually AUG preceded by several bases that pair with 16S rRNA	934	31.3 Regulatory Circuits Can Result in Switching Between Patterns of Gene Expression	964
Bacterial protein synthesis is initiated by formylmethionyl transfer RNA	935	Lambda repressor regulates its own expression	964
Formylmethionyl-tRNA _f is placed in the P site of the ribosome in the formation of the 70S initiation complex	936	A circuit based on lambda repressor and Cro form a genetic switch	965
Elongation factors deliver aminoacyl-tRNA to the ribosome	936	Many prokaryotic cells release chemical signals that regulate gene expression in other cells	965
Peptidyl transferase catalyzes peptide-bond synthesis	937	Biofilms are complex communities of prokaryotes	966
The formation of a peptide bond is followed by the GTP-driven translocation of tRNAs and mRNA	938	31.4 Gene Expression Can Be Controlled at Posttranscriptional Levels	967
Protein synthesis is terminated by release factors that read stop codons	940	Attenuation is a prokaryotic mechanism for regulating transcription through the modulation of nascent RNA secondary structure	967
30.4 Eukaryotic Protein Synthesis Differs from Prokaryotic Protein Synthesis Primarily in Translation Initiation	941	<hr/> Chapter 32 The Control of Gene Expression in Eukaryotes	973
Mutations in initiation factor 2 cause a curious pathological condition	942	32.1 Eukaryotic DNA Is Organized into Chromatin	974
30.5 A Variety of Antibiotics and Toxins Can Inhibit Protein Synthesis	943	Nucleosomes are complexes of DNA and histones	975
Some antibiotics inhibit protein synthesis	943	DNA wraps around histone octamers to form nucleosomes	975
Diphtheria toxin blocks protein synthesis in eukaryotes by inhibiting translocation	944	32.2 Transcription Factors Bind DNA and Regulate Transcription Initiation	977
Ricin fatally modifies 28S ribosomal RNA	945	A range of DNA-binding structures are employed by eukaryotic DNA-binding proteins	977
30.6 Ribosomes Bound to the Endoplasmic Reticulum Manufacture Secretory and Membrane Proteins	945	Activation domains interact with other proteins	978
Signal sequences mark proteins for translocation across the endoplasmic reticulum membrane	945	Multiple transcription factors interact with eukaryotic regulatory regions	979
Transport vesicles carry cargo proteins to their final destination	947	Enhancers can stimulate transcription in specific cell types	979
<hr/> Chapter 31 The Control of Gene Expression in Prokaryotes	957	Induced pluripotent stem cells can be generated by introducing four transcription factors into differentiated cells	980
31.1 Many DNA-Binding Proteins Recognize Specific DNA Sequences	958	32.3 The Control of Gene Expression Can Require Chromatin Remodeling	980
The helix-turn-helix motif is common to many prokaryotic DNA-binding proteins	959	The methylation of DNA can alter patterns of gene expression	981
31.2 Prokaryotic DNA-Binding Proteins Bind Specifically to Regulatory Sites in Operons	959	Steroids and related hydrophobic molecules pass through membranes and bind to DNA-binding receptors	982
An operon consists of regulatory elements and protein-encoding genes	960	Nuclear hormone receptors regulate transcription by recruiting coactivators to the transcription complex	982
The <i>lac</i> repressor protein in the absence of lactose binds to the operator and blocks transcription	961	Steroid-hormone receptors are targets for drugs	984
Ligand binding can induce structural changes in regulatory proteins	962	Chromatin structure is modulated through covalent modifications of histone tails	985
The operon is a common regulatory unit in prokaryotes	962	Histone deacetylases contribute to transcriptional repression	986
Transcription can be stimulated by proteins that contact RNA polymerase	963	32.4 Eukaryotic Gene Expression Can Be Controlled at Posttranscriptional Levels	987
		Genes associated with iron metabolism are translationally regulated in animals	987
		Small RNAs regulate the expression of many eukaryotic genes	989

Part IV RESPONDING TO ENVIRONMENTAL CHANGES

Chapter 33 Sensory Systems

995

33.1 A Wide Variety of Organic Compounds Are Detected by Olfaction

996

Olfaction is mediated by an enormous family of seven-transmembrane-helix receptors

996

Odorants are decoded by a combinatorial mechanism

998

33.2 Taste Is a Combination of Senses That Function by Different Mechanisms

1000

Sequencing of the human genome led to the discovery of a large family of 7TM bitter receptors

1001

A heterodimeric 7TM receptor responds to sweet compounds

1002

Umami, the taste of glutamate and aspartate, is mediated by a heterodimeric receptor related to the sweet receptor

1003

Salty tastes are detected primarily by the passage of sodium ions through channels

1003

Sour tastes arise from the effects of hydrogen ions (acids) on channels

1003

33.3 Photoreceptor Molecules in the Eye Detect Visible Light

1004

Rhodopsin, a specialized 7TM receptor, absorbs visible light

1004

Light absorption induces a specific isomerization of bound 11-*cis*-retinal

1005

Light-induced lowering of the calcium level coordinates recovery

1006

Color vision is mediated by three cone receptors that are homologs of rhodopsin

1007

Rearrangements in the genes for the green and red pigments lead to "color blindness"

1008

33.4 Hearing Depends on the Speedy Detection of Mechanical Stimuli

1009

Hair cells use a connected bundle of stereocilia to detect tiny motions

1009

Mechanosensory channels have been identified in *Drosophila* and vertebrates

1010

33.5 Touch Includes the Sensing of Pressure, Temperature, and Other Factors

1011

Studies of capsaicin reveal a receptor for sensing high temperatures and other painful stimuli

1011

More sensory systems remain to be studied

1012

Chapter 34 The Immune System

1017

Innate immunity is an evolutionarily ancient defense system

1018

The adaptive immune system responds by using the principles of evolution

1019

34.1 Antibodies Possess Distinct Antigen-Binding and Effector Units

1021

34.2 Antibodies Bind Specific Molecules Through Hypervariable Loops

1023

The immunoglobulin fold consists of a beta-sandwich framework with hypervariable loops

1024

X-ray analyses have revealed how antibodies bind antigens

1024

Large antigens bind antibodies with numerous interactions

1026

34.3 Diversity Is Generated by Gene Rearrangements

1027

J (joining) genes and D (diversity) genes increase antibody diversity

1027

More than 10^8 antibodies can be formed by combinatorial association and somatic mutation

1028

The oligomerization of antibodies expressed on the surfaces of immature B cells triggers antibody secretion

1029

Different classes of antibodies are formed by the hopping of V_H genes

1030

34.4 Major-Histocompatibility-Complex Proteins Present Peptide Antigens on Cell Surfaces for Recognition by T-Cell Receptors

1031

Peptides presented by MHC proteins occupy a deep groove flanked by alpha helices

1032

T-cell receptors are antibody-like proteins containing variable and constant regions

1034

CD8 on cytotoxic T cells acts in concert with T-cell receptors

1034

Helper T cells stimulate cells that display foreign peptides bound to class II MHC proteins

1036

Helper T cells rely on the T-cell receptor and CD4 to recognize foreign peptides on antigen-presenting cells

1036

MHC proteins are highly diverse

1038

Human immunodeficiency viruses subvert the immune system by destroying helper T cells

1039

34.5 The Immune System Contributes to the Prevention and the Development of Human Diseases

1040

T cells are subjected to positive and negative selection in the thymus

1040

Autoimmune diseases result from the generation of immune responses against self-antigens

1041

The immune system plays a role in cancer prevention

1041

Vaccines are a powerful means to prevent and eradicate disease

1042

Chapter 35 Molecular Motors

1049

35.1 Most Molecular-Motor Proteins Are Members of the P-Loop NTPase Superfamily

1050

Molecular motors are generally oligomeric proteins with an ATPase core and an extended structure

1050

ATP binding and hydrolysis induce changes in the conformation and binding affinity of motor proteins	1052		
35.2 Myosins Move Along Actin Filaments	1054		
Actin is a polar, self-assembling, dynamic polymer	1054		
Myosin head domains bind to actin filaments	1056		
Motions of single motor proteins can be directly observed	1056		
Phosphate release triggers the myosin power stroke	1057		
Muscle is a complex of myosin and actin	1057		
The length of the lever arm determines motor velocity	1060		
35.3 Kinesin and Dynein Move Along Microtubules	1060		
Microtubules are hollow cylindrical polymers	1060		
Kinesin motion is highly processive	1062		
35.4 A Rotary Motor Drives Bacterial Motion	1064		
Bacteria swim by rotating their flagella	1064		
Proton flow drives bacterial flagellar rotation	1064		
Bacterial chemotaxis depends on reversal of the direction of flagellar rotation	1066		
<hr/>			
Chapter 36 Drug Development	1073		
<hr/>			
36.1 The Development of Drugs Presents Huge Challenges	1074		
Drug candidates must be potent modulators of their targets	1074		
Drugs must have suitable properties to reach their targets	1075		
Toxicity can limit drug effectiveness	1080		
		36.2 Drug Candidates Can Be Discovered by Serendipity, Screening, or Design	1081
		Serendipitous observations can drive drug development	1081
		Screening libraries of compounds can yield drugs or drug leads	1083
		Drugs can be designed on the basis of three-dimensional structural information about their targets	1086
		36.3 Analyses of Genomes Hold Great Promise for Drug Discovery	1089
		Potential targets can be identified in the human proteome	1089
		Animal models can be developed to test the validity of potential drug targets	1090
		Potential targets can be identified in the genomes of pathogens	1090
		Genetic differences influence individual responses to drugs	1091
		36.4 The Development of Drugs Proceeds Through Several Stages	1092
		Clinical trials are time consuming and expensive	1092
		The evolution of drug resistance can limit the utility of drugs for infectious agents and cancer	1094
		Answers to Problems	A1
		Index	B1