

Contents

| | |
|----------------------------|-----|
| Sponsors | I |
| Preface | III |
| Group photo..... | V |
| List of participants | VII |
| Contents | XIX |

Session 1. Biotic and abiotic factors affecting induced resistance

| | |
|---|------|
| Abiotic and biotic influences on the expression of induced resistance <i>Dale R. Walters</i> | 5-10 |
|---|------|

Sub-Session 1.1. Molecular mechanisms of induced defense

| | |
|--|-------|
| Deciphering the role of <i>Arabidopsis thaliana</i> heterotrimeric G protein in induced resistance against necrotrophic fungi <i>Lucía Jordá, Viviana Escudero, Magdalena Delgado-Cerezo, Sara Sopeña, Clara Sánchez-Rodríguez, Eva Miedes, Antonio Molina</i> | 11-15 |
| Up-regulation of defense genes in pepper leaves inoculated with two different Tobamoviruses <i>Csilla Juhász, István Tóbiás, Gábor Gullner</i> | 17-20 |
| Transcriptomic and symptomatic features of PAMP triggered immunity in grapevine leaves <i>Gabriella Varga, Anita Szabó, Ernő Szegedi, Péter G. Ott, Ágnes Szatmári, Zoltán Bozsó, Arianna Lovato and Annalisa Polverari</i> | 21-26 |
| Increased expression of defense-related genes may explain enhanced <i>Botrytis cinerea</i> resistance of <i>Arabidopsis</i> plants grown in compost <i>Guillem Segarra, Gabriel Santpere, Georgina Elena, Isabel Trillas</i> | 27-30 |

Sub-Session 1.2. Priming for enhanced defense

| | |
|--|-------|
| Unraveling the epigenetic basis of transgenerational immune priming <i>Ana López, Estrella Luna, Michael R. Roberts, Pablo Vera, Jurriaan Ton</i> | 33-35 |
| <i>IBI</i> : Master regulator of β -amino butyric acid-induced priming <i>Estrella Luna, Yuhua Zhang, Marieke van Hulst, Jordi Gamir, Oliver Berkowitz, Victoria Pastor, Wouter Kegge, Dusica Acoska, Alison Van de Meere, Corné M. J. Pieterse, Victor Flors, Jurriaan Ton</i> | 37-39 |
| Chemical and constitutive priming share a common metabolomic fingerprint <i>Jordi Gamir, Miguel Cerezo, Victor Flors</i> | 41-46 |

Sub-Session 1.3. Herbivore-induced defenses

| | |
|--|-------|
| Herbivores in disguise: the wolf in sheep's clothing <i>Gary W. Felton, Seung Ho Chung, Cristina Rosa, Kelli Hoover</i> | 49-53 |
| Endogenous plant danger signaling peptides – an unwanted ingredient in a herbivore's diet? <i>Dominik Klausner, Martina Lori, Rainer Böni, Thomas Boller, Sebastian Bartels</i> | 55-59 |
| Plant mediated interactions between root parasitic nematodes and shoot herbivorous insects <i>Tom O. G. Tytgat, Mesfin Wondafrash & Nicole M. van Dam</i> | 61-62 |
| Is there a role for jasmonic acid in induced resistance against broad mites in pot azalea? <i>Gil Luypaert, Ellen De Keyser, Johan Van Huylbroeck, Johan Witters, Martine Maes, Jan De Riek, Patrick De Clercq</i> | 63-66 |
| Aboveground insect feeding leads to the recruitment of rhizobacteria for plant self-protection against subsequent diverse attacks <i>Choong-Min Ryu</i> | 67-70 |

Sub-Session 1.4. Induced defense by beneficials

| | |
|---|-------|
| Transcriptional profile of tomato roots exhibiting <i>Bacillus thuringiensis</i> - induced resistance to <i>Ralstonia solanacearum</i> <i>Shigenobu Yoshida, Hideki Takahashi, Kazuhiro Nakaho, Takeaki Ishihara, Sugihiko Ando, Yoshinori Kanayama, Seiya Tsushima, Mitsuro Hyakumachi</i> | 73-77 |
| Molecular characterization of induced resistance in grapevine: relevance of plant genotype and exposure to abiotic stresses <i>Michele Perazzolli, Maria Cristina Palmieri, Bendetta Roatti, Houda Banani, Luisa Lenzi, Ilaria Pertot</i> | 79-81 |
| Effects of temperature on the proto-cooperation between plants and <i>Bacillus amyloliquefaciens</i> S499 <i>Gerardo Puopolo, Taha Hosni, Lorenzo Pedrotti, Emmanuel Jourdan, Marc Ongena, Ilaria Pertot</i> | 83-86 |
| <i>Trichoderma harzianum</i> induces systemic resistance to <i>Alternaria alternata</i> f. sp. <i>lycopersici</i> leaf necrosis in tomato <i>Gustavo B. Meirelles, Guilherme L. Duarte, Jonathas Johnson, Lourenço M. D'Almeida, Andréa B. da Rocha, Ana P. Trivilin, Marcelo G. Moraes</i> | 87-91 |

Posters

| | |
|---|--------|
| Plant defence inducer properties of the grape marc extract <i>Razik Benouaret, Eric Goujon, Pascale Goupil</i> | 95-97 |
| Role in defense of the two glycosyltransferases <i>UGT74F1</i> and <i>UGT74F2</i> against <i>Pseudomonas syringae</i> <i>Benoît Boachon, Jordi Gamir, Victoria Pastor, Jon V. Dean, Víctor Flors, Brigitte Mauch-Mani</i> | 99-102 |

| | |
|--|---------|
| Maize antioxidant protective potential under abiotic soil drought, biotic two-spotted spider mite feeding and the combination of both stresses <i>Anna Czapla, Małgorzata Grudkowska, Anna Miazek, Małgorzata Kielkiewicz, Barbara Zagdańska</i> | 103-104 |
| Deciphering tomato defense after induction of resistance towards a biotrophic, a hemibiotrophic and a necrotrophic pathogen <i>Hejer Daoulati, Veronika Pleskova, Zuzana Storkova, Benoit Industri, Catherine Mura, Eric Galiana, Aurélie Séassau, Emeline Deleury, Philippe C. Nicot, Marc Bardin, Michel Ponchet</i> | 105-108 |
| Nitric oxide (NO) signaling in plant defense: recent progress in deciphering the mechanisms <i>Sylvain Jeandroz, Carine Fournier, Olivier Lamotte, David Wendehenne</i> | 109-112 |
| Activation of <i>Brassica napus</i> immune response by non-plant antimicrobial peptides <i>Barbora Jindřichová, Anna Macůrková, Lenka Burketová</i> | 113-117 |
| Use of <i>Bacillus amyloliquefaciens</i> to improve plant defence to insects <i>Anna Johansson, Sarosh Bejai, Johan Meijer</i> | 119-122 |
| Nonribosomal lipopeptides from <i>Bacillus subtilis</i> : potential resistance inducers and/or biopesticides for wheat crop? <i>N. G. Khong, B. Randoux, B. Tisserant, J. Deravel, Ch. Tayeh, F. Coutte, N. Bourdon, Ph. Jacques and Ph. Reignault</i> | 123-127 |
| Induced systemic resistance against <i>Pseudomonas syringae</i> pv. <i>maculicola</i> by a long chain bacterial volatile emitted from <i>Paenibacillus polymyxa</i> in <i>Arabidopsis thaliana</i> <i>Boyoung Lee, Mohamed A. Farag, Hyo Bee Park, Joseph W. Kloepper, Soo Hyun Lee, and Choong-Min Ryu</i> | 129-132 |
| Aphid infestation augments induced resistance against leaf and root pathogens in pepper <i>Boyoung Lee, Soohyun Lee, and Choong-Min Ryu</i> | 133-136 |
| Characterization of efficient resistance inducers for control of crop disease <i>Luisa Lenzi, Maria Cristina Palmieri, Carla Caruso, Ilaria Pertot, Michele Perazzolli</i> | 137-138 |
| Changes in actin dynamics are involved in the salicylic acid pathway <i>Matoušková, J., Janda, M., Šásek, V., Burketová, L., Valentová, O.</i> | 139-140 |
| Systemic resistance induced by volatile substances produced by PGPF in <i>Arabidopsis</i> plants <i>Hushna Ara Naznin, Kiyohara Daigo, Minako Kimura, Mitsuo Miyazawa, Masafumi Shimizu, Mitsuro Hyakumachi</i> | 141-144 |
| Study of the persistence of two entomopathogenic bacteria on the vegetation cover of winter soft wheat and soil <i>Hakima Oulebsir-MohandKaci, Fahima Outtar, Bahia Doumandji-Mitiche</i> | 145-147 |
| Effect of temperature on induced systemic resistance to strawberry powdery mildew <i>Dalia Rav-David, Yigal Elad</i> | 149-153 |
| Induction of resistance under different watering regimes <i>Dalia Rav-David, Yigal Elad</i> | 155-159 |

| | |
|--|---------|
| Enhanced resistance due to the simultaneous application of plant growth-promoting fungi (PGPF) and arbuscular mycorrhizal fungus (AMF) in cucumber <i>Mary Grace B. Saldajeno, Mohsen Mohamed Elsharkawy, Weddippuli Arachchige Chandanie, Masanori Ito, Masafumi Shimizu, Mitsuro Hyakumachi</i> | 161-164 |
| The behaviour of the <i>opr3</i> tomato mutant points to a role of OPDA in plant defense <i>Loredana Scalschi, Maite Sanmartin, Gemma Camañes, Pilar Troncho, Emma Fernandez-Crespo, Carmen González-Bosch, José J. Sánchez-Serrano, Begonya Vicedo and Pilar García-Agustín</i> | 165-169 |
| Does leaf position influence induced resistance to grape downy mildew? <i>Emilie Steimetz, Sophie Trouvelot, Adeline Bordier, Marie-Claire Heloir, Benoît Poinssot, Marielle Adrian, Xavier Daire</i> | 171-175 |
| Emission of sulfur-containing volatiles from <i>Arabidopsis thaliana</i> (L.) Heynh Col-0 related to diamondback moth [<i>Plutella xylostella</i> (L.)] infestation <i>Hien Truong Thi Dieu, Pierre Delaplace, Frédéric Francis, Georges Lognay</i> | 177-180 |
| Light-activated compounds may systemically reduce rice blast <i>via</i> ROS <i>Tatiana Zakharenkova, Andrey Aver'yanov, Tatiana Pasechnik, Vera Lapikova, C. Jacyn Baker</i> | 181-183 |

Session 2. Moving from model plants to crop plants

| | |
|--|---------|
| Moving from model plants to crop plants – prevailing or particular systemic defense strategies? <i>Dirk Balmer</i> | 187-191 |
|--|---------|

Sub-Session 2.1. Chemical induction of defense

| | |
|--|---------|
| Vitamin Bx-mediated induced resistance against <i>Cucumber mosaic virus</i> and <i>Xanthomonas axonopodis</i> in the pepper field <i>Geun Cheol Song, Hye Kyung Choi, Choong-Min Ryu</i> | 195-198 |
| Induced systemic resistance against systemic viruses: a feasible approach? <i>Franco Faoro</i> | 199-203 |
| A first approach to the use of hexanoic acid as a resistance inducer against <i>Xanthomonas citri</i> <i>Eugenio Llorens, Leonor Lapeña, James H. Graham, Pilar Garcia-Agustin</i> | 205-209 |
| Time course response analysis of leaf transcriptome to phosphite treatment for induced defence responses in <i>Solanum tuberosum</i> (cv. Des) <i>Dharanidhar Burra, Therese Bengtsson, Laith Moshaiib, Pete Hedley, Erland Liljeroth, Erik Andreasson, Erik Alexandersson</i> | 211-212 |

Sub-Session 2.2. Microbe- and plant-derived elicitation of defense

| | |
|---|---------|
| Bacteria induce systemic acquired resistance (SAR) in barley <i>Sanjukta Dey, Marion Wenig, Claudia Knappe, A. & Corina Vlot</i> | 215-218 |
|---|---------|

| | |
|---|---------|
| Development of a screening system for ISR-inducing <i>Trichoderma</i> spp. based on ISR-marker genes <i>Kaat De Cremer, Barbara De Coninck, Bruno P. A. Cammue &</i> <i>Christine Vos</i> | 219-222 |
| Comparison of plant defense elicitor properties of amphiphilic compounds from plant-associated bacteria <i>Martin Mariutto, Olivier Fernandez, Philippe Thonart,</i> <i>Christophe Clément, Jacques Dommes, Stephan Dorey, Marc Ongena</i> | 223-226 |
| The nature of systemic resistance induced in tomato (<i>Solanum lycopersicum</i>) by biochar soil treatments <i>Zeraye M. Haile, Yael Meller Harel, Dalia Rav-David,</i> <i>Ellen R. Graber, Yigal Elad</i> | 227-230 |

Posters

| | |
|--|---------|
| Systemic reduction of rice blast by inhibitors of antioxidant enzymes <i>Andrey Aver'yanov, Tatiana Pasechnik, Vera Lapikova,</i> <i>Tatiana Zakharenkova, C. Jacyn Baker</i> | 233-235 |
| Rootstock resistance against <i>Tetranychus urticae</i> in citrus is transmitted to grafted varieties probably activating jasmonic acid- dependent signalling <i>Blas Agut, Mónica Hurtado, Josep Jacas, Víctor Flors</i> | 237-241 |
| Plant endogenous danger peptides <i>Martina Lori, Dominik Klauser, Tim Hander, Thomas Boller,</i> <i>Sebastian Bartels</i> | 243-247 |
| Differences in 1,4-benzoxazin-3-one profile and induction during growth of young maize affect foraging patterns of leaf herbivores <i>Daniel Maag, Angela Köhler, Gaëtan Glauser, Matthias Erb,</i> <i>Jean-Luc Wolfender, Ted C. J. Turlings</i> | 249-252 |
| Induction of resistance and stress reduction by <i>Glycyrrhiza glabra</i> L. (licorice) leaf extract in cucumber and tomato <i>Marc Orlik, Andrea Scherf, Sebastian Bartels,</i> <i>Sara Mazzotta & Annegret Schmitt</i> | 253-257 |
| Identification of the VvFLS2 grapevine flagellin receptor by a functional genomics strategy <i>Lucie Trda, Freddy Boutrot, Mireille Perrin, Carine Schmitt,</i> <i>Jean Masson, Jani Kelloniemi, Marie-Claire Héloir, Xavier Daire,</i> <i>Cyril Zipfel, Benoît Poinssot</i> | 259-262 |
| Induced Systemic Resistance (ISR) signaling pathways involved in the <i>Trichoderma hamatum</i> – Tomato – <i>Botrytis cinerea</i> tripartite system <i>Yuxia Yang, Barbara De Coninck, Bruno P. A. Cammue, Christine Vos</i> | 263-266 |

Session 3. How to predict success in the field from success in the lab?

| | |
|--|---------|
| Plant defense activators: can we predict success in the field from success in the lab? <i>Marie-Noëlle Brisset</i> | 269-274 |
|--|---------|

Sub-Session 3.1. In forest trees

| | |
|--|---------|
| Moving to the field: effective protection of conifer seedlings against a forest pest by eliciting jasmonate-induced resistance <i>Rafael Zas, Niklas Björklund, Göran Nordlander, Cesar Cendán, Claes Hellqvist, Luis Sampedro</i> | 277-281 |
| Biological and chemical interventions to enhance disease resistance in radiata pine in New Zealand <i>Tony Reglinski, Joe Taylor, Mike Spiers, Annette Ah Chee, Robert Hill, Nicholas Cummings, Alison Stewart</i> | 283-287 |

Sub-Session 3.2. In agricultural crops

| | |
|--|---------|
| Are green leaf volatiles important for parasitoid wasp attraction in the field? <i>Elvira S. de Lange, Mickaël D. P. Gaillard, Yuanxin Yan, Ted C. J. Turlings, Michael V. Kolomiets, Shawn A. Christensen</i> | 291-294 |
| «BioMolChem», a tool to assess the defense status of grapevines after stimulations or not: from laboratory to the field <i>Marie-France Corio-Costet, Stéphanie Cluzet, Carole Lambert, Marie-Cécile Dufour</i> | 295-301 |
| From the lab to the field scale: dose-dependent efficacy and mode of action of resistance inducers in wheat against <i>Septoria tritici</i> leaf blotch <i>Marie-Eva Ors, Béatrice Randoux, Ali Siah, Sameh Selim, Fabienne Boizet, Céline Ade, Gilles Couleaud, Claude Maumené, Patrice Halama and Philippe Reignault</i> | 303-307 |

Session 4. Breeding for inducible resistance

| | |
|---|---------|
| Breeding for inducible resistance <i>Inger Åhman</i> | 311-317 |
| Phloem-specific resistance in <i>Brassica oleracea</i> against the whitefly <i>Aleyrodes proletella</i> <i>Colette Broekgaarden, Pierre Riviere, Greet Steenhuis, Maria del sol Cuenca, Martine Kos, Koen Pelgrom, Roeland Voorrips, Ben Vosman</i> | 319-320 |
| Increasing resistance of <i>Arabidopsis thaliana</i> towards aphids <i>Xi Chen, Zhao Zhang, Richard G. F. Visser, Colette Broekgaarden, Ben Vosman</i> | 321-322 |
| Expression of the defense gene <i>mir1</i> depends on herbivore feeding guild and maize genotype <i>Dawn S. Luthé, Joe Louis, Shan Jin, Lina Castano-Duque</i> | 323-327 |
| Effect of plant genotype on the efficacy of stimulators of plant defences in two horticultural pathosystems <i>Brigitte Maisonneuve, Caroline Allen-Aubouard, Michel Pitrat</i> | 327-331 |

| | |
|--|---------|
| Cultivar-dependent efficacy and mode of action of plant resistance inducers in wheat against <i>Septoria tritici</i> leaf blotch <i>Marie-Eva Ors, Ali Siah, Béatrice Randoux, Sameh Selim, Gilles Couleaud, Claude Maumené, Philippe Reignault and Patrice Halama</i> | 333-338 |
| Effects of the endophyte <i>Burkholderia phytofirmans</i> PsJN and its lipopolysaccharide on potato leaves <i>Friederike Trognitz, Birgit Mitter, Antonio Molinaro, Valerie Hubalek, Angela Sessitsch</i> | 339-343 |

Posters

| | |
|---|---------|
| Genetic variation in inducibility in two Mediterranean pines <i>Luis Sampedro, Xoaquín Moreira, Rafael Zas</i> | 347-350 |
| A strategy towards bioprotection of tropical crops: Experiences and perspectives with ISR on pineapple and banana in Martinique <i>Alain Soler, Paul-Alex Marie-Alphonsine, Claudine Corbion, Paula Fernandes, Nayanci Portal Gonzalez, Rayza Gonzalez, Anne Repellin, Stephan Declerck, Patrick Quénehervé</i> | 351-356 |

Session 5. Induced resistance and field adventures

| | |
|---|---------|
| Induced resistance and field adventure: from lab success to field reality <i>Sonia Hallier</i> | 359-363 |
|---|---------|

Sub-Session 5.1. Field control of insect pests

| | |
|---|---------|
| Seed treatment with trinexapac-ethyl induces resistance to white fly (<i>Bemisia tabaci</i>) in lettuce (<i>Lactuca sativa</i> L.) seedlings <i>Joshua D. Klein, Hila Duvdevani, Joshua Herskovitz, Peninah Feldman, Yonit Hebbe, Phyllis G. Weintraub</i> | 367-371 |
| Bio-inoculants to boost resistance in crop plants against insects: Field success and struggles in India <i>V. Selvanarayanan</i> | 373-378 |
| Induction of plant systemic defense by a gaseous compound against <i>Pseudomonas syringae</i> and aphid in cucumber in the open field <i>Geun Cheol Song and Choong-Min Ryu</i> | 379-383 |

Sub-Session 5.2. Field control of diseases

| | |
|--|---------|
| Plant defense priming in lab and field <i>Uwe Conrath</i> | 387-390 |
| BioMolChem tools used in the vineyard to predict grapevine protection against downy mildew after elicitation <i>Marie-Cécile Dufour, Lionel Druelle, Marie-France Corio-Costet</i> | 391-396 |

| | |
|--|---------|
| PRB1 induces resistance in garlic, onion and pepper, increasing yield and crop quality <i>Marisé Borja, Julio Bonet, Yolanda Sanz, Antonio Molina</i> | 397-402 |
| COS-OGA, a new oligosaccharidic elicitor that induces protection against a wide range of plant pathogens <i>Géraldine van Aubel, Raffael Buonatesta, Pierre Van Cutsem</i> | 403-407 |
| Use of elicitors to protect melon crops (<i>Cucumis melo</i>) against <i>Podosphaera xanthii</i> : A multisite field experiment <i>Marie Torres</i> | 409-414 |

Posters

| | |
|---|---------|
| Reducing the amounts of copper in vineyards against <i>Plasmopara viticola</i> by the use of a low dose of D-fructose <i>Ingrid Arnault, Arnaud Furet, Marc Chovelon, Christelle Gomez, Sylvie Derridj</i> | 417-420 |
| Influence of inducible benzoxazinoid derivatives in maize on larval dispersal of the European corn borer (<i>Ostrinia nubilalis</i>) <i>Noémie Chervet, Françoise Klötzli and Thomas Steinger</i> | 421-423 |
| Application of bacterial volatiles in the pepper roots primes systemic resistance under field condition <i>Hye Kyung Choi, Geun Cheol Song, Joon-Hui Chung, and Choong-Min Ryu</i> | 425-428 |
| BABA (β -aminobutyric acid) induced resistance against grey mould and virus infection in grapevine <i>Anna Csikász-Krizsics, Anikó Mátai, Ágnes Nagy, Sándor Kovács, Brigitta Végh, János Werner, Gábor Jakab</i> | 429-432 |
| Can plant defence elicitors control bacterial infection in field vegetables? <i>Nicola J. Holden, Ian K. Toth, Adrian Newton and Dale Walters</i> | 433-436 |
| The use of effective microorganisms as plant growth promoters and biotic resistance inducers <i>Jolanta Kowalska, Dorota Remlein-Starosta, Iwona Morkunas</i> | 437-442 |
| Bio-stimulant effect of two Algerian isolates of <i>Trichoderma</i> spp. on tomato (<i>Lycopersicon esculentum</i> Mill.) and their elicitor effect on <i>Tuta absoluta</i> (Povolny, 1994) ex (Meyrick, 1917) <i>Saida Moumene-Messgo, Rym Boukhalfa, Zahia Houmani, Dounia Saddek, Khadidja Bencheikh, Sabrina Zanoune, Mohamed Bellatreche</i> | 443-448 |
| Apple scab control with plant resistance improving product in organic apple growing <i>Václav Psota, Kateřina Rychlá, Michal Katriňák</i> | 449-453 |
| Potential for integrated control of the wheat pathogen, <i>Stagonospora nodorum</i> , by Folicur and extracellular compounds produced by isolate FS-94 of <i>Fusarium sambucinum</i> <i>Larisa Shcherbakova, Yuliay Semina, Tatyana Nazarova, Lubov Dorofeeva, Vitaly Dzhavakhiya, Bruce Campbell</i> | 455-458 |
| Bio-stimulants as inducers of defense reactions against <i>Tetranychus urticae</i> Koch in greenhouse cucumber <i>A. Tomczyk</i> | 459-462 |