

# Brief Contents

CHAPTER 1	<i>Plant and Cell Architecture</i>	1
CHAPTER 2	<i>Genome Structure and Gene Expression</i>	51
<b>UNIT I</b>	<b>Transport and Translocation of Water and Solutes</b>	81
CHAPTER 3	<i>Water and Plant Cells</i>	83
CHAPTER 4	<i>Water Balance of Plants</i>	99
CHAPTER 5	<i>Mineral Nutrition</i>	119
CHAPTER 6	<i>Solute Transport</i>	143
<b>UNIT II</b>	<b>Biochemistry and Metabolism</b>	169
CHAPTER 7	<i>Photosynthesis: The Light Reactions</i>	171
CHAPTER 8	<i>Photosynthesis: The Carbon Reactions</i>	203
CHAPTER 9	<i>Photosynthesis: Physiological and Ecological Considerations</i>	245
CHAPTER 10	<i>Stomatal Biology</i>	269
CHAPTER 11	<i>Translocation in the Phloem</i>	285
CHAPTER 12	<i>Respiration and Lipid Metabolism</i>	317
CHAPTER 13	<i>Assimilation of Inorganic Nutrients</i>	353
<b>UNIT III</b>	<b>Growth and Development</b>	377
CHAPTER 14	<i>Cell Walls: Structure, Formation, and Expansion</i>	379
CHAPTER 15	<i>Signals and Signal Transduction</i>	407
CHAPTER 16	<i>Signals from Sunlight</i>	447
CHAPTER 17	<i>Embryogenesis</i>	477
CHAPTER 18	<i>Seed Dormancy, Germination, and Seedling Establishment</i>	513
CHAPTER 19	<i>Vegetative Growth and Organogenesis</i>	553
CHAPTER 20	<i>The Control of Flowering and Floral Development</i>	591
CHAPTER 21	<i>Gametophytes, Pollination, Seeds, and Fruits</i>	625
CHAPTER 22	<i>Plant Senescence and Cell Death</i>	665
CHAPTER 23	<i>Biotic Interactions</i>	693
CHAPTER 24	<i>Abiotic Stress</i>	731

# Table of Contents

## CHAPTER 1

### Plant and Cell Architecture 1

#### Plant Life Processes: Unifying Principles 2

#### Plant Classification and Life Cycles 2

Plant life cycles alternate between diploid and haploid generations 3

#### Overview of Plant Structure 5

Plant cells are surrounded by rigid cell walls 5

Plasmodesmata allow the free movement of molecules between cells 8

New cells originate in dividing tissues called meristems 8

#### Plant Cell Organelles 10

Biological membranes are phospholipid bilayers that contain proteins 10

#### The Endomembrane System 13

The nucleus contains the majority of the genetic material 13

Gene expression involves both transcription and translation 17

The endoplasmic reticulum is a network of internal membranes 17

Secretion of proteins from cells begins with the rough ER 19

Glycoproteins and polysaccharides destined for secretion are processed in the Golgi apparatus 20

The plasma membrane has specialized regions involved in membrane recycling 22

Vacuoles have diverse functions in plant cells 23

#### Independently Dividing or Fusing Organelles

#### Derived from the Endomembrane System 23

Oil bodies are lipid-storing organelles 23

Microbodies play specialized metabolic roles in leaves and seeds 24

#### Independently Dividing, Semiautonomous Organelles 25

Proplastids mature into specialized plastids in different plant tissues 27

Chloroplast and mitochondrial division are independent of nuclear division 29

#### The Plant Cytoskeleton 29

The plant cytoskeleton consists of microtubules and microfilaments 29

Actin, tubulin, and their polymers are in constant flux in the living cell 31

Cortical microtubules move around the cell by treadmilling 33

Cytoskeletal motor proteins mediate cytoplasmic streaming and directed organelle movement 33

#### Cell Cycle Regulation 35

Each phase of the cell cycle has a specific set of biochemical and cellular activities 35

The cell cycle is regulated by cyclins and cyclin-dependent kinases 36

Mitosis and cytokinesis involve both microtubules and the endomembrane system 37

#### Plant Cell Types 39

Dermal tissues cover the surfaces of plants 39

Ground tissues form the bodies of plants 40

Vascular tissues form transport networks between different parts of the plant 44

## CHAPTER 2

### Genome Structure and Gene Expression 51

#### Nuclear Genome Organization 51

The nuclear genome is packaged into chromatin 52

Centromeres, telomeres, and nucleolar organizer regions contain repetitive sequences 52

Transposons are mobile sequences within the genome 53

Chromosome organization is not random in the interphase nucleus 54

Meiosis halves the number of chromosomes and allows for the recombination of alleles	54	Epigenetic modifications help determine gene activity	65
Polyploids contain multiple copies of the entire genome	56	<b>Posttranscriptional Regulation of Nuclear Gene Expression</b>	67
Phenotypic and physiological responses to polyploidy are unpredictable	58	All RNA molecules are subject to decay	67
The role of polyploidy in evolution is still unclear	60	Noncoding RNAs regulate mRNA activity via the RNA interference (RNAi) pathway	67
<b>Plant Cytoplasmic Genomes: Mitochondria and Plastids</b>	61	Posttranslational regulation determines the life span of proteins	71
The endosymbiotic theory describes the origin of cytoplasmic genomes	61	<b>Tools for Studying Gene Function</b>	72
Organellar genomes vary in size	61	Mutant analysis can help elucidate gene function	72
Organellar genetics do not obey Mendelian principles	61	Molecular techniques can measure the activity of genes	73
<b>Transcriptional Regulation of Nuclear Gene Expression</b>	62	Gene fusions can introduce reporter genes	74
RNA polymerase II binds to the promoter region of most protein-coding genes	62	<b>Genetic Modification of Crop Plants</b>	76
Conserved nucleotide sequences signal transcriptional termination and polyadenylation	64	Transgenes can confer resistance to herbicides or plant pests	77
		Genetically modified organisms are controversial	77

## UNIT I Transport and Translocation of Water and Solutes 81

### CHAPTER 3 Water and Plant Cells 83

#### Water in Plant Life 83

##### The Structure and Properties of Water 84

Water is a polar molecule that forms hydrogen bonds	84	Water enters the cell along a water potential gradient	91
Water is an excellent solvent	85	Water can also leave the cell in response to a water potential gradient	92
Water has distinctive thermal properties relative to its size	85	Water potential and its components vary with growth conditions and location within the plant	93
Water molecules are highly cohesive	85	<b>Cell Wall and Membrane Properties</b>	93
Water has a high tensile strength	86	Small changes in plant cell volume cause large changes in turgor pressure	93

##### Diffusion and Osmosis 87

Diffusion is the net movement of molecules by random thermal agitation	87	The rate at which cells gain or lose water is influenced by cell membrane hydraulic conductivity	94
Diffusion is most effective over short distances	88	Aquaporins facilitate the movement of water across cell membranes	95
Osmosis describes the net movement of water across a selectively permeable barrier	88	<b>Plant Water Status</b>	96

##### Water Potential 89

The chemical potential of water represents the free-energy status of water	89	Physiological processes are affected by plant water status	96
Three major factors contribute to cell water potential	90	Solute accumulation helps cells maintain turgor and volume	96
Water potentials can be measured	90		

##### Water Potential of Plant Cells 91

Water enters the cell along a water potential gradient	91
Water can also leave the cell in response to a water potential gradient	92

Water potential and its components vary with growth conditions and location within the plant	93
--	----

##### **Cell Wall and Membrane Properties**

Small changes in plant cell volume cause large changes in turgor pressure	93
The rate at which cells gain or lose water is influenced by cell membrane hydraulic conductivity	94
Aquaporins facilitate the movement of water across cell membranes	95

##### **Plant Water Status**

Physiological processes are affected by plant water status	96
Solute accumulation helps cells maintain turgor and volume	96

### CHAPTER 4

### Water Balance of Plants 99

#### Water in the Soil 99

A negative hydrostatic pressure in soil water lowers soil water potential	100
---	-----

Water moves through the soil by bulk flow 101	Crop yields can be improved by the addition of fertilizers 130
<b>Water Absorption by Roots 101</b>	Some mineral nutrients can be absorbed by leaves 131
Water moves in the root via the apoplast, symplast, and transmembrane pathways 102	<b>Soil, Roots, and Microbes 131</b>
Solute accumulation in the xylem can generate “root pressure” 103	Negatively charged soil particles affect the adsorption of mineral nutrients 131
<b>Water Transport through the Xylem 104</b>	Soil pH affects nutrient availability, soil microbes, and root growth 132
The xylem consists of two types of transport cells 104	Excess mineral ions in the soil limit plant growth 133
Water moves through the xylem by pressure-driven bulk flow 105	Some plants develop extensive root systems 133
Water movement through the xylem requires a smaller pressure gradient than movement through living cells 106	Root systems differ in form but are based on common structures 134
What pressure difference is needed to lift water 100 meters to a treetop? 107	Different areas of the root absorb different mineral ions 135
The cohesion–tension theory explains water transport in the xylem 107	Nutrient availability influences root growth 137
Xylem transport of water in trees faces physical challenges 108	Mycorrhizal symbioses facilitate nutrient uptake by roots 137
Plants minimize the consequences of xylem cavitation 110	Nutrients move between mycorrhizal fungi and root cells 140
<b>Water Movement from the Leaf to the Atmosphere 110</b>	<b>CHAPTER 6 Solute Transport 143</b>
Leaves have a large hydraulic resistance 111	<b>Passive and Active Transport 144</b>
The driving force for transpiration is the difference in water vapor concentration 111	<b>Transport of Ions across Membrane Barriers 145</b>
Water loss is also regulated by the pathway resistances 112	Different diffusion rates for cations and anions produce diffusion potentials 146
Stomatal control couples leaf transpiration to leaf photosynthesis 112	How does membrane potential relate to ion distribution? 146
The cell walls of guard cells have specialized features 113	The Nernst equation distinguishes between active and passive transport 147
An increase in guard cell turgor pressure opens the stomata 115	Proton transport is a major determinant of the membrane potential 148
The transpiration ratio measures the relationship between water loss and carbon gain 116	<b>Membrane Transport Processes 149</b>
<b>Overview: The Soil–Plant–Atmosphere Continuum 116</b>	Channels enhance diffusion across membranes 150
<b>CHAPTER 5 Mineral Nutrition 119</b>	Carriers bind and transport specific substances 151
<b>Essential Nutrients, Deficiencies, and Plant Disorders 120</b>	Primary active transport requires energy 151
Special techniques are used in nutritional studies 122	Kinetic analyses can elucidate transport mechanisms 154
Nutrient solutions can sustain rapid plant growth 122	<b>Membrane Transport Proteins 155</b>
Mineral deficiencies disrupt plant metabolism and function 125	The genes for many transporters have been identified 157
Analysis of plant tissues reveals mineral deficiencies 129	Transporters exist for diverse nitrogen-containing compounds 157
<b>Treating Nutritional Deficiencies 129</b>	Cation transporters are diverse 158
	Anion transporters have been identified 160
	Transporters for metal and metalloid ions transport essential micronutrients 160
	Aquaporins have diverse functions 160
	Plasma membrane H <sup>+</sup> -ATPases are highly regulated P-type ATPases 161

The tonoplast H<sup>+</sup>-ATPase drives solute accumulation in vacuoles 162  
H<sup>+</sup>-pyrophosphatases also pump protons at the tonoplast 163

### **Ion Transport in Roots 163**

Solutes move through both apoplast and symplast 164  
Ions cross both symplast and apoplast 164  
Xylem parenchyma cells participate in xylem loading 164

## **UNIT II**

# **Biochemistry and Metabolism 169**

## **CHAPTER 7**

### **Photosynthesis: The Light Reactions 171**

#### **Photosynthesis in Higher Plants 171**

##### **General Concepts 172**

Light has characteristics of both a particle and a wave 172  
When molecules absorb or emit light, they change their electronic state 173  
Photosynthetic pigments absorb the light that powers photosynthesis 175

##### **Key Experiments in Understanding Photosynthesis 175**

Action spectra relate light absorption to photosynthetic activity 176  
Photosynthesis takes place in complexes containing light-harvesting antennas and photochemical reaction centers 176  
The chemical reaction of photosynthesis is driven by light 178  
Light drives the reduction of NADP<sup>+</sup> and the formation of ATP 178  
Oxygen-evolving organisms have two photosystems that operate in series 179

##### **Organization of the Photosynthetic Apparatus 180**

The chloroplast is the site of photosynthesis 180  
Thylakoids contain integral membrane proteins 181  
Photosystems I and II are spatially separated in the thylakoid membrane 181  
Anoxygenic photosynthetic bacteria have a single reaction center 182

##### **Organization of Light-Absorbing Antenna Systems 183**

Antenna systems contain chlorophyll and are membrane-associated 183  
The antenna funnels energy to the reaction center 183  
Many antenna pigment–protein complexes have a common structural motif 183

### **Mechanisms of Electron Transport 185**

Electrons from chlorophyll travel through the carriers organized in the Z scheme 185  
Energy is captured when an excited chlorophyll reduces an electron acceptor molecule 186  
The reaction center chlorophylls of the two photosystems absorb at different wavelengths 187  
The PSII reaction center is a multi-subunit pigment–protein complex 188  
Water is oxidized to oxygen by PSII 188  
Pheophytin and two quinones accept electrons from PSII 189  
Electron flow through the cytochrome *b*<sub>6</sub>*f* complex also transports protons 191  
Plastoquinone and plastocyanin carry electrons between photosystems II and I 192  
The PSI reaction center reduces NADP<sup>+</sup> 192  
Cyclic electron flow generates ATP but no NADPH 193  
Some herbicides block photosynthetic electron flow 193

### **Proton Transport and ATP Synthesis in the Chloroplast 193**

### **Repair and Regulation of the Photosynthetic Machinery 195**

Carotenoids serve as photoprotective agents 196  
Some xanthophylls also participate in energy dissipation 197  
The PSII reaction center is easily damaged 197  
PSI is protected from active oxygen species 198  
Thylakoid stacking permits energy partitioning between the photosystems 198

### **Genetics, Assembly, and Evolution of Photosynthetic Systems 198**

Chloroplast genes exhibit non-Mendelian patterns of inheritance 198  
Most chloroplast proteins are imported from the cytoplasm 199  
The biosynthesis and breakdown of chlorophyll are complex pathways 199  
Complex photosynthetic organisms have evolved from simpler forms 199

# CHAPTER 8

## Photosynthesis: The Carbon Reactions 203

### The Calvin–Benson Cycle 204

- The Calvin–Benson cycle has three phases: carboxylation, reduction, and regeneration 204
- The fixation of CO<sub>2</sub> via carboxylation of ribulose 1,5-bisphosphate and the reduction of the product 3-phosphoglycerate yield triose phosphates 206
- The regeneration of ribulose 1,5-bisphosphate ensures the continuous assimilation of CO<sub>2</sub> 207
- An induction period precedes the steady state of photosynthetic CO<sub>2</sub> assimilation 208
- Many mechanisms regulate the Calvin–Benson cycle 209
- Rubisco-activase regulates the catalytic activity of rubisco 209
- Light regulates the Calvin–Benson cycle via the ferredoxin–thioredoxin system 210
- Light-dependent ion movements modulate enzymes of the Calvin–Benson cycle 211
- Light controls the assembly of chloroplast enzymes into supramolecular complexes 211

### The C<sub>2</sub> Oxidative Photosynthetic Carbon Cycle 211

- The oxygenation of ribulose 1,5-bisphosphate sets in motion the C<sub>2</sub> oxidative photosynthetic carbon cycle 213
- Photorespiration is linked to the photosynthetic electron transport system 217
- Enzymes of the plant C<sub>2</sub> oxidative photosynthetic carbon cycle derive from different ancestors 217
- Cyanobacteria use a proteobacterial pathway for bringing carbon atoms of 2-phosphoglycolate back to the Calvin–Benson cycle 217
- The C<sub>2</sub> oxidative photosynthetic carbon cycle interacts with many metabolic pathways 218
- Production of biomass may be enhanced by engineering photorespiration 219

### Inorganic Carbon–Concentrating Mechanisms 220

#### Inorganic Carbon–Concentrating Mechanisms: The C<sub>4</sub> Carbon Cycle 220

- Malate and aspartate are the primary carboxylation products of the C<sub>4</sub> cycle 221
- The C<sub>4</sub> cycle assimilates CO<sub>2</sub> by the concerted action of two different types of cells 222
- The C<sub>4</sub> cycle uses different mechanisms for decarboxylation of four-carbon acids transported to bundle sheath cells 224

- Bundle sheath cells and mesophyll cells exhibit anatomical and biochemical differences 224
- The C<sub>4</sub> cycle also concentrates CO<sub>2</sub> in single cells 225
- Light regulates the activity of key C<sub>4</sub> enzymes 225
- Photosynthetic assimilation of CO<sub>2</sub> in C<sub>4</sub> plants demands more transport processes than in C<sub>3</sub> plants 225
- In hot, dry climates, the C<sub>4</sub> cycle reduces photorespiration 228

### Inorganic Carbon–Concentrating Mechanisms: Crassulacean Acid Metabolism (CAM) 228

- Different mechanisms regulate C<sub>4</sub> PEPCase and CAM PEPCase 230
- CAM is a versatile mechanism sensitive to environmental stimuli 230

### Accumulation and Partitioning of Photosynthates—Starch and Sucrose 230

#### Formation and Mobilization of Chloroplast Starch 231

- Chloroplast stroma accumulates starch as insoluble granules during the day 233
- Starch degradation at night requires the phosphorylation of amylopectin 236
- The export of maltose prevails in the nocturnal breakdown of transitory starch 237
- The synthesis and degradation of the starch granule are regulated by multiple mechanisms 237

#### Sucrose Biosynthesis and Signaling 238

- Triose phosphates from the Calvin–Benson cycle build up the cytosolic pool of three important hexose phosphates in the light 238
- Fructose 2,6-bisphosphate regulates the hexose phosphate pool in the light 239
- Sucrose is continuously synthesized in the cytosol 239

# CHAPTER 9

## Photosynthesis: Physiological and Ecological Considerations 245

### Photosynthesis Is Influenced by Leaf Properties 246

- Leaf anatomy and canopy structure maximize light absorption 247
- Leaf angle and leaf movement can control light absorption 249
- Leaves acclimate to sun and shade environments 249

### Effects of Light on Photosynthesis in the Intact Leaf 250

- Light-response curves reveal photosynthetic properties 250

Leaves must dissipate excess light energy 252	<b>CHAPTER 11</b>
Absorption of too much light can lead to photoinhibition 254	
<b>Effects of Temperature on Photosynthesis in the Intact Leaf 255</b>	<b>Translocation in the Phloem 285</b>
Leaves must dissipate vast quantities of heat 255	<b>Pathways of Translocation 286</b>
There is an optimal temperature for photosynthesis 256	Sugar is translocated in phloem sieve elements 286
Photosynthesis is sensitive to both high and low temperatures 256	Mature sieve elements are living cells specialized for translocation 287
Photosynthetic efficiency is temperature-sensitive 257	Large pores in cell walls are the prominent feature of sieve elements 288
<b>Effects of Carbon Dioxide on Photosynthesis in the Intact Leaf 258</b>	Damaged sieve elements are sealed off 289
Atmospheric CO <sub>2</sub> concentration keeps rising 258	Companion cells aid the highly specialized sieve elements 290
CO <sub>2</sub> diffusion to the chloroplast is essential to photosynthesis 258	<b>Patterns of Translocation: Source to Sink 291</b>
CO <sub>2</sub> imposes limitations on photosynthesis 260	<b>Materials Translocated in the Phloem 292</b>
How will photosynthesis and respiration change in the future under elevated CO <sub>2</sub> conditions? 262	Phloem sap can be collected and analyzed 292
<b>Stable Isotopes Record Photosynthetic Properties 264</b>	Sugars are translocated in a nonreducing form 293
How do we measure the stable carbon isotopes of plants? 264	Other solutes are translocated in the phloem 293
Why are there carbon isotope ratio variations in plants? 265	<b>Rates of Movement 295</b>
<b>CHAPTER 10</b>	<b>The Pressure-Flow Model, a Passive Mechanism for Phloem Transport 295</b>
<b>Stomatal Biology 269</b>	An osmotically generated pressure gradient drives translocation in the pressure-flow model 295
<b>Light-dependent Stomatal Opening 270</b>	Some predictions of pressure flow have been confirmed, while others require further experimentation 296
Guard cells respond to blue light 270	There is no bidirectional transport in single sieve elements, and solutes and water move at the same velocity 297
Blue light activates a proton pump at the guard cell plasma membrane 271	The energy requirement for transport through the phloem pathway is small in herbaceous plants 297
Blue-light responses have characteristic kinetics and lag times 273	Sieve plate pores appear to be open channels 298
Blue light regulates the osmotic balance of guard cells 273	Pressure gradients in the sieve elements may be modest; pressures in herbaceous plants and trees appear to be similar 298
Sucrose is an osmotically active solute in guard cells 275	Alternative models for translocation by mass flow have been suggested 299
<b>Mediation of Blue-light Photoreception in Guard Cells by Zeaxanthin 276</b>	Does translocation in gymnosperms involve a different mechanism? 299
<b>Reversal of Blue Light-Stimulated Opening by Green Light 278</b>	<b>Phloem Loading 300</b>
A carotenoid–protein complex senses light intensity 280	Phloem loading can occur via the apoplast or symplast 300
<b>The Resolving Power of Photophysiology 280</b>	Abundant data support the existence of apoplastic loading in some species 301
	Sucrose uptake in the apoplastic pathway requires metabolic energy 301
	Phloem loading in the apoplastic pathway involves a sucrose–H <sup>+</sup> symporter 302
	Phloem loading is symplastic in some species 302
	The polymer-trapping model explains symplastic loading in plants with intermediary-type companion cells 303

Phloem loading is passive in several tree species 304  
 The type of phloem loading is correlated with several significant characteristics 304

### **Phloem Unloading and Sink-to-Source Transition 305**

Phloem unloading and short-distance transport can occur via symplastic or apoplastic pathways 305  
 Transport into sink tissues requires metabolic energy 306  
 The transition of a leaf from sink to source is gradual 307

### **Photosynthate Distribution: Allocation and Partitioning 309**

Allocation includes storage, utilization, and transport 309  
 Various sinks partition transport sugars 309  
 Source leaves regulate allocation 310  
 Sink tissues compete for available translocated photosynthate 310  
 Sink strength depends on sink size and activity 311  
 The source adjusts over the long term to changes in the source-to-sink ratio 311

### **Transport of Signaling Molecules 311**

Turgor pressure and chemical signals coordinate source and sink activities 312  
 Proteins and RNAs function as signal molecules in the phloem to regulate growth and development 312  
 Plasmodesmata function in phloem signaling 313

## **CHAPTER 12**

### **Respiration and Lipid Metabolism 317**

#### **Overview of Plant Respiration 317**

##### **Glycolysis 321**

Glycolysis metabolizes carbohydrates from several sources 321  
 The energy-conserving phase of glycolysis extracts usable energy 322  
 Plants have alternative glycolytic reactions 322  
 In the absence of oxygen, fermentation regenerates the NAD<sup>+</sup> needed for glycolysis 323  
 Plant glycolysis is controlled by its products 324

##### **The Oxidative Pentose Phosphate Pathway 324**

The oxidative pentose phosphate pathway produces NADPH and biosynthetic intermediates 326  
 The oxidative pentose phosphate pathway is redox-regulated 326

##### **The Citric Acid Cycle 326**

Mitochondria are semiautonomous organelles 327

Pyruvate enters the mitochondrion and is oxidized via the citric acid cycle 328

The citric acid cycle of plants has unique features 329

### **Mitochondrial Electron Transport and ATP Synthesis 329**

The electron transport chain catalyzes a flow of electrons from NADH to O<sub>2</sub> 330  
 The electron transport chain has supplementary branches 332  
 ATP synthesis in the mitochondrion is coupled to electron transport 333  
 Transporters exchange substrates and products 334  
 Aerobic respiration yields about 60 molecules of ATP per molecule of sucrose 334  
 Several subunits of respiratory complexes are encoded by the mitochondrial genome 336  
 Plants have several mechanisms that lower the ATP yield 336  
 Short-term control of mitochondrial respiration occurs at different levels 338  
 Respiration is tightly coupled to other pathways 339

### **Respiration in Intact Plants and Tissues 340**

Plants respire roughly half of the daily photosynthetic yield 340  
 Respiration operates during photosynthesis 341  
 Different tissues and organs respire at different rates 341  
 Environmental factors alter respiration rates 342

### **Lipid Metabolism 343**

Fats and oils store large amounts of energy 343  
 Triacylglycerols are stored in oil bodies 343  
 Polar glycerolipids are the main structural lipids in membranes 344  
 Fatty acid biosynthesis consists of cycles of two-carbon addition 344  
 Glycerolipids are synthesized in the plastids and the ER 346  
 Lipid composition influences membrane function 348  
 Membrane lipids are precursors of important signaling compounds 348  
 Storage lipids are converted into carbohydrates in germinating seeds 348

## **CHAPTER 13**

### **Assimilation of Inorganic Nutrients 353**

#### **Nitrogen in the Environment 354**

Nitrogen passes through several forms in a biogeochemical cycle 354  
 Unassimilated ammonium or nitrate may be dangerous 355

**Nitrate Assimilation 356**

Many factors regulate nitrate reductase 356  
Nitrite reductase converts nitrite to ammonium 357  
Both roots and shoots assimilate nitrate 357

**Ammonium Assimilation 358**

Converting ammonium to amino acids requires two enzymes 358  
Ammonium can be assimilated via an alternative pathway 360  
Transamination reactions transfer nitrogen 360  
Asparagine and glutamine link carbon and nitrogen metabolism 360

**Amino Acid Biosynthesis 360****Biological Nitrogen Fixation 360**

Free-living and symbiotic bacteria fix nitrogen 361  
Nitrogen fixation requires microanaerobic or anaerobic conditions 362  
Symbiotic nitrogen fixation occurs in specialized structures 363  
Establishing symbiosis requires an exchange of signals 364

Nod factors produced by bacteria act as signals for symbiosis 364

Nodule formation involves phytohormones 365  
The nitrogenase enzyme complex fixes N<sub>2</sub> 366  
Amides and ureides are the transported forms of nitrogen 367

**Sulfur Assimilation 367**

Sulfate is the form of sulfur transported into plants 368  
Sulfate assimilation requires the reduction of sulfate to cysteine 368  
Sulfate assimilation occurs mostly in leaves 369  
Methionine is synthesized from cysteine 369

**Phosphate Assimilation 369****Cation Assimilation 370**

Cations form noncovalent bonds with carbon compounds 370  
Roots modify the rhizosphere to acquire iron 371  
Iron cations form complexes with carbon and phosphate 372

**Oxygen Assimilation 372****The Energetics of Nutrient Assimilation 372****UNIT  
III****Growth and Development 377****CHAPTER 14****Cell Walls: Structure, Formation, and Expansion 379****Overview of Plant Cell Wall Functions and Structures 380**

Plants vary in structure and function 380  
Components differ for primary and secondary cell walls 382  
Cellulose microfibrils have an ordered structure and are synthesized at the plasma membrane 384  
Matrix polymers are synthesized in the Golgi apparatus and secreted via vesicles 387  
Pectins are hydrophilic gel-forming components of the primary cell wall 388  
Hemicelluloses are matrix polysaccharides that bind to cellulose 390

**Primary Cell Wall Structure and Function 392**

The primary cell wall is composed of cellulose microfibrils embedded in a matrix of pectins and hemicelluloses 392

New primary cell walls are assembled during cytokinesis and continue to be assembled during growth 392

**Mechanisms of Cell Expansion 393**

Microfibril orientation influences growth directionality of cells with diffuse growth 394  
Cortical microtubules influence the orientation of newly deposited microfibrils 395

**The Extent and Rate of Cell Growth 397**

Stress relaxation of the cell wall drives water uptake and cell expansion 397

Acid-induced growth and wall stress relaxation are mediated by expansins 397

Cell wall models are hypotheses about how molecular components fit together to make a functional wall 399

Many structural changes accompany the cessation of wall expansion 400

**Secondary Cell Wall Structure and Function 400**

Secondary cell walls are rich in cellulose and hemi-cellulose and often have a hierarchical organization 400

Lignification transforms the SCW into a hydrophobic structure resistant to deconstruction 402

## CHAPTER 15

### Signals and Signal Transduction 407

#### Temporal and Spatial Aspects of Signaling 408

##### Signal Perception and Amplification 409

Receptors are located throughout the cell and are conserved across kingdoms 409

Signals must be amplified intracellularly to regulate their target molecules 411

The MAP kinase signal amplification cascade is present in all eukaryotes 411

$\text{Ca}^{2+}$  is the most ubiquitous second messenger in plants and other eukaryotes 411

Changes in the cytosolic or cell wall pH can serve as second messengers for hormonal and stress responses 412

Reactive oxygen species act as second messengers mediating both environmental and developmental signals 413

Lipid signaling molecules act as second messengers that regulate a variety of cellular processes 414

#### Hormones and Plant Development 414

Auxin was discovered in early studies of coleoptile bending during phototropism 417

Gibberellins promote stem growth and were discovered in relation to the “foolish seedling disease” of rice 417

Cytokinins were discovered as cell division-promoting factors in tissue culture experiments 418

Ethylene is a gaseous hormone that promotes fruit ripening and other developmental processes 419

Abscisic acid regulates seed maturation and stomatal closure in response to water stress 419

Brassinosteroids regulate photomorphogenesis, germination, and other developmental processes 420

Strigolactones suppress branching and promote rhizosphere interactions 421

#### Phytohormone Metabolism and Homeostasis 421

Indole-3-pyruvate is the primary intermediate in auxin biosynthesis 421

Gibberellins are synthesized by oxidation of the diterpene *ent*-kaurene 422

Cytokinins are adenine derivatives with isoprene side chains 423

Ethylene is synthesized from methionine via the intermediate ACC 426

Abscisic acid is synthesized from a carotenoid intermediate 426

Brassinosteroids are derived from the sterol campesterol 428

Strigolactones are synthesized from  $\beta$ -carotene 429

#### Signal Transmission and Cell–Cell Communication 429

#### Hormonal Signaling Pathways 431

The cytokinin and ethylene signal transduction pathways are derived from the bacterial two-component regulatory system 431

Receptor-like kinases mediate brassinosteroid and certain auxin signaling pathways 433

The core ABA signaling components include phosphatases and kinases 436

Plant hormone signaling pathways generally employ negative regulation 436

Several plant hormone receptors encode components of the ubiquitination machinery and mediate signaling via protein degradation 437

Plants have evolved mechanisms for switching off or attenuating signaling responses 439

The cellular response output to a signal is often tissue-specific 441

Cross-regulation allows signal transduction pathways to be integrated 441

## CHAPTER 16

### Signals from Sunlight 447

#### Plant Photoreceptors 448

Photoreponses are driven by light quality or spectral properties of the energy absorbed 449

Plants responses to light can be distinguished by the amount of light required 450

#### Phytochromes 452

Phytochrome is the primary photoreceptor for red and far-red light 452

Phytochrome can interconvert between Pr and Pfr forms 452

Pfr is the physiologically active form of phytochrome 453

The phytochrome chromophore and protein both undergo conformational changes in response to red light 453

Pfr is partitioned between the cytosol and the nucleus 454

#### Phytochrome Responses 457

Phytochrome responses vary in lag time and escape time 457

Phytochrome responses fall into three main categories based on the amount of light required 457

Phytochrome A mediates responses to continuous far-red light 459

Phytochrome B mediates responses to continuous red or white light 459

Roles for phytochromes C, D, and E are emerging 459

### **Phytochrome Signaling Pathways 459**

Phytochrome regulates membrane potentials and ion fluxes 459

Phytochrome regulates gene expression 460

Phytochrome interacting factors (PIFs) act early in signaling 460

Phytochrome signaling involves protein phosphorylation and dephosphorylation 461

Phytochrome-induced photomorphogenesis involves protein degradation 461

### **Blue-Light Responses and Photoreceptors 462**

Blue-light responses have characteristic kinetics and lag times 462

### **Cryptochromes 463**

The activated FAD chromophore of cryptochrome causes a conformational change in the protein 463

*cry1* and *cry2* have different developmental effects 465

Nuclear cryptochromes inhibit COP1-induced protein degradation 465

Cryptochrome can also bind to transcriptional regulators directly 465

### **The Coaction of Cryptochrome, Phytochrome, and Phototropins 466**

Stem elongation is inhibited by both red and blue photoreceptors 466

Phytochrome interacts with cryptochrome to regulate flowering 467

The circadian clock is regulated by multiple aspects of light 467

### **Phototropins 467**

Blue light induces changes in FMN absorption maxima associated with conformation changes 468

The LOV2 domain is primarily responsible for kinase activation in response to blue light 469

Blue light induces a conformational change that “uncages” the kinase domain of phototropin and leads to autophosphorylation 469

Phototropism requires changes in auxin mobilization 469

Phototropins regulate chloroplast movements via F-actin filament assembly 470

Stomatal opening is regulated by blue light, which activates the plasma membrane H<sup>+</sup>-ATPase 471

The main signal transduction events of phototropin-mediated stomatal opening have been identified 472

### **Responses to Ultraviolet Radiation 473**

## **CHAPTER 17**

### **Embryogenesis 477**

#### **Overview of Plant Growth and Development 478**

Sporophytic development can be divided into three major stages 479

#### **Embryogenesis: The Origins of Polarity 480**

Embryogenesis differs between eudicots and monocots, but also features common fundamental processes 480

Apical-basal polarity is established early in embryogenesis 481

Position-dependent mechanisms guide embryogenesis 483

Intercellular signaling processes play key roles in guiding position-dependent development 484

Embryo development features regulate communication between cells 484

The analysis of mutants identifies genes for signaling processes that are essential for embryo organization 485

Auxin functions as a mobile chemical signal during embryogenesis 487

Plant polarity is maintained by polar auxin streams 487

Auxin transport is regulated by multiple mechanisms 489

The GNOM protein establishes a polar distribution of PIN auxin efflux proteins 491

MONOPTEROS encodes a transcription factor that is activated by auxin 492

Radial patterning guides formation of tissue layers 492

The origin of epidermis: a boundary and interface at the edge of the radial axis 492

Procambial precursors for the vascular stele lie at the center of the radial axis 493

The differentiation of cortical and endodermal cells involves the intercellular movement of a transcription factor 494

#### **Meristematic Tissues: Foundations for Indeterminate Growth 495**

The root and shoot apical meristems use similar strategies to enable indeterminate growth 495

#### **The Root Apical Meristem 496**

The root tip has four developmental zones 497

The origin of different root tissues can be traced to specific initial cells 497

Cell ablation experiments implicate directional signaling processes in determination of cell identity 499

Auxin contributes to the formation and maintenance of the RAM 499

Responses to auxin are mediated by several distinct families of transcription factors 499

Cytokinin is required for normal root development	500
<b>The Shoot Apical Meristem 500</b>	
The shoot apical meristem has distinct zones and layers	502
Shoot tissues are derived from several discrete sets of apical initials	502
Factors involved in auxin movement and responses influence SAM formation	503
Embryonic SAM formation requires the coordinated expression of transcription factors	503
A combination of positive and negative interactions determines apical meristem size	505
KNOX class homeodomain genes help maintain the proliferative ability of the SAM through regulation of cytokinin and GA levels	506
Localized zones of auxin accumulation promote leaf initiation	507
<b>The Vascular Cambium 508</b>	
The maintenance of undetermined initials in various meristem types depends on similar mechanisms	508
<b>CHAPTER 18</b>	
<b>Seed Dormancy, Germination, and Seedling Establishment 513</b>	
<b>Seed Structure 514</b>	
Seed anatomy varies widely among different plant groups	514
<b>Seed Dormancy 515</b>	
Dormancy can be imposed on the embryo by the surrounding tissues	516
Embryo dormancy may be caused by physiological or morphological factors	516
Non-dormant seeds can exhibit vivipary and precocious germination	516
The ABA:GA ratio is the primary determinant of seed dormancy	517
<b>Release from Dormancy 519</b>	
Light is an important signal that breaks dormancy in small seeds	519
Some seeds require either chilling or after-ripening to break dormancy	519
Seed dormancy can be broken by various chemical compounds	520
<b>Seed Germination 520</b>	
Germination can be divided into three phases corresponding to the phases of water uptake	520
<b>Mobilization of Stored Reserves 522</b>	
The cereal aleurone layer is a specialized digestive tissue surrounding the starchy endosperm	522

Gibberellins enhance the transcription of $\alpha$ -amylase mRNA	522
The gibberellin receptor, GID1, promotes the degradation of negative regulators of the gibberellin response	523
GA-MYB is a positive regulator of $\alpha$ -amylase transcription	524
DELLA repressor proteins are rapidly degraded	524
ABA inhibits gibberellin-induced enzyme production	524
<b>Seedling Growth and Establishment 526</b>	
Auxin promotes growth in stems and coleoptiles, while inhibiting growth in roots	526
The outer tissues of eudicot stems are the targets of auxin action	526
The minimum lag time for auxin-induced elongation is 10 minutes	526
Auxin-induced proton extrusion induces cell wall creep and cell elongation	528
<b>Tropisms: Growth in Response to Directional Stimuli 528</b>	
Gravitropism involves the lateral redistribution of auxin	528
Polar auxin transport requires energy and is gravity independent	529
According to the starch–statolith hypothesis, specialized amyloplasts serve as gravity sensors in root caps	530
Auxin movements in the root are regulated by specific transporters	532
The gravitropic stimulus perturbs the symmetric movement of auxin from the root tip	533
Gravity perception in eudicot stems and stemlike organs occurs in the starch sheath	533
Gravity sensing may involve pH and calcium ions ( $\text{Ca}^{2+}$ ) as second messengers	533
<b>Phototropism 535</b>	
Phototropism is mediated by the lateral redistribution of auxin	535
Phototropism occurs in a series of posttranslational events	536
<b>Photomorphogenesis 537</b>	
Gibberellins and brassinosteroids both suppress photomorphogenesis in the dark	538
Hook opening is regulated by phytochrome and auxin	539
Ethylene induces lateral cell expansion	539
<b>Shade Avoidance 540</b>	
Phytochrome enables plants to adapt to changes in light quality	540
Decreasing the R:FR ratio causes elongation in sun plants	540

Reducing shade avoidance responses can improve crop yields 542

### **Vascular Tissue Differentiation 542**

Auxin and cytokinin are required for normal vascular development 543

Zinnia suspension-cultured cells can be induced to undergo xylogenesis 544

Xylogenesis involves chemical signaling between neighboring cells 544

### **Root Growth and Differentiation 545**

Root epidermal development follows three basic patterns 546

Auxin and other hormones regulate root hair development 546

Lateral root formation and emergence depend on endogenous and exogenous signals 547

Regions of lateral root emergence correspond with regions of auxin maxima 548

Lateral roots and shoots have gravitropic setpoint angles 549

## **CHAPTER 19**

### **Vegetative Growth and Organogenesis 553**

#### **Leaf Development 553**

##### **The Establishment of Leaf Polarity 554**

Hormonal signals play key roles in regulating leaf primordia emergence 555

A signal from the SAM initiates adaxial-abaxial polarity 555

ARP genes promote adaxial identity and repress the *KNOX1* gene 556

Adaxial leaf development requires HD-ZIP III transcription factors 556

The expression of HD-ZIP III genes is antagonized by miR166 in abaxial regions of the leaf 558

Antagonism between KANADI and HD-ZIP III is a key determinant of adaxial-abaxial leaf polarity 558

Interactions between adaxial and abaxial tissues are required for blade outgrowth 558

Blade outgrowth is auxin dependent and regulated by the YABBY and WOX genes 558

Leaf proximal-distal polarity also depends on specific gene expression 559

In compound leaves, de-repression of the *KNOX1* gene promotes leaflet formation 559

#### **Differentiation of Epidermal Cell Types 561**

Guard cell fate is ultimately determined by a specialized epidermal lineage 562

Two groups of bHLH transcription factors govern stomatal cell fate transitions 563

Peptide signals regulate stomatal patterning by interacting with cell surface receptors 563

Genetic screens have led to the identification of positive and negative regulators of trichome initiation 563

*GLABRA2* acts downstream of the GL1-GL3-TTG1 complex to promote trichome formation 565

Jasmonic acid regulates *Arabidopsis* leaf trichome development 565

#### **Venation Patterns in Leaves 565**

The primary leaf vein is initiated discontinuously from the preexisting vascular system 566

Auxin canalization initiates development of the leaf trace 566

Basipetal auxin transport from the L1 layer of the leaf primordium initiates development of the leaf trace procambium 568

The existing vasculature guides the growth of the leaf trace 568

Higher-order leaf veins differentiate in a predictable hierarchical order 569

Auxin canalization regulates higher-order vein formation 570

Localized auxin biosynthesis is critical for higher-order venation patterns 571

#### **Shoot Branching and Architecture 572**

Axillary meristem initiation involves many of the same genes as leaf initiation and lamina outgrowth 573

Auxin, cytokinins, and strigolactones regulate axillary bud outgrowth 573

Auxin from the shoot tip maintains apical dominance 574

Strigolactones act locally to repress axillary bud growth 574

Cytokinins antagonize the effects of strigolactones 576

The initial signal for axillary bud growth may be an increase in sucrose availability to the bud 577

Integration of environmental and hormonal branching signals is required for plant fitness 577

Axillary bud dormancy in woody plants is affected by season, position, and age factors 578

#### **Root System Architecture 579**

Plants can modify their root system architecture to optimize water and nutrient uptake 579

Monocots and eudicots differ in their root system architecture 580

Root system architecture changes in response to phosphorous deficiencies 580

Root system architecture responses to phosphorus deficiency involve both local and systemic regulatory networks 582

Mycorrhizal networks augment root system architecture in all major terrestrial ecosystems 583

**Secondary Growth 583**

The vascular cambium and cork cambium are the secondary meristems where secondary growth originates 584

Secondary growth evolved early in the evolution of land plants 585

Secondary growth from the vascular cambium gives rise to secondary xylem and phloem 585

Phytohormones have important roles in regulating vascular cambium activity and differentiation of secondary xylem and phloem 585

Genes involved in stem cell maintenance, proliferation, and differentiation regulate secondary growth 586

Environmental factors influence vascular cambium activity and wood properties 587

**CHAPTER 20****The Control of Flowering and Floral Development 591****Floral Evocation: Integrating Environmental Cues 592****The Shoot Apex and Phase Changes 592**

Plant development has three phases 592

Juvenile tissues are produced first and are located at the base of the shoot 592

Phase changes can be influenced by nutrients, gibberellins, and other signals 593

**Circadian Rhythms: The Clock Within 594**

Circadian rhythms exhibit characteristic features 595

Phase shifting adjusts circadian rhythms to different day-night cycles 596

Phytochromes and cryptochromes entrain the clock 596

**Photoperiodism: Monitoring Day Length 597**

Plants can be classified according to their photoperiodic responses 597

The leaf is the site of perception of the photoperiodic signal 599

Plants monitor day length by measuring the length of the night 599

Night breaks can cancel the effect of the dark period 599

Photoperiodic timekeeping during the night depends on a circadian clock 599

The coincidence model is based on oscillating light sensitivity 600

The coincidence of *CONSTANS* expression and light promotes flowering in LDPs 601

SDPs use a coincidence mechanism to inhibit flowering in long days 603

Phytochrome is the primary photoreceptor in photoperiodism 603

A blue-light photoreceptor regulates flowering in some LDPs 604

**Vernalization: Promoting Flowering with Cold 605**

Vernalization results in competence to flower at the shoot apical meristem 605

Vernalization can involve epigenetic changes in gene expression 606

A range of vernalization pathways may have evolved 607

**Long-Distance Signaling Involved in Flowering 608**

Grafting studies provided the first evidence for a transmissible floral stimulus 608

Florigen is translocated in the phloem 609

**The Identification of Florigen 610**

The *Arabidopsis* protein FLOWERING LOCUS T (FT) is florigen 610

Gibberellins and ethylene can induce flowering 610

The transition to flowering involves multiple factors and pathways 612

**Floral Meristems and Floral Organ Development 612**

The shoot apical meristem in *Arabidopsis* changes with development 613

The four different types of floral organs are initiated as separate whorls 613

Two major categories of genes regulate floral development 614

Floral meristem identity genes regulate meristem function 614

Homeotic mutations led to the identification of floral organ identity genes 616

The ABC model partially explains the determination of floral organ identity 616

*Arabidopsis* Class E genes are required for the activities of the A, B, and C genes 618

According to the Quartet Model, floral organ identity is regulated by tetrameric complexes of the ABCE proteins 618

Class D genes are required for ovule formation 619

Floral asymmetry in flowers is regulated by gene expression 620

<b>CHAPTER 21</b>	Gametophytic self-incompatibility (GSI) is mediated by cytotoxic S-RNases and F-box proteins 642
<b>Gametophytes, Pollination, Seeds, and Fruits 625</b>	<b>Apomixis: Asexual Reproduction by Seed 643</b>
<b>Development of the Male and Female Gametophyte Generations 625</b>	<b>Endosperm Development 643</b>
<b>Formation of Male Gametophytes in the Stamen 626</b>	Cellularization of coenocytic endosperm in <i>Arabidopsis</i> progresses from the micropylar to the chalazal region 645
Pollen grain formation occurs in two successive stages 627	Cellularization of the coenocytic endosperm of cereals progresses centripetally 646
The multilayered pollen cell wall is surprisingly complex 628	Endosperm development and embryogenesis can occur autonomously 646
<b>Female Gametophyte Development in the Ovule 630</b>	Many of the genes that control endosperm development are maternally expressed genes 647
The <i>Arabidopsis</i> gynoecium is an important model system for studying ovule development 630	The FIS proteins are members of a Polycomb repressive complex (PRC2) that represses endosperm development 647
The vast majority of angiosperms exhibit <i>Polygonum</i> -type embryo sac development 630	Cells of the starchy endosperm and aleurone layer follow divergent developmental pathways 649
Functional megasporogenesis undergoes a series of free nuclear mitotic divisions followed by cellularization 631	Two genes, <i>DEK1</i> and <i>CR4</i> , have been implicated in aleurone layer differentiation 649
Embryo sac development involves hormonal signaling between sporophytic and gametophytic generations 632	<b>Seed Coat Development 650</b>
<b>Pollination and Fertilization in Flowering Plants 632</b>	Seed coat development appears to be regulated by the endosperm 650
Delivery of sperm cells to the female gametophyte by the pollen tube occurs in six phases 633	<b>Seed Maturation and Desiccation Tolerance 652</b>
Adhesion and hydration of a pollen grain on a compatible flower depend on recognition between pollen and stigma surfaces 634	Seed filling and desiccation tolerance phases overlap in most species 652
$\text{Ca}^{2+}$ -triggered polarization of the pollen grain precedes tube formation 635	The acquisition of desiccation tolerance involves many metabolic pathways 653
Pollen tubes grow by tip growth 635	During the acquisition of desiccation tolerance, the cells of the embryo acquire a glassy state 653
Receptor-like kinases are thought to regulate the ROP1 GTPase switch, a master regulator of tip growth 635	LEA proteins and nonreducing sugars have been implicated in seed desiccation tolerance 653
Pollen tube tip growth in the pistil is directed by both physical and chemical cues 637	Specific LEA proteins have been implicated in desiccation tolerance in <i>Medicago truncatula</i> 653
Style tissue conditions the pollen tube to respond to attractants produced by the synergids of the embryo sac 637	Abscisic acid plays a key role in seed maturation 654
Double fertilization occurs in three distinct stages 638	Coat-imposed dormancy is correlated with long-term seed-viability 654
<b>Selfing versus Outcrossing 639</b>	<b>Fruit Development and Ripening 655</b>
Hermaphroditic and monoecious species have evolved floral features to ensure outcrossing 639	<i>Arabidopsis</i> and tomato are model systems for the study of fruit development 655
Cytoplasmic male sterility (CMS) occurs in the wild and is of great utility in agriculture 640	Fleshy fruits undergo ripening 657
Self-incompatibility (SI) is the primary mechanism that enforces outcrossing in angiosperms 640	Ripening involves changes in the color of fruit 657
The Brassicaceae sporophytic SI system requires two S-locus genes 641	Fruit softening involves the coordinated action of many cell wall-degrading enzymes 658
	Taste and flavor reflect changes in acids, sugars, and aroma compounds 658
	The causal link between ethylene and ripening was demonstrated in transgenic and mutant tomatoes 658
	Climacteric and non-climacteric fruit differ in their ethylene responses 658
	The ripening process is transcriptionally regulated 660

Angiosperms share a range of common molecular mechanisms controlling fruit development and ripening 660  
 Fruit ripening is under epigenetic control 660  
 A mechanistic understanding of the ripening process has commercial applications 661

## CHAPTER 22

### Plant Senescence and Cell Death 665

**Programmed Cell Death and Autolysis 666**  
 PCD during normal development differs from that of the hypersensitive response 668  
 The autophagy pathway captures and degrades cellular constituents within lytic compartments 669  
 A subset of the autophagy-related genes controls the formation of the autophagosome 669  
 The autophagy pathway plays a dual role in plant development 671

#### The Leaf Senescence Syndrome 671

The developmental age of a leaf may differ from its chronological age 672  
 Leaf senescence may be sequential, seasonal, or stress-induced 672  
 Developmental leaf senescence consists of three distinct phases 673  
 The earliest cellular changes during leaf senescence occur in the chloroplast 675  
 The autolysis of chloroplast proteins occurs in multiple compartments 675  
 The STAY-GREEN (SGR) protein is required for both LHCP II protein recycling and chlorophyll catabolism 676  
 Leaf senescence is preceded by a massive reprogramming of gene expression 677

#### Leaf Senescence: The Regulatory Network 678

The NAC and WRKY gene families are the most abundant transcription factors regulating leaf senescence 678  
 ROS serve as internal signaling agents in leaf senescence 680  
 Sugars accumulate during leaf senescence and may serve as a signal 681  
 Plant hormones interact in the regulation of leaf senescence 681

#### Leaf Abscission 684

The timing of leaf abscission is regulated by the interaction of ethylene and auxin 685

#### Whole Plant Senescence 686

Angiosperm life cycles may be annual, biennial, or perennial 687

Whole plant senescence differs from aging in animals 688  
 The determinacy of shoot apical meristems is developmentally regulated 688  
 Nutrient or hormonal redistribution may trigger senescence in monocarpic plants 689  
 The rate of carbon accumulation in trees increases continuously with tree size 689

## CHAPTER 23

### Biotic Interactions 693

#### Beneficial Interactions between Plants and Microorganisms 695

Nod factors are recognized by the Nod factor receptor (NFR) in legumes 695  
 Arbuscular mycorrhizal associations and nitrogen-fixing symbioses involve related signaling pathways 695  
 Rhizobacteria can increase nutrient availability, stimulate root branching, and protect against pathogens 697

#### Harmful Interactions between Plants, Pathogens, and Herbivores 697

Mechanical barriers provide a first line of defense against insect pests and pathogens 698  
 Plant secondary metabolites can deter insect herbivores 700  
 Plants store constitutive toxic compounds in specialized structures 701  
 Plants often store defensive chemicals as nontoxic water-soluble sugar conjugates in the vacuole 703  
 Constitutive levels of secondary compounds are higher in young developing leaves than in older tissues 705

#### Inducible Defense Responses to Insect Herbivores 705

Plants can recognize specific components of insect saliva 706  
 Modified fatty acids secreted by grasshoppers act as elicitors of jasmonic acid accumulation and ethylene emission 706  
 Phloem feeders activate defense signaling pathways similar to those activated by pathogen infections 707  
 Calcium signaling and activation of the MAP kinase pathway are early events associated with insect herbivory 707  
 Jasmonic acid activates defense responses against insect herbivores 708  
 Jasmonic acid acts through a conserved ubiquitin ligase signaling mechanism 709  
 Hormonal interactions contribute to plant-insect herbivore interactions 709

- JA initiates the production of defense proteins that inhibit herbivore digestion 710
- Herbivore damage induces systemic defenses 710
- Glutamate receptor-like (GLR) genes are required for long-distance electrical signaling during herbivory 712
- Herbivore-induced volatiles can repel herbivores and attract natural enemies 712
- Herbivore-induced volatiles can serve as long-distance signals between plants 713
- Herbivore-induced volatiles can also act as systemic signals within a plant 714
- Defense responses to herbivores and pathogens are regulated by circadian rhythms 714
- Insects have evolved mechanisms to defeat plant defenses 715

### **Plant Defenses against Pathogens 715**

- Microbial pathogens have evolved various strategies to invade host plants 715
- Pathogens produce effector molecules that aid in the colonization of their plant host cells 716
- Pathogen infection can give rise to molecular “danger signals” that are perceived by cell surface pattern recognition receptors (PRRs) 717
- R genes provide resistance to individual pathogens by recognizing strain-specific effectors 718
- Exposure to elicitors induces a signal transduction cascade 719
- Effectors released by phloem-feeding insects also activate NBS-LRR receptors 719
- The hypersensitive response is a common defense against pathogens 720
- Phytoalexins with antimicrobial activity accumulate after pathogen attack 721
- A single encounter with a pathogen may increase resistance to future attacks 721
- The main components of the salicylic acid signaling pathway for SAR have been identified 723
- Interactions of plants with nonpathogenic bacteria can trigger systemic resistance through a process called induced systemic resistance (ISR) 723

### **Plant Defenses against Other Organisms 724**

- Some plant parasitic nematodes form specific associations through the formation of distinct feeding structures 724
- Plants compete with other plants by secreting allelopathic secondary metabolites into the soil 725
- Some plants are biotrophic pathogens of other plants 726

## **CHAPTER 24**

### **Abiotic Stress 731**

#### **Defining Plant Stress 732**

Physiological adjustment to abiotic stress involves trade-offs between vegetative and reproductive development 732

#### **Acclimation and Adaptation 733**

- Adaptation to stress involves genetic modification over many generations 733
- Acclimation allows plants to respond to environmental fluctuations 733

#### **Environmental Factors and Their Biological Impacts on Plants 734**

- Water deficit decreases turgor pressure, increases ion toxicity, and inhibits photosynthesis 735
- Salinity stress has both osmotic and cytotoxic effects 736
- Light stress can occur when shade-adapted or shade-acclimated plants are subjected to full sunlight 736
- Temperature stress affects a broad spectrum of physiological processes 736
- Flooding results in anaerobic stress to the root 737
- During freezing stress, extracellular ice crystal formation causes cell dehydration 737
- Heavy metals can both mimic essential mineral nutrients and generate ROS 737
- Mineral nutrient deficiencies are a cause of stress 737
- Ozone and ultraviolet light generate ROS that cause lesions and induce PCD 737
- Combinations of abiotic stresses can induce unique signaling and metabolic pathways 738
- Sequential exposure to different abiotic stresses sometimes confers cross-protection 739

#### **Stress-Sensing Mechanisms in Plants 739**

Early-acting stress sensors provide the initial signal for the stress response 740

#### **Signaling Pathways Activated in Response to Abiotic Stress 740**

- The signaling intermediates of many stress-response pathways can interact 740
- Acclimation to stress involves transcriptional regulatory networks called *regulons* 743
- Chloroplast genes respond to high-intensity light by sending stress signals to the nucleus 744
- A self-propagating wave of ROS mediates systemic acquired acclimation 745
- Epigenetic mechanisms and small RNAs provide additional protection against stress 745

Hormonal interactions regulate normal development and abiotic stress responses 745

## **Developmental and Physiological Mechanisms That Protect Plants against Abiotic Stress 747**

Plants adjust osmotically to drying soil by accumulating solutes 748

Submerged organs develop aerenchyma tissue in response to hypoxia 749

Antioxidants and ROS-scavenging pathways protect cells from oxidative stress 750

Molecular chaperones and molecular shields protect proteins and membranes during abiotic stress 751

Plants can alter their membrane lipids in response to temperature and other abiotic stresses 752

Exclusion and internal tolerance mechanisms allow plants to cope with toxic ions 753

Phytochelatins and other chelators contribute to internal tolerance of toxic metal ions 754

Plants use cryoprotectant molecules and antifreeze proteins to prevent ice crystal formation 754

ABA signaling during water stress causes the massive efflux of K<sup>+</sup> and anions from guard cells 755

Plants can alter their morphology in response to abiotic stress 757

Metabolic shifts enable plants to cope with a variety of abiotic stresses 759

The process of recovery from stress can be dangerous to the plant and requires a coordinated adjustment of plant metabolism and physiology 759

Developing crops with enhanced tolerance to abiotic stress conditions is a major goal of agricultural research 759

---

## **Glossary G-1**

## **Illustration Credits IC-1**

## **Photo Credits PC-1**

## **Subject Index SI-1**