

Contents

Introduction	1
Outline	8
Some notations and assumptions	15
Chapter I. Geometry of the Mumford-Tate domains	17
I.A. Generalities on homogeneous complex manifolds	17
I.B. The first example ($\mathcal{U}(2, 1)$)	18
I.C. The second example ($\mathrm{Sp}(4)$)	28
Chapter II. Homogeneous line bundles over the Mumford-Tate domains	37
II.A. Generalities on homogeneous vector bundles	37
II.B. Homogeneous line bundles over the first example	37
Appendix to section II.B: Sign conventions	44
II.C. Homogeneous line bundles in the second example	45
Chapter III. Correspondence and cycle spaces; Penrose transforms	49
III.A. Introduction	49
III.B. Basic definitions and examples	49
III.C. The basic example	59
III.D. The Penrose transform in the compact case	68
Appendix to section III.D: Arithmetic aspects of the Penrose transform in the compact case	73
III.E. The Penrose transform in the first example	77
III.F. The Penrose transform in the second example	86
Chapter IV. The Penrose transform in the automorphic case and the main result	93
IV.A. Cuspidal automorphic cohomology	93
IV.B. Picard and Siegel cuspidal automorphic forms	98
IV.C. Arithmetic structures on vector spaces	102
Appendix to section IV.C: Explicit canonical models for the two examples	108
IV.D. Special values of cuspidal automorphic cohomology classes	110
Appendix to section IV.D: An alternate method for evaluating cohomology classes and a question	113
IV.E. CM points on correspondence spaces	117
IV.F. On a result of Carayol	123
Appendix to section IV.F: Geometric construction of K -types and discussion of totally degenerate limits of discrete series	130
Bibliography	143