

# BRIEF CONTENTS

<b>Part I THE MOLECULAR DESIGN OF LIFE</b>	
1 Biochemistry: An Evolving Science	1
2 Protein Composition and Structure	27
3 Exploring Proteins and Proteomes	65
4 DNA, RNA, and the Flow of Genetic Information	105
5 Exploring Genes and Genomes	135
6 Exploring Evolution and Bioinformatics	169
7 Hemoglobin: Portrait of a Protein in Action	191
8 Enzymes: Basic Concepts and Kinetics	215
9 Catalytic Strategies	251
10 Regulatory Strategies	285
11 Carbohydrates	315
12 Lipids and Cell Membranes	341
13 Membrane Channels and Pumps	367
14 Signal-Transduction Pathways	397
<b>Part II TRANSDUCING AND STORING ENERGY</b>	
15 Metabolism: Basic Concepts and Design	423
16 Glycolysis and Gluconeogenesis	449
17 The Citric Acid Cycle	495
18 Oxidative Phosphorylation	523
19 The Light Reactions of Photosynthesis	565
20 The Calvin Cycle and the Pentose Phosphate Pathway	589
21 Glycogen Metabolism	617
22 Fatty Acid Metabolism	643
23 Protein Turnover and Amino Acid Catabolism	681
<b>Part III SYNTHESIZING THE MOLECULES OF LIFE</b>	
24 The Biosynthesis of Amino Acids	713
25 Nucleotide Biosynthesis	743
26 The Biosynthesis of Membrane Lipids and Steroids	767
27 The Integration of Metabolism	801
28 DNA Replication, Repair, and Recombination	827
29 RNA Synthesis and Processing	859
30 Protein Synthesis	893
31 The Control of Gene Expression in Prokaryotes	925
32 The Control of Gene Expression in Eukaryotes	941
<b>Part IV RESPONDING TO ENVIRONMENTAL CHANGES</b>	
33 Sensory Systems	961
34 The Immune System	981
35 Molecular Motors	1011
36 Drug Development	1033

# CONTENTS

Preface	v
<b>Part I THE MOLECULAR DESIGN OF LIFE</b>	<b>1</b>
<b>CHAPTER 1 Biochemistry: An Evolving Science</b>	<b>1</b>
1.1 Biochemical Unity Underlies Biological Diversity	1
1.2 DNA Illustrates the Interplay Between Form and Function	4
DNA is constructed from four building blocks	4
Two single strands of DNA combine to form a double helix	5
DNA structure explains heredity and the storage of information	5
1.3 Concepts from Chemistry Explain the Properties of Biological Molecules	6
The formation of the DNA double helix as a key example	6
The double helix can form from its component strands	6
Covalent and noncovalent bonds are important for the structure and stability of biological molecules	6
The double helix is an expression of the rules of chemistry	9
The laws of thermodynamics govern the behavior of biochemical systems	10
Heat is released in the formation of the double helix	12
Acid-base reactions are central in many biochemical processes	13
Acid-base reactions can disrupt the double helix	14
Buffers regulate pH in organisms and in the laboratory	15
1.4 The Genomic Revolution Is Transforming Biochemistry, Medicine, and Other Fields	17
Genome sequencing has transformed biochemistry and other fields	17
Environmental factors influence human biochemistry	20
Genome sequences encode proteins and patterns of expression	21
APPENDIX: Visualizing Molecular Structures I: Small Molecules	22
<b>CHAPTER 2 Protein Composition and Structure</b>	<b>27</b>
2.1 Proteins Are Built from a Repertoire of 20 Amino Acids	29
2.2 Primary Structure: Amino Acids Are Linked by Peptide Bonds to Form Polypeptide Chains	35
Proteins have unique amino acid sequences specified by genes	37
Polypeptide chains are flexible yet conformationally restricted	38
2.3 Secondary Structure: Polypeptide Chains Can Fold into Regular Structures Such As the Alpha Helix, the Beta Sheet, and Turns and Loops	40
The alpha helix is a coiled structure stabilized by intrachain hydrogen bonds	40
Beta sheets are stabilized by hydrogen bonding between polypeptide strands	42

Polypeptide chains can change direction by making reverse turns and loops	44		
Fibrous proteins provide structural support for cells and tissues	44		
<b>2.4 Tertiary Structure: Water-Soluble Proteins Fold into Compact Structures with Nonpolar Cores</b>	<b>46</b>		
<b>2.5 Quaternary Structure: Polypeptide Chains Can Assemble into Multisubunit Structures</b>	<b>48</b>		
<b>2.6 The Amino Acid Sequence of a Protein Determines Its Three-Dimensional Structure</b>	<b>49</b>		
Amino acids have different propensities for forming $\alpha$ helices, $\beta$ sheets, and turns	51		
Protein folding is a highly cooperative process	52		
Proteins fold by progressive stabilization of intermediates rather than by random search	53		
Prediction of three-dimensional structure from sequence remains a great challenge	54		
Some proteins are inherently unstructured and can exist in multiple conformations	55		
Protein misfolding and aggregation are associated with some neurological diseases	56		
Protein modification and cleavage confer new capabilities	57		
APPENDIX: Visualizing Molecular Structures II: Proteins	61		
<b>CHAPTER 3 Exploring Proteins and Proteomes</b>	<b>65</b>		
The proteome is the functional representation of the genome	66		
<b>3.1 The Purification of Proteins Is an Essential First Step in Understanding Their Function</b>	<b>66</b>		
The assay: How do we recognize the protein that we are looking for?	67		
Proteins must be released from the cell to be purified	67		
Proteins can be purified according to solubility, size, charge, and binding affinity	68		
Proteins can be separated by gel electrophoresis and displayed	71		
A protein purification scheme can be quantitatively evaluated	75		
Ultracentrifugation is valuable for separating biomolecules and determining their masses	76		
Protein purification can be made easier with the use of recombinant DNA technology	78		
<b>3.2 Immunology Provides Important Techniques with Which to Investigate Proteins</b>	<b>79</b>		
Antibodies to specific proteins can be generated	79		
Monoclonal antibodies with virtually any desired specificity can be readily prepared	80		
Proteins can be detected and quantified by using an enzyme-linked immunosorbent assay	82		
Western blotting permits the detection of proteins separated by gel electrophoresis	83		
Fluorescent markers make the visualization of proteins in the cell possible	84		
		<b>3.3 Mass Spectrometry Is a Powerful Technique for the Identification of Peptides and Proteins</b>	<b>85</b>
		Peptides can be sequenced by mass spectrometry	87
		Proteins can be specifically cleaved into small peptides to facilitate analysis	88
		Genomic and proteomic methods are complementary	89
		The amino acid sequence of a protein provides valuable information	90
		Individual proteins can be identified by mass spectrometry	91
		<b>3.4 Peptides Can Be Synthesized by Automated Solid-Phase Methods</b>	<b>92</b>
		<b>3.5 Three-Dimensional Protein Structure Can Be Determined by X-ray Crystallography and NMR Spectroscopy</b>	<b>95</b>
		X-ray crystallography reveals three-dimensional structure in atomic detail	95
		Nuclear magnetic resonance spectroscopy can reveal the structures of proteins in solution	97
		<b>CHAPTER 4 DNA, RNA, and the Flow of Genetic Information</b>	<b>105</b>
		<b>4.1 A Nucleic Acid Consists of Four Kinds of Bases Linked to a Sugar-Phosphate Backbone</b>	<b>106</b>
		RNA and DNA differ in the sugar component and one of the bases	106
		Nucleotides are the monomeric units of nucleic acids	107
		DNA molecules are very long and have directionality	108
		<b>4.2 A Pair of Nucleic Acid Strands with Complementary Sequences Can Form a Double-Helical Structure</b>	<b>109</b>
		The double helix is stabilized by hydrogen bonds and van der Waals interactions	109
		DNA can assume a variety of structural forms	111
		Z-DNA is a left-handed double helix in which backbone phosphates zigzag	112
		Some DNA molecules are circular and supercoiled	113
		Single-stranded nucleic acids can adopt elaborate structures	113
		<b>4.3 The Double Helix Facilitates the Accurate Transmission of Hereditary Information</b>	<b>114</b>
		Differences in DNA density established the validity of the semiconservative replication hypothesis	115
		The double helix can be reversibly melted	116
		<b>4.4 DNA Is Replicated by Polymerases That Take Instructions from Templates</b>	<b>117</b>
		DNA polymerase catalyzes phosphodiester-bridge formation	117
		The genes of some viruses are made of RNA	118
		<b>4.5 Gene Expression Is the Transformation of DNA Information into Functional Molecules</b>	<b>119</b>
		Several kinds of RNA play key roles in gene expression	119

All cellular RNA is synthesized by RNA polymerases	120	Next-generation sequencing methods enable the rapid determination of a complete genome sequence	155
RNA polymerases take instructions from DNA templates	121	Comparative genomics has become a powerful research tool	156
Transcription begins near promoter sites and ends at terminator sites	122	<b>5.4 Eukaryotic Genes Can Be Quantitated and Manipulated with Considerable Precision</b>	157
Transfer RNAs are the adaptor molecules in protein synthesis	123	Gene-expression levels can be comprehensively examined	157
<b>4.6 Amino Acids Are Encoded by Groups of Three Bases Starting from a Fixed Point</b>	124	New genes inserted into eukaryotic cells can be efficiently expressed	159
Major features of the genetic code	125	Transgenic animals harbor and express genes introduced into their germ lines	160
Messenger RNA contains start and stop signals for protein synthesis	126	Gene disruption and genome editing provide clues to gene function and opportunities for new therapies	160
The genetic code is nearly universal	126	RNA interference provides an additional tool for disrupting gene expression	162
<b>4.7 Most Eukaryotic Genes Are Mosaics of Introns and Exons</b>	127	Tumor-inducing plasmids can be used to introduce new genes into plant cells	163
RNA processing generates mature RNA	127	Human gene therapy holds great promise for medicine	164
Many exons encode protein domains	128		
<b>CHAPTER 5 Exploring Genes and Genomes</b>	<b>135</b>	<b>CHAPTER 6 Exploring Evolution and Bioinformatics</b>	<b>169</b>
<b>5.1 The Exploration of Genes Relies on Key Tools</b>	<b>136</b>	<b>6.1 Homologs Are Descended from a Common Ancestor</b>	<b>170</b>
Restriction enzymes split DNA into specific fragments	137	<b>6.2 Statistical Analysis of Sequence Alignments Can Detect Homology</b>	<b>171</b>
Restriction fragments can be separated by gel electrophoresis and visualized	137	The statistical significance of alignments can be estimated by shuffling	173
DNA can be sequenced by controlled termination of replication	138	Distant evolutionary relationships can be detected through the use of substitution matrices	174
DNA probes and genes can be synthesized by automated solid-phase methods	139	Databases can be searched to identify homologous sequences	177
Selected DNA sequences can be greatly amplified by the polymerase chain reaction	141	<b>6.3 Examination of Three-Dimensional Structure Enhances Our Understanding of Evolutionary Relationships</b>	<b>177</b>
PCR is a powerful technique in medical diagnostics, forensics, and studies of molecular evolution	142	Tertiary structure is more conserved than primary structure	178
The tools for recombinant DNA technology have been used to identify disease-causing mutations	143	Knowledge of three-dimensional structures can aid in the evaluation of sequence alignments	179
<b>5.2 Recombinant DNA Technology Has Revolutionized All Aspects of Biology</b>	<b>143</b>	Repeated motifs can be detected by aligning sequences with themselves	180
Restriction enzymes and DNA ligase are key tools in forming recombinant DNA molecules	143	Convergent evolution illustrates common solutions to biochemical challenges	181
Plasmids and $\lambda$ phage are choice vectors for DNA cloning in bacteria	144	Comparison of RNA sequences can be a source of insight into RNA secondary structures	182
Bacterial and yeast artificial chromosomes	147	<b>6.4 Evolutionary Trees Can Be Constructed on the Basis of Sequence Information</b>	<b>183</b>
Specific genes can be cloned from digests of genomic DNA	147	Horizontal gene transfer events may explain unexpected branches of the evolutionary tree	184
Complementary DNA prepared from mRNA can be expressed in host cells	149	<b>6.5 Modern Techniques Make the Experimental Exploration of Evolution Possible</b>	<b>185</b>
Proteins with new functions can be created through directed changes in DNA	150	Ancient DNA can sometimes be amplified and sequenced	185
Recombinant methods enable the exploration of the functional effects of disease-causing mutations	152	Molecular evolution can be examined experimentally	185
<b>5.3 Complete Genomes Have Been Sequenced and Analyzed</b>	<b>152</b>		
The genomes of organisms ranging from bacteria to multicellular eukaryotes have been sequenced	153		
The sequence of the human genome has been completed	154		

<b>CHAPTER 7 Hemoglobin: Portrait of a Protein in Action</b>	<b>191</b>		
<b>7.1 Myoglobin and Hemoglobin Bind Oxygen at Iron Atoms in Heme</b>	<b>192</b>		
Changes in heme electronic structure upon oxygen binding are the basis for functional imaging studies	193		
The structure of myoglobin prevents the release of reactive oxygen species	194		
Human hemoglobin is an assembly of four myoglobin-like subunits	195		
<b>7.2 Hemoglobin Binds Oxygen Cooperatively</b>	<b>195</b>		
Oxygen binding markedly changes the quaternary structure of hemoglobin	197		
Hemoglobin cooperativity can be potentially explained by several models	198		
Structural changes at the heme groups are transmitted to the $\alpha_1\beta_1$ - $\alpha_2\beta_2$ interface	200		
2,3-Bisphosphoglycerate in red cells is crucial in determining the oxygen affinity of hemoglobin	200		
Carbon monoxide can disrupt oxygen transport by hemoglobin	201		
<b>7.3 Hydrogen Ions and Carbon Dioxide Promote the Release of Oxygen: The Bohr Effect</b>	<b>202</b>		
<b>7.4 Mutations in Genes Encoding Hemoglobin Subunits Can Result in Disease</b>	<b>204</b>		
Sickle-cell anemia results from the aggregation of mutated deoxyhemoglobin molecules	205		
Thalassemia is caused by an imbalanced production of hemoglobin chains	207		
The accumulation of free alpha-hemoglobin chains is prevented	207		
Additional globins are encoded in the human genome	208		
APPENDIX: Binding Models Can Be Formulated in Quantitative Terms: The Hill Plot and the Concerted Model	210		
<b>CHAPTER 8 Enzymes: Basic Concepts and Kinetics</b>	<b>215</b>		
<b>8.1 Enzymes are Powerful and Highly Specific Catalysts</b>	<b>216</b>		
Many enzymes require cofactors for activity	217		
Enzymes can transform energy from one form into another	217		
<b>8.2 Gibbs Free Energy Is a Useful Thermodynamic Function for Understanding Enzymes</b>	<b>218</b>		
The free-energy change provides information about the spontaneity but not the rate of a reaction	218		
The standard free-energy change of a reaction is related to the equilibrium constant	219		
Enzymes alter only the reaction rate and not the reaction equilibrium	220		
<b>8.3 Enzymes Accelerate Reactions by Facilitating the Formation of the Transition State</b>	<b>221</b>		
		The formation of an enzyme-substrate complex is the first step in enzymatic catalysis	222
		The active sites of enzymes have some common features	223
		The binding energy between enzyme and substrate is important for catalysis	225
		<b>8.4 The Michaelis-Menten Model Accounts for the Kinetic Properties of Many Enzymes</b>	<b>225</b>
		Kinetics is the study of reaction rates	225
		The steady-state assumption facilitates a description of enzyme kinetics	226
		Variations in $K_M$ can have physiological consequences	228
		$K_M$ and $V_{max}$ values can be determined by several means	228
		$K_M$ and $V_{max}$ values are important enzyme characteristics	229
		$k_{cat}/K_M$ is a measure of catalytic efficiency	230
		Most biochemical reactions include multiple substrates	231
		Allosteric enzymes do not obey Michaelis-Menten kinetics	233
		<b>8.5 Enzymes Can Be Inhibited by Specific Molecules</b>	<b>234</b>
		The different types of reversible inhibitors are kinetically distinguishable	235
		Irreversible inhibitors can be used to map the active site	237
		Penicillin irreversibly inactivates a key enzyme in bacterial cell-wall synthesis	239
		Transition-state analogs are potent inhibitors of enzymes	240
		Catalytic antibodies demonstrate the importance of selective binding of the transition state to enzymatic activity	241
		<b>8.6 Enzymes Can Be Studied One Molecule at a Time</b>	<b>242</b>
		APPENDIX: Enzymes are Classified on the Basis of the Types of Reactions That They Catalyze	245
		<b>CHAPTER 9 Catalytic Strategies</b>	<b>251</b>
		A few basic catalytic principles are used by many enzymes	252
		<b>9.1 Proteases Facilitate a Fundamentally Difficult Reaction</b>	<b>253</b>
		Chymotrypsin possesses a highly reactive serine residue	253
		Chymotrypsin action proceeds in two steps linked by a covalently bound intermediate	254
		Serine is part of a catalytic triad that also includes histidine and aspartate	255
		Catalytic triads are found in other hydrolytic enzymes	258
		The catalytic triad has been dissected by site-directed mutagenesis	260
		Cysteine, aspartyl, and metalloproteases are other major classes of peptide-cleaving enzymes	260
		Protease inhibitors are important drugs	263

<b>9.2 Carbonic Anhydrases Make a Fast Reaction Faster</b>	264	Cyclic AMP activates protein kinase A by altering the quaternary structure	297
Carbonic anhydrase contains a bound zinc ion essential for catalytic activity	265	ATP and the target protein bind to a deep cleft in the catalytic subunit of protein kinase A	298
Catalysis entails zinc activation of a water molecule	265	<b>10.4 Many Enzymes Are Activated by Specific Proteolytic Cleavage</b>	299
A proton shuttle facilitates rapid regeneration of the active form of the enzyme	267	Chymotrypsinogen is activated by specific cleavage of a single peptide bond	299
<b>9.3 Restriction Enzymes Catalyze Highly Specific DNA-Cleavage Reactions</b>	269	Proteolytic activation of chymotrypsinogen leads to the formation of a substrate-binding site	300
Cleavage is by in-line displacement of 3'-oxygen from phosphorus by magnesium-activated water	269	The generation of trypsin from trypsinogen leads to the activation of other zymogens	301
Restriction enzymes require magnesium for catalytic activity	271	Some proteolytic enzymes have specific inhibitors	302
The complete catalytic apparatus is assembled only within complexes of cognate DNA molecules, ensuring specificity	272	Blood clotting is accomplished by a cascade of zymogen activations	303
Host-cell DNA is protected by the addition of methyl groups to specific bases	274	Prothrombin requires a vitamin K-dependent modification for activation	304
Type II restriction enzymes have a catalytic core in common and are probably related by horizontal gene transfer	275	Fibrinogen is converted by thrombin into a fibrin clot	304
<b>9.4 Myosins Harness Changes in Enzyme Conformation to Couple ATP Hydrolysis to Mechanical Work</b>	275	Vitamin K is required for the formation of $\gamma$ -carboxylglutamate	306
ATP hydrolysis proceeds by the attack of water on the gamma-phosphoryl group	276	The clotting process must be precisely regulated	307
Formation of the transition state for ATP hydrolysis is associated with a substantial conformational change	277	Hemophilia revealed an early step in clotting	308
The altered conformation of myosin persists for a substantial period of time	278	<b>CHAPTER 11 Carbohydrates</b>	<b>315</b>
Scientists can watch single molecules of myosin move	279	<b>11.1 Monosaccharides Are the Simplest Carbohydrates</b>	<b>316</b>
Myosins are a family of enzymes containing P-loop structures	280	Many common sugars exist in cyclic forms	318
<b>CHAPTER 10 Regulatory Strategies</b>	<b>285</b>	Pyranose and furanose rings can assume different conformations	320
<b>10.1 Aspartate Transcarbamoylase Is Allosterically Inhibited by the End Product of Its Pathway</b>	<b>286</b>	Glucose is a reducing sugar	321
Allosterically regulated enzymes do not follow Michaelis-Menten kinetics	287	Monosaccharides are joined to alcohols and amines through glycosidic bonds	322
ATCase consists of separable catalytic and regulatory subunits	287	Phosphorylated sugars are key intermediates in energy generation and biosyntheses	322
Allosteric interactions in ATCase are mediated by large changes in quaternary structure	288	<b>11.2 Monosaccharides Are Linked to Form Complex Carbohydrates</b>	<b>323</b>
Allosteric regulators modulate the T-to-R equilibrium	291	Sucrose, lactose, and maltose are the common disaccharides	323
<b>10.2 Isozymes Provide a Means of Regulation Specific to Distinct Tissues and Developmental Stages</b>	<b>292</b>	Glycogen and starch are storage forms of glucose	324
<b>10.3 Covalent Modification Is a Means of Regulating Enzyme Activity</b>	<b>293</b>	Cellulose, a structural component of plants, is made of chains of glucose	324
Kinases and phosphatases control the extent of protein phosphorylation	294	<b>11.3 Carbohydrates Can Be Linked to Proteins to Form Glycoproteins</b>	<b>325</b>
Phosphorylation is a highly effective means of regulating the activities of target proteins	296	Carbohydrates can be linked to proteins through asparagine ( <i>N</i> -linked) or through serine or threonine ( <i>O</i> -linked) residues	326
		The glycoprotein erythropoietin is a vital hormone	327
		Glycosylation functions in nutrient sensing	327
		Proteoglycans, composed of polysaccharides and protein, have important structural roles	327
		Proteoglycans are important components of cartilage	328
		Mucins are glycoprotein components of mucus	329
		Protein glycosylation takes place in the lumen of the endoplasmic reticulum and in the Golgi complex	330

Specific enzymes are responsible for oligosaccharide assembly	331	<b>12.6 Eukaryotic Cells Contain Compartments Bounded by Internal Membranes</b>	<b>359</b>
Blood groups are based on protein glycosylation patterns	331		
Errors in glycosylation can result in pathological conditions	332	<b>CHAPTER 13 Membrane Channels and Pumps</b>	<b>367</b>
Oligosaccharides can be “sequenced”	332	The expression of transporters largely defines the metabolic activities of a given cell type	368
<b>11.4 Lectins Are Specific Carbohydrate-Binding Proteins</b>	<b>333</b>	<b>13.1 The Transport of Molecules Across a Membrane May Be Active or Passive</b>	<b>368</b>
Lectins promote interactions between cells	334	Many molecules require protein transporters to cross membranes	368
Lectins are organized into different classes	334	Free energy stored in concentration gradients can be quantified	369
Influenza virus binds to sialic acid residues	335	<b>13.2 Two Families of Membrane Proteins Use ATP Hydrolysis to Pump Ions and Molecules Across Membranes</b>	<b>370</b>
<b>CHAPTER 12 Lipids and Cell Membranes</b>	<b>341</b>	P-type ATPases couple phosphorylation and conformational changes to pump calcium ions across membranes	370
Many common features underlie the diversity of biological membranes	342	Digitalis specifically inhibits the Na <sup>+</sup> -K <sup>+</sup> pump by blocking its dephosphorylation	373
<b>12.1 Fatty Acids Are Key Constituents of Lipids</b>	<b>342</b>	P-type ATPases are evolutionarily conserved and play a wide range of roles	374
Fatty acid names are based on their parent hydrocarbons	342	Multidrug resistance highlights a family of membrane pumps with ATP-binding cassette domains	374
Fatty acids vary in chain length and degree of unsaturation	343	<b>13.3 Lactose Permease Is an Archetype of Secondary Transporters That Use One Concentration Gradient to Power the Formation of Another</b>	<b>376</b>
<b>12.2 There Are Three Common Types of Membrane Lipids</b>	<b>344</b>	<b>13.4 Specific Channels Can Rapidly Transport Ions Across Membranes</b>	<b>378</b>
Phospholipids are the major class of membrane lipids	344	Action potentials are mediated by transient changes in Na <sup>+</sup> and K <sup>+</sup> permeability	378
Membrane lipids can include carbohydrate moieties	345	Patch-clamp conductance measurements reveal the activities of single channels	379
Cholesterol is a lipid based on a steroid nucleus	346	The structure of a potassium ion channel is an archetype for many ion-channel structures	379
Archaeal membranes are built from ether lipids with branched chains	346	The structure of the potassium ion channel reveals the basis of ion specificity	380
A membrane lipid is an amphipathic molecule containing a hydrophilic and a hydrophobic moiety	347	The structure of the potassium ion channel explains its rapid rate of transport	383
<b>12.3 Phospholipids and Glycolipids Readily Form Bimolecular Sheets in Aqueous Media</b>	<b>348</b>	Voltage gating requires substantial conformational changes in specific ion-channel domains	383
Lipid vesicles can be formed from phospholipids	348	A channel can be inactivated by occlusion of the pore: the ball-and-chain model	384
Lipid bilayers are highly impermeable to ions and most polar molecules	349	The acetylcholine receptor is an archetype for ligand-gated ion channels	385
<b>12.4 Proteins Carry Out Most Membrane Processes</b>	<b>350</b>	Action potentials integrate the activities of several ion channels working in concert	387
Proteins associate with the lipid bilayer in a variety of ways	351	Disruption of ion channels by mutations or chemicals can be potentially life-threatening	388
Proteins interact with membranes in a variety of ways	351	<b>13.5 Gap Junctions Allow Ions and Small Molecules to Flow Between Communicating Cells</b>	<b>389</b>
Some proteins associate with membranes through covalently attached hydrophobic groups	354	<b>13.6 Specific Channels Increase the Permeability of Some Membranes to Water</b>	<b>390</b>
Transmembrane helices can be accurately predicted from amino acid sequences	354		
<b>12.5 Lipids and Many Membrane Proteins Diffuse Rapidly in the Plane of the Membrane</b>	<b>356</b>		
The fluid mosaic model allows lateral movement but not rotation through the membrane	357		
Membrane fluidity is controlled by fatty acid composition and cholesterol content	357		
Lipid rafts are highly dynamic complexes formed between cholesterol and specific lipids	358		
All biological membranes are asymmetric	358		

<b>CHAPTER 14 Signal-Transduction Pathways</b>	<b>397</b>		
Signal transduction depends on molecular circuits	398		
<b>14.1 Heterotrimeric G Proteins Transmit Signals and Reset Themselves</b>	<b>399</b>		
Ligand binding to 7TM receptors leads to the activation of heterotrimeric G proteins	400		
Activated G proteins transmit signals by binding to other proteins	402		
Cyclic AMP stimulates the phosphorylation of many target proteins by activating protein kinase A	403		
G proteins spontaneously reset themselves through GTP hydrolysis	403		
Some 7TM receptors activate the phosphoinositide cascade	404		
Calcium ion is a widely used second messenger	405		
Calcium ion often activates the regulatory protein calmodulin	407		
<b>14.2 Insulin Signaling: Phosphorylation Cascades Are Central to Many Signal-Transduction Processes</b>	<b>407</b>		
The insulin receptor is a dimer that closes around a bound insulin molecule	408		
Insulin binding results in the cross-phosphorylation and activation of the insulin receptor	408		
The activated insulin-receptor kinase initiates a kinase cascade	409		
Insulin signaling is terminated by the action of phosphatases	411		
<b>14.3 EGF Signaling: Signal-Transduction Pathways Are Poised to Respond</b>	<b>411</b>		
EGF binding results in the dimerization of the EGF receptor	411		
The EGF receptor undergoes phosphorylation of its carboxyl-terminal tail	413		
EGF signaling leads to the activation of Ras, a small G protein	413		
Activated Ras initiates a protein kinase cascade	414		
EGF signaling is terminated by protein phosphatases and the intrinsic GTPase activity of Ras	414		
<b>14.4 Many Elements Recur with Variation in Different Signal-Transduction Pathways</b>	<b>415</b>		
<b>14.5 Defects in Signal-Transduction Pathways Can Lead to Cancer and Other Diseases</b>	<b>416</b>		
Monoclonal antibodies can be used to inhibit signal-transduction pathways activated in tumors	416		
Protein kinase inhibitors can be effective anticancer drugs	417		
Cholera and whooping cough are the result of altered G-protein activity	417		
<b>Part II TRANSDUCING AND STORING ENERGY</b>			
<b>CHAPTER 15 Metabolism: Basic Concepts and Design</b>	<b>423</b>		
<b>15.1 Metabolism Is Composed of Many Coupled, Interconnecting Reactions</b>	<b>424</b>		
Metabolism consists of energy-yielding and energy-requiring reactions	424		
		A thermodynamically unfavorable reaction can be driven by a favorable reaction	425
<b>15.2 ATP Is the Universal Currency of Free Energy in Biological Systems</b>	<b>426</b>		
ATP hydrolysis is exergonic	426		
ATP hydrolysis drives metabolism by shifting the equilibrium of coupled reactions	427		
The high phosphoryl potential of ATP results from structural differences between ATP and its hydrolysis products	429		
Phosphoryl-transfer potential is an important form of cellular energy transformation	430		
<b>15.3 The Oxidation of Carbon Fuels Is an Important Source of Cellular Energy</b>	<b>432</b>		
Compounds with high phosphoryl-transfer potential can couple carbon oxidation to ATP synthesis	432		
Ion gradients across membranes provide an important form of cellular energy that can be coupled to ATP synthesis	433		
Phosphates play a prominent role in biochemical processes	434		
Energy from foodstuffs is extracted in three stages	434		
<b>15.4 Metabolic Pathways Contain Many Recurring Motifs</b>	<b>435</b>		
Activated carriers exemplify the modular design and economy of metabolism	435		
Many activated carriers are derived from vitamins	438		
Key reactions are reiterated throughout metabolism	440		
Metabolic processes are regulated in three principal ways	442		
Aspects of metabolism may have evolved from an RNA world	444		
<b>CHAPTER 16 Glycolysis and Gluconeogenesis</b>	<b>449</b>		
Glucose is generated from dietary carbohydrates	450		
Glucose is an important fuel for most organisms	451		
<b>16.1 Glycolysis Is an Energy-Conversion Pathway in Many Organisms</b>	<b>451</b>		
Hexokinase traps glucose in the cell and begins glycolysis	451		
Fructose 1,6-bisphosphate is generated from glucose 6-phosphate	453		
The six-carbon sugar is cleaved into two three-carbon fragments	454		
Mechanism: Triose phosphate isomerase salvages a three-carbon fragment	455		
The oxidation of an aldehyde to an acid powers the formation of a compound with high phosphoryl-transfer potential	457		
Mechanism: Phosphorylation is coupled to the oxidation of glyceraldehyde 3-phosphate by a thioester intermediate	458		
ATP is formed by phosphoryl transfer from 1,3-bisphosphoglycerate	459		
Additional ATP is generated with the formation of pyruvate	460		
Two ATP molecules are formed in the conversion of glucose into pyruvate	461		

NAD <sup>+</sup> is regenerated from the metabolism of pyruvate	462	Flexible linkages allow lipoamide to move between different active sites	500
Fermentations provide usable energy in the absence of oxygen	464	<b>17.2 The Citric Acid Cycle Oxidizes Two-Carbon Units</b>	<b>501</b>
The binding site for NAD <sup>+</sup> is similar in many dehydrogenases	465	Citrate synthase forms citrate from oxaloacetate and acetyl coenzyme A	502
Fructose is converted into glycolytic intermediates by fructokinase	465	Mechanism: The mechanism of citrate synthase prevents undesirable reactions	502
Excessive fructose consumption can lead to pathological conditions	466	Citrate is isomerized into isocitrate	504
Galactose is converted into glucose 6-phosphate	466	Isocitrate is oxidized and decarboxylated to alpha-ketoglutarate	504
Many adults are intolerant of milk because they are deficient in lactase	467	Succinyl coenzyme A is formed by the oxidative decarboxylation of alpha-ketoglutarate	505
Galactose is highly toxic if the transferase is missing	468	A compound with high phosphoryl-transfer potential is generated from succinyl coenzyme A	505
<b>16.2 The Glycolytic Pathway Is Tightly Controlled</b>	<b>469</b>	Mechanism: Succinyl coenzyme A synthetase transforms types of biochemical energy	506
Glycolysis in muscle is regulated to meet the need for ATP	469	Oxaloacetate is regenerated by the oxidation of succinate	507
The regulation of glycolysis in the liver illustrates the biochemical versatility of the liver	472	The citric acid cycle produces high-transfer-potential electrons, ATP, and CO <sub>2</sub>	508
A family of transporters enables glucose to enter and leave animal cells	473	<b>17.3 Entry to the Citric Acid Cycle and Metabolism Through It Are Controlled</b>	<b>510</b>
Aerobic glycolysis is a property of rapidly growing cells	474	The pyruvate dehydrogenase complex is regulated allosterically and by reversible phosphorylation	511
Cancer and endurance training affect glycolysis in a similar fashion	476	The citric acid cycle is controlled at several points	512
<b>16.3 Glucose Can Be Synthesized from Noncarbohydrate Precursors</b>	<b>476</b>	Defects in the citric acid cycle contribute to the development of cancer	513
Gluconeogenesis is not a reversal of glycolysis	478	<b>17.4 The Citric Acid Cycle Is a Source of Biosynthetic Precursors</b>	<b>514</b>
The conversion of pyruvate into phosphoenolpyruvate begins with the formation of oxaloacetate	478	The citric acid cycle must be capable of being rapidly replenished	514
Oxaloacetate is shuttled into the cytoplasm and converted into phosphoenolpyruvate	480	The disruption of pyruvate metabolism is the cause of beriberi and poisoning by mercury and arsenic	515
The conversion of fructose 1,6-bisphosphate into fructose 6-phosphate and orthophosphate is an irreversible step	480	The citric acid cycle may have evolved from preexisting pathways	516
The generation of free glucose is an important control point	481	<b>17.5 The Glyoxylate Cycle Enables Plants and Bacteria to Grow on Acetate</b>	<b>516</b>
Six high-transfer-potential phosphoryl groups are spent in synthesizing glucose from pyruvate	481	<b>CHAPTER 18 Oxidative Phosphorylation</b>	<b>523</b>
<b>16.4 Gluconeogenesis and Glycolysis Are Reciprocally Regulated</b>	<b>482</b>	<b>18.1 Eukaryotic Oxidative Phosphorylation Takes Place in Mitochondria</b>	<b>524</b>
Energy charge determines whether glycolysis or gluconeogenesis will be most active	482	Mitochondria are bounded by a double membrane	524
The balance between glycolysis and gluconeogenesis in the liver is sensitive to blood-glucose concentration	483	Mitochondria are the result of an endosymbiotic event	525
Substrate cycles amplify metabolic signals and produce heat	485	<b>18.2 Oxidative Phosphorylation Depends on Electron Transfer</b>	<b>526</b>
Lactate and alanine formed by contracting muscle are used by other organs	485	The electron-transfer potential of an electron is measured as redox potential	526
Glycolysis and gluconeogenesis are evolutionarily intertwined	487	A 1.14-volt potential difference between NADH and molecular oxygen drives electron transport through the chain and favors the formation of a proton gradient	528
<b>CHAPTER 17 The Citric Acid Cycle</b>	<b>495</b>	<b>18.3 The Respiratory Chain Consists of Four Complexes: Three Proton Pumps and a Physical Link to the Citric Acid Cycle</b>	<b>529</b>
The citric acid cycle harvests high-energy electrons	496	Iron-sulfur clusters are common components of the electron transport chain	531
<b>17.1 The Pyruvate Dehydrogenase Complex Links Glycolysis to the Citric Acid Cycle</b>	<b>497</b>		
Mechanism: The synthesis of acetyl coenzyme A from pyruvate requires three enzymes and five coenzymes	498		



The high-potential electrons of NADH enter the respiratory chain at NADH-Q oxidoreductase	532	Chloroplasts arose from an endosymbiotic event	568
Ubiquinol is the entry point for electrons from FADH <sub>2</sub> of flavoproteins	533	<b>19.2 Light Absorption by Chlorophyll Induces Electron Transfer</b>	568
Electrons flow from ubiquinol to cytochrome <i>c</i> through Q-cytochrome <i>c</i> oxidoreductase	533	A special pair of chlorophylls initiate charge separation	569
The Q cycle funnels electrons from a two-electron carrier to a one-electron carrier and pumps protons	535	Cyclic electron flow reduces the cytochrome of the reaction center	572
Cytochrome <i>c</i> oxidase catalyzes the reduction of molecular oxygen to water	535	<b>19.3 Two Photosystems Generate a Proton Gradient and NADPH in Oxygenic Photosynthesis</b>	572
Toxic derivatives of molecular oxygen such as superoxide radicals are scavenged by protective enzymes	538	Photosystem II transfers electrons from water to plastoquinone and generates a proton gradient	572
Electrons can be transferred between groups that are not in contact	540	Cytochrome <i>bf</i> links photosystem II to photosystem I	575
The conformation of cytochrome <i>c</i> has remained essentially constant for more than a billion years	541	Photosystem I uses light energy to generate reduced ferredoxin, a powerful reductant	575
<b>18.4 A Proton Gradient Powers the Synthesis of ATP</b>	541	Ferredoxin-NADP <sup>+</sup> reductase converts NADP <sup>+</sup> into NADPH	576
ATP synthase is composed of a proton-conducting unit and a catalytic unit	543	<b>19.4 A Proton Gradient across the Thylakoid Membrane Drives ATP Synthesis</b>	578
Proton flow through ATP synthase leads to the release of tightly bound ATP: The binding-change mechanism	544	The ATP synthase of chloroplasts closely resembles those of mitochondria and prokaryotes	578
Rotational catalysis is the world's smallest molecular motor	546	The activity of chloroplast ATP synthase is regulated	579
Proton flow around the <i>c</i> ring powers ATP synthesis	546	Cyclic electron flow through photosystem I leads to the production of ATP instead of NADPH	580
ATP synthase and G proteins have several common features	548	The absorption of eight photons yields one O <sub>2</sub> , two NADPH, and three ATP molecules	581
<b>18.5 Many Shuttles Allow Movement Across Mitochondrial Membranes</b>	549	<b>19.5 Accessory Pigments Funnel Energy into Reaction Centers</b>	581
Electrons from cytoplasmic NADH enter mitochondria by shuttles	549	Resonance energy transfer allows energy to move from the site of initial absorbance to the reaction center	582
The entry of ADP into mitochondria is coupled to the exit of ATP by ATP-ADP translocase	550	The components of photosynthesis are highly organized	583
Mitochondrial transporters for metabolites have a common tripartite structure	551	Many herbicides inhibit the light reactions of photosynthesis	584
<b>18.6 The Regulation of Cellular Respiration Is Governed Primarily by the Need for ATP</b>	552	<b>19.6 The Ability to Convert Light into Chemical Energy Is Ancient</b>	584
The complete oxidation of glucose yields about 30 molecules of ATP	552	Artificial photosynthetic systems may provide clean, renewable energy	585
The rate of oxidative phosphorylation is determined by the need for ATP	553	<b>CHAPTER 20 The Calvin Cycle and the Pentose Phosphate Pathway</b>	589
ATP synthase can be regulated	554	<b>20.1 The Calvin Cycle Synthesizes Hexoses from Carbon Dioxide and Water</b>	590
Regulated uncoupling leads to the generation of heat	554	Carbon dioxide reacts with ribulose 1,5-bisphosphate to form two molecules of 3-phosphoglycerate	591
Oxidative phosphorylation can be inhibited at many stages	556	Rubisco activity depends on magnesium and carbamate	592
Mitochondrial diseases are being discovered	557	Rubisco activase is essential for rubisco activity	593
Mitochondria play a key role in apoptosis	557	Rubisco also catalyzes a wasteful oxygenase reaction: Catalytic imperfection	593
Power transmission by proton gradients is a central motif of bioenergetics	558	Hexose phosphates are made from phosphoglycerate, and ribulose 1,5-bisphosphate is regenerated	594
<b>CHAPTER 19 The Light Reactions of Photosynthesis</b>	565	Three ATP and two NADPH molecules are used to bring carbon dioxide to the level of a hexose	597
Photosynthesis converts light energy into chemical energy	566	Starch and sucrose are the major carbohydrate stores in plants	597
<b>19.1 Photosynthesis Takes Place in Chloroplasts</b>	567	<b>20.2 The Activity of the Calvin Cycle Depends on Environmental Conditions</b>	598
The primary events of photosynthesis take place in thylakoid membranes	567		

Rubisco is activated by light-driven changes in proton and magnesium ion concentrations	598	Muscle phosphorylase is regulated by the intracellular energy charge	625
Thioredoxin plays a key role in regulating the Calvin cycle	599	Biochemical characteristics of muscle fiber types differ	625
The C <sub>4</sub> pathway of tropical plants accelerates photosynthesis by concentrating carbon dioxide	599	Phosphorylation promotes the conversion of phosphorylase <i>b</i> to phosphorylase <i>a</i>	626
Crassulacean acid metabolism permits growth in arid ecosystems	601	Phosphorylase kinase is activated by phosphorylation and calcium ions	626
<b>20.3 The Pentose Phosphate Pathway Generates NADPH and Synthesizes Five-Carbon Sugars</b>	<b>601</b>	<b>21.3 Epinephrine and Glucagon Signal the Need for Glycogen Breakdown</b>	<b>627</b>
Two molecules of NADPH are generated in the conversion of glucose 6-phosphate into ribulose 5-phosphate	602	G proteins transmit the signal for the initiation of glycogen breakdown	627
The pentose phosphate pathway and glycolysis are linked by <i>transketolase</i> and <i>transaldolase</i>	602	Glycogen breakdown must be rapidly turned off when necessary	629
Mechanism: <i>Transketolase</i> and <i>transaldolase</i> stabilize carbanionic intermediates by different mechanisms	605	The regulation of glycogen phosphorylase became more sophisticated as the enzyme evolved	629
<b>20.4 The Metabolism of Glucose 6-Phosphate by the Pentose Phosphate Pathway Is Coordinated with Glycolysis</b>	<b>607</b>	<b>21.4 Glycogen Is Synthesized and Degraded by Different Pathways</b>	<b>630</b>
The rate of the pentose phosphate pathway is controlled by the level of NADP <sup>+</sup>	607	UDP-glucose is an activated form of glucose	630
The flow of glucose 6-phosphate depends on the need for NADPH, ribose 5-phosphate, and ATP	608	Glycogen synthase catalyzes the transfer of glucose from UDP-glucose to a growing chain	630
The pentose phosphate pathway is required for rapid cell growth	610	A branching enzyme forms $\alpha$ -1,6 linkages	631
Through the looking-glass: The Calvin cycle and the pentose phosphate pathway are mirror images	610	Glycogen synthase is the key regulatory enzyme in glycogen synthesis	632
<b>20.5 Glucose 6-Phosphate Dehydrogenase Plays a Key Role in Protection Against Reactive Oxygen Species</b>	<b>610</b>	Glycogen is an efficient storage form of glucose	632
Glucose 6-phosphate dehydrogenase deficiency causes a drug-induced hemolytic anemia	610	<b>21.5 Glycogen Breakdown and Synthesis Are Reciprocally Regulated</b>	<b>632</b>
A deficiency of glucose 6-phosphate dehydrogenase confers an evolutionary advantage in some circumstances	612	Protein phosphatase 1 reverses the regulatory effects of kinases on glycogen metabolism	633
<b>CHAPTER 21 Glycogen Metabolism</b>	<b>617</b>	Insulin stimulates glycogen synthesis by inactivating glycogen synthase kinase	635
Glycogen metabolism is the regulated release and storage of glucose	618	Glycogen metabolism in the liver regulates the blood-glucose level	635
<b>21.1 Glycogen Breakdown Requires the Interplay of Several Enzymes</b>	<b>619</b>	A biochemical understanding of glycogen-storage diseases is possible	637
Phosphorylase catalyzes the phosphorolytic cleavage of glycogen to release glucose 1-phosphate	619	<b>CHAPTER 22 Fatty Acid Metabolism</b>	<b>643</b>
Mechanism: Pyridoxal phosphate participates in the phosphorolytic cleavage of glycogen	620	Fatty acid degradation and synthesis mirror each other in their chemical reactions	644
A debranching enzyme also is needed for the breakdown of glycogen	621	<b>22.1 Triacylglycerols Are Highly Concentrated Energy Stores</b>	<b>645</b>
Phosphoglucomutase converts glucose 1-phosphate into glucose 6-phosphate	622	Dietary lipids are digested by pancreatic lipases	645
The liver contains glucose 6-phosphatase, a hydrolytic enzyme absent from muscle	622	Dietary lipids are transported in chylomicrons	646
<b>21.2 Phosphorylase Is Regulated by Allosteric Interactions and Reversible Phosphorylation</b>	<b>623</b>	<b>22.2 The Use of Fatty Acids as Fuel Requires Three Stages of Processing</b>	<b>647</b>
Liver phosphorylase produces glucose for use by other tissues	623	Triacylglycerols are hydrolyzed by hormone-stimulated lipases	647
		Free fatty acids and glycerol are released into the blood	648
		Fatty acids are linked to coenzyme A before they are oxidized	648
		Carnitine carries long-chain activated fatty acids into the mitochondrial matrix	649
		Acetyl CoA, NADH, and FADH <sub>2</sub> are generated in each round of fatty acid oxidation	650

The complete oxidation of palmitate yields 106 molecules of ATP	652		
<b>22.3 Unsaturated and Odd-Chain Fatty Acids Require Additional Steps for Degradation</b>	652		
An isomerase and a reductase are required for the oxidation of unsaturated fatty acids	652		
Odd-chain fatty acids yield propionyl CoA in the final thiolysis step	654		
Vitamin B <sub>12</sub> contains a corrin ring and a cobalt atom	654		
Mechanism: Methylmalonyl CoA mutase catalyzes a rearrangement to form succinyl CoA	655		
Fatty acids are also oxidized in peroxisomes	656		
Ketone bodies are formed from acetyl CoA when fat breakdown predominates	657		
Ketone bodies are a major fuel in some tissues	658		
Animals cannot convert fatty acids into glucose	660		
Some fatty acids may contribute to the development of pathological conditions	661		
<b>22.4 Fatty Acids Are Synthesized by Fatty Acid Synthase</b>	661		
Fatty acids are synthesized and degraded by different pathways	661		
The formation of malonyl CoA is the committed step in fatty acid synthesis	662		
Intermediates in fatty acid synthesis are attached to an acyl carrier protein	662		
Fatty acid synthesis consists of a series of condensation, reduction, dehydration, and reduction reactions	662		
Fatty acids are synthesized by a multifunctional enzyme complex in animals	664		
The synthesis of palmitate requires 8 molecules of acetyl CoA, 14 molecules of NADPH, and 7 molecules of ATP	666		
Citrate carries acetyl groups from mitochondria to the cytoplasm for fatty acid synthesis	666		
Several sources supply NADPH for fatty acid synthesis	667		
Fatty acid metabolism is altered in tumor cells	667		
<b>22.5 The Elongation and Unsaturation of Fatty Acids are Accomplished by Accessory Enzyme Systems</b>	668		
Membrane-bound enzymes generate unsaturated fatty acids	668		
Eicosanoid hormones are derived from polyunsaturated fatty acids	669		
Variations on a theme: Polyketide and nonribosomal peptide synthetases resemble fatty acid synthase	670		
<b>22.6 Acetyl CoA Carboxylase Plays a Key Role in Controlling Fatty Acid Metabolism</b>	670		
Acetyl CoA carboxylase is regulated by conditions in the cell	671		
Acetyl CoA carboxylase is regulated by a variety of hormones	671		
<b>CHAPTER 23 Protein Turnover and Amino Acid Catabolism</b>	<b>681</b>		
<b>23.1 Proteins are Degraded to Amino Acids</b>	682		
The digestion of dietary proteins begins in the stomach and is completed in the intestine	682		
Cellular proteins are degraded at different rates	682		
<b>23.2 Protein Turnover Is Tightly Regulated</b>	683		
Ubiquitin tags proteins for destruction	683		
The proteasome digests the ubiquitin-tagged proteins	685		
The ubiquitin pathway and the proteasome have prokaryotic counterparts	686		
Protein degradation can be used to regulate biological function	687		
<b>23.3 The First Step in Amino Acid Degradation Is the Removal of Nitrogen</b>	687		
Alpha-amino groups are converted into ammonium ions by the oxidative deamination of glutamate	687		
Mechanism: Pyridoxal phosphate forms Schiff-base intermediates in aminotransferases	689		
Aspartate aminotransferase is an archetypal pyridoxal-dependent transaminase	690		
Blood levels of aminotransferases serve a diagnostic function	691		
Pyridoxal phosphate enzymes catalyze a wide array of reactions	691		
Serine and threonine can be directly deaminated	692		
Peripheral tissues transport nitrogen to the liver	692		
<b>23.4 Ammonium Ion Is Converted into Urea in Most Terrestrial Vertebrates</b>	693		
The urea cycle begins with the formation of carbamoyl phosphate	693		
Carbamoyl phosphate synthetase is the key regulatory enzyme for urea synthesis	694		
Carbamoyl phosphate reacts with ornithine to begin the urea cycle	694		
The urea cycle is linked to gluconeogenesis	696		
Urea-cycle enzymes are evolutionarily related to enzymes in other metabolic pathways	696		
Inherited defects of the urea cycle cause hyperammonemia and can lead to brain damage	697		
Urea is not the only means of disposing of excess nitrogen	698		
<b>23.5 Carbon Atoms of Degraded Amino Acids Emerge as Major Metabolic Intermediates</b>	698		
Pyruvate is an entry point into metabolism for a number of amino acids	699		
Oxaloacetate is an entry point into metabolism for aspartate and asparagine	700		
Alpha-ketoglutarate is an entry point into metabolism for five-carbon amino acids	700		
Succinyl coenzyme A is a point of entry for several nonpolar amino acids	701		
Methionine degradation requires the formation of a key methyl donor, S-adenosylmethionine	701		
The branched-chain amino acids yield acetyl CoA, acetoacetate, or propionyl CoA	701		
Oxygenases are required for the degradation of aromatic amino acids	703		
<b>23.6 Inborn Errors of Metabolism Can Disrupt Amino Acid Degradation</b>	705		
Phenylketonuria is one of the most common metabolic disorders	706		
Determining the basis of the neurological symptoms of phenylketonuria is an active area of research	706		

**Part III SYNTHESIZING THE MOLECULES OF LIFE**
**CHAPTER 24 The Biosynthesis of Amino Acids 713**

Amino acid synthesis requires solutions to three key biochemical problems	714
<b>24.1 Nitrogen Fixation: Microorganisms Use ATP and a Powerful Reductant to Reduce Atmospheric Nitrogen to Ammonia</b>	<b>714</b>
The iron–molybdenum cofactor of nitrogenase binds and reduces atmospheric nitrogen	715
Ammonium ion is assimilated into an amino acid through glutamate and glutamine	717
<b>24.2 Amino Acids Are Made from Intermediates of the Citric Acid Cycle and Other Major Pathways</b>	<b>719</b>
Human beings can synthesize some amino acids but must obtain others from their diet	719
Aspartate, alanine, and glutamate are formed by the addition of an amino group to an alpha-ketoacid	720
A common step determines the chirality of all amino acids	721
The formation of asparagine from aspartate requires an adenylated intermediate	721
Glutamate is the precursor of glutamine, proline, and arginine	722
3-Phosphoglycerate is the precursor of serine, cysteine, and glycine	722
Tetrahydrofolate carries activated one-carbon units at several oxidation levels	723
S-Adenosylmethionine is the major donor of methyl groups	724
Cysteine is synthesized from serine and homocysteine	726
High homocysteine levels correlate with vascular disease	726
Shikimate and chorismate are intermediates in the biosynthesis of aromatic amino acids	727
Tryptophan synthase illustrates substrate channeling in enzymatic catalysis	729
<b>24.3 Feedback Inhibition Regulates Amino Acid Biosynthesis</b>	<b>730</b>
Branched pathways require sophisticated regulation	731
The sensitivity of glutamine synthetase to allosteric regulation is altered by covalent modification	732
<b>24.4 Amino Acids Are Precursors of Many Biomolecules</b>	<b>734</b>
Glutathione, a gamma-glutamyl peptide, serves as a sulfhydryl buffer and an antioxidant	734
Nitric oxide, a short-lived signal molecule, is formed from arginine	735
Porphyrins are synthesized from glycine and succinyl coenzyme A	736
Porphyrins accumulate in some inherited disorders of porphyrin metabolism	737

**CHAPTER 25 Nucleotide Biosynthesis 743**

Nucleotides can be synthesized by de novo or salvage pathways	744
---	-----

<b>25.1 The Pyrimidine Ring Is Assembled de Novo or Recovered by Salvage Pathways</b>	<b>744</b>
Bicarbonate and other oxygenated carbon compounds are activated by phosphorylation	745
The side chain of glutamine can be hydrolyzed to generate ammonia	745
Intermediates can move between active sites by channeling	745
Orotate acquires a ribose ring from PRPP to form a pyrimidine nucleotide and is converted into uridylylate	746
Nucleotide mono-, di-, and triphosphates are interconvertible	747
CTP is formed by amination of UTP	747
Salvage pathways recycle pyrimidine bases	748
<b>25.2 Purine Bases Can Be Synthesized de Novo or Recycled by Salvage Pathways</b>	<b>748</b>
The purine ring system is assembled on ribose phosphate	749
The purine ring is assembled by successive steps of activation by phosphorylation followed by displacement	749
AMP and GMP are formed from IMP	751
Enzymes of the purine synthesis pathway associate with one another in vivo	752
Salvage pathways economize intracellular energy expenditure	752
<b>25.3 Deoxyribonucleotides Are Synthesized by the Reduction of Ribonucleotides Through a Radical Mechanism</b>	<b>753</b>
Mechanism: A tyrosyl radical is critical to the action of ribonucleotide reductase	753
Stable radicals other than tyrosyl radical are employed by other ribonucleotide reductases	755
Thymidylate is formed by the methylation of deoxyuridylylate	755
Dihydrofolate reductase catalyzes the regeneration of tetrahydrofolate, a one-carbon carrier	756
Several valuable anticancer drugs block the synthesis of thymidylate	757
<b>25.4 Key Steps in Nucleotide Biosynthesis Are Regulated by Feedback Inhibition</b>	<b>758</b>
Pyrimidine biosynthesis is regulated by aspartate transcarbamoylase	758
The synthesis of purine nucleotides is controlled by feedback inhibition at several sites	758
The synthesis of deoxyribonucleotides is controlled by the regulation of ribonucleotide reductase	759
<b>25.5 Disruptions in Nucleotide Metabolism Can Cause Pathological Conditions</b>	<b>760</b>
The loss of adenosine deaminase activity results in severe combined immunodeficiency	760
Gout is induced by high serum levels of urate	761
Lesch–Nyhan syndrome is a dramatic consequence of mutations in a salvage-pathway enzyme	761
Folic acid deficiency promotes birth defects such as spina bifida	762

---

**CHAPTER 26 The Biosynthesis of Membrane Lipids and Steroids** 767
 

---

**26.1 Phosphatidate Is a Common Intermediate in the Synthesis of Phospholipids and Triacylglycerols** 768

- The synthesis of phospholipids requires an activated intermediate 769
- Some phospholipids are synthesized from an activated alcohol 770
- Phosphatidylcholine is an abundant phospholipid 770
- Excess choline is implicated in the development of heart disease 771
- Base-exchange reactions can generate phospholipids 771
- Sphingolipids are synthesized from ceramide 772
- Gangliosides are carbohydrate-rich sphingolipids that contain acidic sugars 772
- Sphingolipids confer diversity on lipid structure and function 773
- Respiratory distress syndrome and Tay-Sachs disease result from the disruption of lipid metabolism 774
- Ceramide metabolism stimulates tumor growth 774
- Phosphatidic acid phosphatase is a key regulatory enzyme in lipid metabolism 775

**26.2 Cholesterol Is Synthesized from Acetyl Coenzyme A in Three Stages** 776

- The synthesis of mevalonate, which is activated as isopentenyl pyrophosphate, initiates the synthesis of cholesterol 776
- Squalene (C<sub>30</sub>) is synthesized from six molecules of isopentenyl pyrophosphate (C<sub>5</sub>) 777
- Squalene cyclizes to form cholesterol 778

**26.3 The Complex Regulation of Cholesterol Biosynthesis Takes Place at Several Levels** 779

- Lipoproteins transport cholesterol and triacylglycerols throughout the organism 782
- Low-density lipoproteins play a central role in cholesterol metabolism 784
- The absence of the LDL receptor leads to hypercholesterolemia and atherosclerosis 784
- Mutations in the LDL receptor prevent LDL release and result in receptor destruction 785
- Cycling of the LDL receptor is regulated 787
- HDL appears to protect against atherosclerosis 787
- The clinical management of cholesterol levels can be understood at a biochemical level 788

**26.4 Important Derivatives of Cholesterol Include Bile Salts and Steroid Hormones** 788

- Letters identify the steroid rings and numbers identify the carbon atoms 790
- Steroids are hydroxylated by cytochrome P450 monooxygenases that use NADPH and O<sub>2</sub> 790
- The cytochrome P450 system is widespread and performs a protective function 791
- Pregnenolone, a precursor of many other steroids, is formed from cholesterol by cleavage of its side chain 792

- Pregesterone and corticosteroids are synthesized from pregnenolone 792
- Androgens and estrogens are synthesized from pregnenolone 792
- Vitamin D is derived from cholesterol by the ring-splitting activity of light 794

---

**CHAPTER 27 The Integration of Metabolism** 801
 

---

**27.1 Caloric Homeostasis Is a Means of Regulating Body Weight** 802

**27.2 The Brain Plays a Key Role in Caloric Homeostasis** 804

- Signals from the gastrointestinal tract induce feelings of satiety 804
- Leptin and insulin regulate long-term control over caloric homeostasis 805
- Leptin is one of several hormones secreted by adipose tissue 806
- Leptin resistance may be a contributing factor to obesity 806
- Dieting is used to combat obesity 807

**27.3 Diabetes Is a Common Metabolic Disease Often Resulting from Obesity** 807

- Insulin initiates a complex signal-transduction pathway in muscle 808
- Metabolic syndrome often precedes type 2 diabetes 809
- Excess fatty acids in muscle modify metabolism 810
- Insulin resistance in muscle facilitates pancreatic failure 810
- Metabolic derangements in type 1 diabetes result from insulin insufficiency and glucagon excess 812

**27.4 Exercise Beneficially Alters the Biochemistry of Cells** 813

- Mitochondrial biogenesis is stimulated by muscular activity 813
- Fuel choice during exercise is determined by the intensity and duration of activity 813

**27.5 Food Intake and Starvation Induce Metabolic Changes** 816

- The starved-fed cycle is the physiological response to a fast 816
- Metabolic adaptations in prolonged starvation minimize protein degradation 818

**27.6 Ethanol Alters Energy Metabolism in the Liver** 819

- Ethanol metabolism leads to an excess of NADH 820
- Excess ethanol consumption disrupts vitamin metabolism 821

---

**CHAPTER 28 DNA Replication, Repair, and Recombination** 827
 

---

**28.1 DNA Replication Proceeds by the Polymerization of Deoxyribonucleoside Triphosphates Along a Template** 828

DNA polymerases require a template and a primer	829	<b>CHAPTER 29 RNA Synthesis and Processing</b>	<b>859</b>
All DNA polymerases have structural features in common	829	RNA synthesis comprises three stages: Initiation, elongation, and termination	860
Two bound metal ions participate in the polymerase reaction	829	<b>29.1 RNA Polymerases Catalyze Transcription</b>	<b>861</b>
The specificity of replication is dictated by complementarity of shape between bases	830	RNA chains are formed <i>de novo</i> and grow in the 5'-to-3' direction	862
An RNA primer synthesized by primase enables DNA synthesis to begin	831	RNA polymerases backtrack and correct errors	863
One strand of DNA is made continuously, whereas the other strand is synthesized in fragments	831	RNA polymerase binds to promoter sites on the DNA template to initiate transcription	864
DNA ligase joins ends of DNA in duplex regions	832	Sigma subunits of RNA polymerase recognize promoter sites	865
The separation of DNA strands requires specific helicases and ATP hydrolysis	832	RNA polymerases must unwind the template double helix for transcription to take place	865
<b>28.2 DNA Unwinding and Supercoiling Are Controlled by Topoisomerases</b>	<b>833</b>	Elongation takes place at transcription bubbles that move along the DNA template	866
The linking number of DNA, a topological property, determines the degree of supercoiling	835	Sequences within the newly transcribed RNA signal termination	866
Topoisomerases prepare the double helix for unwinding	836	Some messenger RNAs directly sense metabolite concentrations	867
Type I topoisomerases relax supercoiled structures	836	The <i>rho</i> protein helps to terminate the transcription of some genes	868
Type II topoisomerases can introduce negative supercoils through coupling to ATP hydrolysis	837	Some antibiotics inhibit transcription	869
<b>28.3 DNA Replication Is Highly Coordinated</b>	<b>839</b>	Precursors of transfer and ribosomal RNA are cleaved and chemically modified after transcription in prokaryotes	870
DNA replication requires highly processive polymerases	839	<b>29.2 Transcription in Eukaryotes Is Highly Regulated</b>	<b>871</b>
The leading and lagging strands are synthesized in a coordinated fashion	840	Three types of RNA polymerase synthesize RNA in eukaryotic cells	872
DNA replication in <i>Escherichia coli</i> begins at a unique site	842	Three common elements can be found in the RNA polymerase II promoter region	874
DNA synthesis in eukaryotes is initiated at multiple sites	843	The TFIID protein complex initiates the assembly of the active transcription complex	874
Telomeres are unique structures at the ends of linear chromosomes	844	Multiple transcription factors interact with eukaryotic promoters	875
Telomeres are replicated by telomerase, a specialized polymerase that carries its own RNA template	845	Enhancer sequences can stimulate transcription at start sites thousands of bases away	876
<b>28.4 Many Types of DNA Damage Can Be Repaired</b>	<b>845</b>	<b>29.3 The Transcription Products of Eukaryotic Polymerases Are Processed</b>	<b>876</b>
Errors can arise in DNA replication	846	RNA polymerase I produces three ribosomal RNAs	877
Bases can be damaged by oxidizing agents, alkylating agents, and light	846	RNA polymerase III produces transfer RNA	877
DNA damage can be detected and repaired by a variety of systems	847	The product of RNA polymerase II, the pre-mRNA transcript, acquires a 5' cap and a 3' poly(A) tail	878
The presence of thymine instead of uracil in DNA permits the repair of deaminated cytosine	849	Small regulatory RNAs are cleaved from larger precursors	879
Some genetic diseases are caused by the expansion of repeats of three nucleotides	850	RNA editing changes the proteins encoded by mRNA	879
Many cancers are caused by the defective repair of DNA	850	Sequences at the ends of introns specify splice sites in mRNA precursors	880
Many potential carcinogens can be detected by their mutagenic action on bacteria	852	Splicing consists of two sequential transesterification reactions	881
<b>28.5 DNA Recombination Plays Important Roles in Replication, Repair, and Other Processes</b>	<b>852</b>	Small nuclear RNAs in spliceosomes catalyze the splicing of mRNA precursors	882
RecA can initiate recombination by promoting strand invasion	853	Transcription and processing of mRNA are coupled	883
Some recombination reactions proceed through Holliday-junction intermediates	854	Mutations that affect pre-mRNA splicing cause disease	884
		Most human pre-mRNAs can be spliced in alternative ways to yield different proteins	885
		<b>29.4 The Discovery of Catalytic RNA was Revealing in Regard to Both Mechanism and Evolution</b>	<b>886</b>

<b>CHAPTER 30 Protein Synthesis</b>	<b>893</b>	Transport vesicles carry cargo proteins to their final destination	918
<b>30.1 Protein Synthesis Requires the Translation of Nucleotide Sequences into Amino Acid Sequences</b>	<b>894</b>	<b>CHAPTER 31 The Control of Gene Expression in Prokaryotes</b>	<b>925</b>
The synthesis of long proteins requires a low error frequency	894	<b>31.1 Many DNA-Binding Proteins Recognize Specific DNA Sequences</b>	<b>926</b>
Transfer RNA molecules have a common design	895	The helix-turn-helix motif is common to many prokaryotic DNA-binding proteins	927
Some transfer RNA molecules recognize more than one codon because of wobble in base-pairing	897	<b>31.2 Prokaryotic DNA-Binding Proteins Bind Specifically to Regulatory Sites in Operons</b>	<b>927</b>
<b>30.2 Aminoacyl Transfer RNA Synthetases Read the Genetic Code</b>	<b>898</b>	An operon consists of regulatory elements and protein-encoding genes	928
Amino acids are first activated by adenylation	898	The <i>lac</i> repressor protein in the absence of lactose binds to the operator and blocks transcription	929
Aminoacyl-tRNA synthetases have highly discriminating amino acid activation sites	899	Ligand binding can induce structural changes in regulatory proteins	930
Proofreading by aminoacyl-tRNA synthetases increases the fidelity of protein synthesis	900	The operon is a common regulatory unit in prokaryotes	930
Synthetases recognize various features of transfer RNA molecules	901	Transcription can be stimulated by proteins that contact RNA polymerase	931
Aminoacyl-tRNA synthetases can be divided into two classes	901	<b>31.3 Regulatory Circuits Can Result in Switching Between Patterns of Gene Expression</b>	<b>932</b>
<b>30.3 The Ribosome Is the Site of Protein Synthesis</b>	<b>902</b>	The $\lambda$ repressor regulates its own expression	932
Ribosomal RNAs (5S, 16S, and 23S rRNA) play a central role in protein synthesis	903	A circuit based on the $\lambda$ repressor and Cro forms a genetic switch	933
Ribosomes have three tRNA-binding sites that bridge the 30s and 50s subunits	905	Many prokaryotic cells release chemical signals that regulate gene expression in other cells	933
The start signal is usually AUG preceded by several bases that pair with 16S rRNA	905	Biofilms are complex communities of prokaryotes	934
Bacterial protein synthesis is initiated by formylmethionyl transfer RNA	906	<b>31.4 Gene Expression Can Be Controlled at Posttranscriptional Levels</b>	<b>935</b>
Formylmethionyl-tRNA <sub>f</sub> is placed in the P site of the ribosome in the formation of the 70S initiation complex	907	Attenuation is a prokaryotic mechanism for regulating transcription through the modulation of nascent RNA secondary structure	935
Elongation factors deliver aminoacyl-tRNA to the ribosome	907	<b>CHAPTER 32 The Control of Gene Expression in Eukaryotes</b>	<b>941</b>
Peptidyl transferase catalyzes peptide-bond synthesis	908	<b>32.1 Eukaryotic DNA Is Organized into Chromatin</b>	<b>943</b>
The formation of a peptide bond is followed by the GTP-driven translocation of tRNAs and mRNA	909	Nucleosomes are complexes of DNA and histones	943
Protein synthesis is terminated by release factors that read stop codons	910	DNA wraps around histone octamers to form nucleosomes	943
<b>30.4 Eukaryotic Protein Synthesis Differs from Bacterial Protein Synthesis Primarily in Translation Initiation</b>	<b>911</b>	<b>32.2 Transcription Factors Bind DNA and Regulate Transcription Initiation</b>	<b>945</b>
Mutations in initiation factor 2 cause a curious pathological condition	913	A range of DNA-binding structures are employed by eukaryotic DNA-binding proteins	945
<b>30.5 A Variety of Antibiotics and Toxins Can Inhibit Protein Synthesis</b>	<b>913</b>	Activation domains interact with other proteins	946
Some antibiotics inhibit protein synthesis	914	Multiple transcription factors interact with eukaryotic regulatory regions	946
Diphtheria toxin blocks protein synthesis in eukaryotes by inhibiting translocation	914	Enhancers can stimulate transcription in specific cell types	946
Ricin fatally modifies 28S ribosomal RNA	915	Induced pluripotent stem cells can be generated by introducing four transcription factors into differentiated cells	947
<b>30.6 Ribosomes Bound to the Endoplasmic Reticulum Manufacture Secretory and Membrane Proteins</b>	<b>915</b>	<b>32.3 The Control of Gene Expression Can Require Chromatin Remodeling</b>	<b>948</b>
Protein synthesis begins on ribosomes that are free in the cytoplasm	916		
Signal sequences mark proteins for translocation across the endoplasmic reticulum membrane	916		

The methylation of DNA can alter patterns of gene expression	949	Mechanosensory channels have been identified in <i>Drosophila</i> and vertebrates	976
Steroids and related hydrophobic molecules pass through membranes and bind to DNA-binding receptors	949	<b>33.5 Touch Includes the Sensing of Pressure, Temperature, and Other Factors</b>	<b>977</b>
Nuclear hormone receptors regulate transcription by recruiting coactivators to the transcription complex	950	Studies of capsaicin reveal a receptor for sensing high temperatures and other painful stimuli	977
Steroid-hormone receptors are targets for drugs	951	<b>CHAPTER 34 The Immune System</b>	<b>981</b>
Chromatin structure is modulated through covalent modifications of histone tails	952	Innate immunity is an evolutionarily ancient defense system	982
Histone deacetylases contribute to transcriptional repression	953	The adaptive immune system responds by using the principles of evolution	984
<b>32.4 Eukaryotic Gene Expression Can Be Controlled at Posttranscriptional Levels</b>	<b>954</b>	<b>34.1 Antibodies Possess Distinct Antigen-Binding and Effector Units</b>	<b>985</b>
Genes associated with iron metabolism are translationally regulated in animals	954	<b>34.2 Antibodies Bind Specific Molecules Through Hypervariable Loops</b>	<b>988</b>
Small RNAs regulate the expression of many eukaryotic genes	956	The immunoglobulin fold consists of a beta-sandwich framework with hypervariable loops	988
<b>Part IV RESPONDING TO ENVIRONMENTAL CHANGES</b>		X-ray analyses have revealed how antibodies bind antigens	989
<b>CHAPTER 33 Sensory Systems</b>	<b>961</b>	Large antigens bind antibodies with numerous interactions	990
<b>33.1 A Wide Variety of Organic Compounds Are Detected by Olfaction</b>	<b>962</b>	<b>34.3 Diversity Is Generated by Gene Rearrangements</b>	<b>991</b>
Olfaction is mediated by an enormous family of seven-transmembrane-helix receptors	962	<i>J</i> ( <i>joining</i> ) genes and <i>D</i> ( <i>diversity</i> ) genes increase antibody diversity	991
Odorants are decoded by a combinatorial mechanism	964	More than 10 <sup>8</sup> antibodies can be formed by combinatorial association and somatic mutation	992
<b>33.2 Taste Is a Combination of Senses That Function by Different Mechanisms</b>	<b>966</b>	The oligomerization of antibodies expressed on the surfaces of immature B cells triggers antibody secretion	993
Sequencing of the human genome led to the discovery of a large family of 7TM bitter receptors	967	Different classes of antibodies are formed by the hopping of V <sub>H</sub> genes	994
A heterodimeric 7TM receptor responds to sweet compounds	968	<b>34.4 Major-Histocompatibility-Complex Proteins Present Peptide Antigens on Cell Surfaces for Recognition by T-Cell Receptors</b>	<b>995</b>
Umami, the taste of glutamate and aspartate, is mediated by a heterodimeric receptor related to the sweet receptor	969	Peptides presented by MHC proteins occupy a deep groove flanked by alpha helices	996
Salty tastes are detected primarily by the passage of sodium ions through channels	969	T-cell receptors are antibody-like proteins containing variable and constant regions	998
Sour tastes arise from the effects of hydrogen ions (acids) on channels	969	CD8 on cytotoxic T cells acts in concert with T-cell receptors	998
<b>33.3 Photoreceptor Molecules in the Eye Detect Visible Light</b>	<b>970</b>	Helper T cells stimulate cells that display foreign peptides bound to class II MHC proteins	1000
Rhodopsin, a specialized 7TM receptor, absorbs visible light	970	Helper T cells rely on the T-cell receptor and CD4 to recognize foreign peptides on antigen-presenting cells	1000
Light absorption induces a specific isomerization of bound 11- <i>cis</i> -retinal	971	MHC proteins are highly diverse	1002
Light-induced lowering of the calcium level coordinates recovery	972	Human immunodeficiency viruses subvert the immune system by destroying helper T cells	1003
Color vision is mediated by three cone receptors that are homologs of rhodopsin	973	<b>34.5 The Immune System Contributes to the Prevention and the Development of Human Diseases</b>	<b>1004</b>
Rearrangements in the genes for the green and red pigments lead to "color blindness"	974	T cells are subjected to positive and negative selection in the thymus	1004
<b>33.4 Hearing Depends on the Speedy Detection of Mechanical Stimuli</b>	<b>975</b>	Autoimmune diseases result from the generation of immune responses against self-antigens	1005
Hair cells use a connected bundle of stereocilia to detect tiny motions	975		



The immune system plays a role in cancer prevention	1005	Drugs must have suitable properties to reach their targets	1036
Vaccines are a powerful means to prevent and eradicate disease	1006	Toxicity can limit drug effectiveness	1040
<b>CHAPTER 35 Molecular Motors</b>	<b>1011</b>	<b>36.2 Drug Candidates Can Be Discovered by Serendipity, Screening, or Design</b>	<b>1041</b>
<b>35.1 Most Molecular-Motor Proteins Are Members of the P-Loop NTPase Superfamily</b>	<b>1012</b>	Serendipitous observations can drive drug development	1041
Molecular motors are generally oligomeric proteins with an ATPase core and an extended structure	1012	Natural products are a valuable source of drugs and drug leads	1043
ATP binding and hydrolysis induce changes in the conformation and binding affinity of motor proteins	1014	Screening libraries of synthetic compounds expands the opportunity for identification of drug leads	1044
<b>35.2 Myosins Move Along Actin Filaments</b>	<b>1016</b>	Drugs can be designed on the basis of three-dimensional structural information about their targets	1046
Actin is a polar, self-assembling, dynamic polymer	1016	<b>36.3 Analyses of Genomes Hold Great Promise for Drug Discovery</b>	<b>1048</b>
Myosin head domains bind to actin filaments	1018	Potential targets can be identified in the human proteome	1048
Motions of single motor proteins can be directly observed	1018	Animal models can be developed to test the validity of potential drug targets	1049
Phosphate release triggers the myosin power stroke	1019	Potential targets can be identified in the genomes of pathogens	1050
Muscle is a complex of myosin and actin	1019	Genetic differences influence individual responses to drugs	1050
The length of the lever arm determines motor velocity	1022	<b>36.4 The Clinical Development of Drugs Proceeds Through Several Phases</b>	<b>1051</b>
<b>35.3 Kinesin and Dynein Move Along Microtubules</b>	<b>1022</b>	Clinical trials are time consuming and expensive	1052
Microtubules are hollow cylindrical polymers	1022	The evolution of drug resistance can limit the utility of drugs for infectious agents and cancer	1053
Kinesin motion is highly processive	1024	<b>Answers to Problems</b>	<b>A1</b>
<b>35.4 A Rotary Motor Drives Bacterial Motion</b>	<b>1026</b>	<b>Selected Readings</b>	<b>B1</b>
Bacteria swim by rotating their flagella	1026	<b>Index</b>	<b>C1</b>
Proton flow drives bacterial flagellar rotation	1026		
Bacterial chemotaxis depends on reversal of the direction of flagellar rotation	1028		
<b>CHAPTER 36 Drug Development</b>	<b>1033</b>		
<b>36.1 The Development of Drugs Presents Huge Challenges</b>	<b>1034</b>		
Drug candidates must be potent and selective modulators of their targets	1035		