

# FRUITS OF ANGIOSPERMS

by

Prof. Dr. INGRID ROTH

Universidad Central de Venezuela, Caracas

With 232 figures



1977

---

GEBRÜDER BORNTRÄEGER · BERLIN · STUTTGART

## Contents

<b>A. General part . . . . .</b>	<b>1</b>
1 The concept "fruit" . . . . .	1
1.1 Definition of the fruit . . . . .	1
1.2 Classification of fruits . . . . .	1
2 The carpel . . . . .	6
3 Fruit symmetry . . . . .	8
4 The pericarp . . . . .	9
4.1 Definition of exo-, meso-, endocarp . . . . .	9
5 Relation between anatomical structure and functions of the pericarp . . . . .	11
6 Fruit growth . . . . .	12
6.1 Growth stimulation . . . . .	13
6.2 Cell division . . . . .	14
6.3 Cell enlargement . . . . .	15
6.4 Periods of growth . . . . .	16
6.5 Cell size in relation to fruit size . . . . .	19
7 Meristems and fruit growth . . . . .	21
8 Cell and tissue differentiation . . . . .	26
9 Cell contents . . . . .	27
10 Maturation . . . . .	30
11 Senescence . . . . .	33
12 Texture and edibility of fruits . . . . .	34
13 Pattern formation . . . . .	37
Outer epidermis 37, Inner epidermis 37, Interwoven pattern 38, Formation of radial cell rows 38, Starlike arrangement of cells 38, Giant cells 38	
 <b>B. Tissues composing the pericarp . . . . .</b>	<b>39</b>
14 Epidermis . . . . .	39
14.1 Outer epidermis . . . . .	39
Shape of epidermal cells . . . . .	41
Cell wall structure . . . . .	41
Cutine and wax formations . . . . .	42
Cell contents . . . . .	42
Multiseriate outer epidermis . . . . .	42

14.2	Inner epidermis . . . . .	44
	Multiseriate inner epidermis . . . . .	47
	Differentiation of the endocarp . . . . .	48
14.3	Stomata . . . . .	49
	Occurrence of stomata . . . . .	51
	Stomata distribution and density . . . . .	52
	Stomata development . . . . .	54
	Anomalous stoma formation and reduction of stomata . . . . .	54
	Function of stomata . . . . .	57
14.4	Trichomes . . . . .	57
	Shape of trichomes (unicellular hairs, pluricellular hairs) . . . . .	57
	Internal or locular hairs . . . . .	60
	Development of hairs . . . . .	62
	Density of hairs . . . . .	62
	Systematic value of hairs . . . . .	63
	Function of hairs . . . . .	63
14.5	Emergences . . . . .	65
15	Periderm and lenticel formation . . . . .	66
16	Hypodermis . . . . .	69
17	Parenchyma . . . . .	70
17.1	Special cell arrangements and cell patterns . . . . .	71
17.2	Aerenchyma . . . . .	73
17.3	Air containing parenchyma . . . . .	73
17.4	Cell contents . . . . .	74
18	Collenchyma . . . . .	74
19	Sclerenchyma . . . . .	75
19.1	Sclereids . . . . .	77
	Cell shape . . . . .	77
	Distribution of sclereids . . . . .	79
	Special stone cell patterns in Solanaceae . . . . .	80
	Taxonomic and phylogenetic value of stone cell concretions . . . . .	80
	Transitional forms between sclereids and fibers or parenchyma cells . . . . .	82
19.2	Fibers . . . . .	82
	Shape of fibers . . . . .	83
	Distribution and arrangement of fibers . . . . .	83
	Origin and differentiation of stone cells and fibers . . . . .	86
20	Vascular system . . . . .	87
21	Secretory structures . . . . .	93
21.1	External secretory structures . . . . .	93
	Nectaries hydathodes, glandular hairs . . . . .	93
21.2	Internal secretory structures . . . . .	94
	Secretory cells 94, Secretory cavities 95, Secretory canals 96, Laticifers 97	
<b>C. Structures beyond the pericarp</b>	. . . . .	98
22	Placentae and septa . . . . .	98
22.1	Placentae and septa as sources of pulp formation . . . . .	98

22.2	False partition walls and seed pockets . . . . .	100
23	The inferior ovary . . . . .	100
24	The fruits of cacti . . . . .	106
25	Accessory fruit parts . . . . .	118
25.1	Sterile projections of the gymnoecium ("Grannen") . . . . .	119
25.2	Receptacle . . . . .	121
25.3	Sepals . . . . .	122
	Pappus . . . . .	126
25.4	Petals . . . . .	128
	Anthocarps . . . . .	128
25.5	Stamens . . . . .	131
25.6	Fruit scales . . . . .	131
25.7	Prophylls . . . . .	131
25.8	Pedicel and peduncle of the fruit . . . . .	132
25.9	Parts of the inflorescence . . . . .	142
	Inflorescence axis 142, Glumes spikelets 142, Subtending bracts 146, Involucral bracts of inflorescences 146	
 <b>D. Fruit dehiscence . . . . .</b>		<b>148</b>
26	Types of dehiscence . . . . .	148
26.1	General considerations . . . . .	148
26.2	Ventricidal ("marginicidal") dehiscence . . . . .	152
26.3	Septicidal dehiscence . . . . .	153
26.4	Loculicidal dehiscence . . . . .	153
26.5	Dehiscence of the legume . . . . .	156
26.6	Pore capsule . . . . .	163
26.7	Circumscissile dehiscence . . . . .	166
	Pyxidia which develop sclerenchyma and a zone of mechanical weakness Pyxidia which dehisce by formation of meristematic cells in combination with lignified cells . . . . .	166
		168
27	Mechanisms of dehiscence . . . . .	171
27.1	Hygroscopic mechanisms . . . . .	171
	Xerochasy . . . . .	172
	Hygrochasy . . . . .	175
27.2	Turgor mechanisms . . . . .	181
	Spluttermechanisms (Spritzmechanismen) . . . . .	181
	Throwing mechanisms of turgescent fleshy fruits (Schleudermechanismen) Lever catapults (Hebelschleudern) . . . . .	182
	<i>Cyclanthera explodens</i> 182, <i>Impatiens</i> 182, <i>Cardamine</i> 184, <i>Cory-</i> <i>dalis</i> 184, <i>Lathraea</i> 184	182
	Squeezing catapults (Quetschschieleudern) . . . . .	185
	<i>Dorstenia</i> 185, <i>Arctothobium</i> 185	
	Pushing catapults (Stoßschleudern) . . . . .	187
	<i>Polygonum</i> 187	
28	Indehiscent fruits with delayed dehiscence . . . . .	187

<b>E. Types of fruits . . . . .</b>	<b>191</b>
29 The dehiscent fruit: the capsule sensu lato . . . . .	191
29.1 General considerations . . . . .	191
29.2 Special examples . . . . .	194
Solanaceae . . . . .	197
Caryophyllaceae . . . . .	200
The legume . . . . .	203
Vascular anatomy . . . . .	209
Development . . . . .	210
Formation of the "seed-cushion" . . . . .	210
Formation of false septa . . . . .	212
Follicular types with ventricidal dehiscence only . . . . .	215
Evolutionary trends in fruit development of Leguminosae . . . . .	216
Cruciferae (Silique and silicle) . . . . .	218
Acanthaceae . . . . .	225
Scrophulariaceae . . . . .	228
Orchidaceae . . . . .	231
Liliaceae . . . . .	231
<i>Linum usitatissimum</i> . . . . .	233
Fleshy capsules . . . . .	235
<i>Myristica fragrans</i> (the nutmeg) . . . . .	235
<i>Punica granatum</i> (the pomegranate) . . . . .	239
<i>Scyphostegia borneensis</i> . . . . .	244
30 The indehiscent sclerocarpium: the nut . . . . .	244
30.1 The nut sensu stricto . . . . .	246
30.2 The achene . . . . .	258
The genus <i>Centaurea</i> . . . . .	261
Distribution of the parenchyma in the pericarp . . . . .	264
The aerenchyma . . . . .	266
The sclerenchyma . . . . .	267
The "carbon layer" ("Kohleschicht") or black pigment layer . . . . .	270
The hypodermis . . . . .	271
Secretory structures . . . . .	273
Cell contents . . . . .	273
Formation of intercellular spaces . . . . .	274
The epidermis . . . . .	274
Slime cells . . . . .	276
Slime trichomes . . . . .	277
Twin hairs and other trichomes . . . . .	278
Periderm . . . . .	280
Emergences . . . . .	281
Vascular system . . . . .	281
The apical plate . . . . .	281
The receptacle of the inflorescence . . . . .	283
The integument . . . . .	284
Structure of the pappus . . . . .	284
Chaffy bracts and involucral leaves . . . . .	286

The abscission region . . . . .	287
Heterocarpous inflorescences . . . . .	290
Taxonomic significance of the pericarp structure . . . . .	290
30.3 The caryopsis of Gramineae . . . . .	291
General structure . . . . .	291
<i>Zea mays</i> . . . . .	293
Pericarp development . . . . .	296
Semipermeable membrane . . . . .	304
Bracts (glumes and palets) taking part in fruit formation . . . . .	307
30.4 The schizocarpium of Umbelliferae . . . . .	311
General pericarp structure . . . . .	313
The schizocarpium of Hydrocotyloideae . . . . .	314
Secretory spaces . . . . .	318
Interpretation of the inferior ovary . . . . .	320
The carpophore . . . . .	320
Classification based on fruit anatomy . . . . .	324
Evolutionary trends of the schizocarp and derivation from fruits of Araliaceae . . . . .	325
30.5 The mericarps or nutlets ("Klausen") of Boraginaceae . . . . .	330
30.6 The mericarps or nutlets ("Klausen") of Labiatae . . . . .	336
30.7 <i>Anacardium occidentale</i> . . . . .	344
30.8 Winged fruits (samaras) . . . . .	358
31 The fleshy-stony indehiscent fruit: the drupe (sarco-sclerocarpium) . . . . .	368
31.1 General characteristics . . . . .	368
31.2 Zonation of meso- and endocarp . . . . .	372
31.3 The <i>Prunus</i> fruits . . . . .	374
<i>Prunus domestica</i> (The plum) . . . . .	374
<i>Prunus cerasus</i> (The sour cherry) . . . . .	376
<i>Prunus persica</i> (The peach) . . . . .	378
Vascular system . . . . .	379
Cell contents . . . . .	379
Tissue texture . . . . .	380
Size and form of fruits . . . . .	380
Growth periods . . . . .	381
31.4 Species of <i>Rubus</i> . . . . .	381
<i>Rubus idaeus</i> (Red raspberry) . . . . .	381
<i>Rubus fruticosus</i> (Black raspberry) . . . . .	383
<i>Rubus strigosus</i> (American red raspberry) . . . . .	383
31.5 <i>Olea europaea</i> (The olive) . . . . .	386
31.6 <i>Coffea arabica</i> , <i>C. canephora</i> (The coffee) . . . . .	388
31.7 <i>Cocos nucifera</i> (The coconut) . . . . .	400
31.8 Other palm fruits . . . . .	412
31.9 <i>Casimiroa edulis</i> (The white sapote) . . . . .	420
31.10 <i>Mangifera indica</i> (The mango) . . . . .	421
31.11 <i>Ribes</i> (Saxifragaceae) . . . . .	423
31.12 Fruits of Ericaceae: <i>Vaccinium</i> , <i>Gaylussacia</i> . . . . .	423
31.13 <i>Symporicarpus rivularis</i> (cit. <i>S. racemosus</i> ), <i>Cornus mas</i> , <i>Geum urbanum</i>	424

31.14 Juglandaceae . . . . .	425
<i>Carya illinoensis</i> (cit. <i>Hicoria pecan</i> ) 425, <i>Carya ovata</i> and <i>Carya cordiformis</i> 427, <i>Juglans</i> 428	
32 The fleshy indehiscent fruit: the berry (sarcocarpium) . . . . .	431
32.1 General characteristic . . . . .	431
32.2 Types of berries . . . . .	433
<i>Ribes</i> . . . . .	433
<i>Phoenix dactylifera</i> (The date) . . . . .	434
<i>Achras zapota</i> (The sapote) . . . . .	435
<i>Persea americana</i> (The avocado) . . . . .	445
Solanaceae . . . . .	451
General characteristics . . . . .	451
<i>Lycopersicon esculentum</i> (The tomato) . . . . .	454
<i>Capsicum annuum</i> (The red Pepper) . . . . .	457
<i>Musa acuminata</i> and <i>M. balbisiana</i> — cultivars (The banana) . . . . .	461
Fruits of Cucurbitaceae (Type: Pepo) . . . . .	471
Exocarp . . . . .	472
Cork formation . . . . .	475
Mesocarp . . . . .	475
Endocarp . . . . .	477
Vascular bundles . . . . .	477
<i>Carica papaya</i> (The paw paw) . . . . .	477
Anatomy of the ripe fruit . . . . .	480
Development of the fruit . . . . .	480
<i>Theobroma cacao</i> (The Cacao) . . . . .	483
Development of the fruit . . . . .	483
Anatomy of the mature fruit . . . . .	489
Species of <i>Citrus</i> (Type: hesperidium) . . . . .	494
General structure of the <i>Citrus</i> fruit . . . . .	494
Development of the <i>Citrus</i> fruit . . . . .	502
Cell contents . . . . .	511
Other members of Rutaceae . . . . .	516
Loranthaceae . . . . .	517
The clove (dried flower buds of <i>Syzygium aromaticum</i> or <i>Eugenia caryophyllata</i> ) . . . . .	519
33 Aggregate fruits and apocarpous gynoecia . . . . .	522
33.1 Ranunculaceae . . . . .	522
33.2 Annonaceae . . . . .	524
<i>Annona cherimola</i> (The cherimoya) . . . . .	524
Morphology . . . . .	524
Anatomy . . . . .	526
Outer epidermis 526, Mesocarp 526, Inner epidermis 526, Vascular anatomy 526, Cell contents 527	
33.3 Rosaceae . . . . .	527
<i>Fragaria</i> sp. (The strawberry) . . . . .	530
<i>Rosa</i> . . . . .	531

The pome fruits . . . . .	533
<i>Malus sylvestris</i> (The apple) . . . . .	533
Vascular pattern of the apple flower . . . . .	533
Histogenesis . . . . .	540
Histology of the ripe apples . . . . .	540
Epidermis 540, Cork and lenticels 541, Hypodermis 543, Fruit flesh 543, Endocarp 543	
Calyx . . . . .	543
Texture of flesh and air spaces . . . . .	543
Cell contents . . . . .	543
Growth rate, cell size and cell number . . . . .	544
Fruit abscission . . . . .	546
<i>Pyrus communis</i> (The pear) . . . . .	546
Vascular supply . . . . .	546
Histogenesis . . . . .	546
Histology . . . . .	548
Outer epidermis 548, Cork formation 548, Hypodermis 550, Flesh 550, Sclereid differentiation 550, Endocarp 550	
Growth rate . . . . .	552
33.4 The pseudo-apocarpous fruit of <i>Ochna</i> . . . . .	552
34 Infrutescences (collective or multiple fruits) . . . . .	553
34.1 <i>Ananas comosus</i> (The pineapple) . . . . .	553
34.2 <i>Artocarpus altilis</i> (the bread fruit) and related species . . . . .	557
34.3 <i>Ficus carica</i> and related species (The fig) . . . . .	560
<b>F. Fruit abscission</b> . . . . .	565
<b>G. Special dispersal structures and heterodiaspory</b> . . . . .	572
35 Elaiosomes . . . . .	572
36 Floating fruits . . . . .	579
36.1 Floating tissue without air spaces . . . . .	579
36.2 Floating tissue in the form of aerenchyma . . . . .	581
36.3 Floating bladders filled with air . . . . .	582
36.4 Floating tissue developed outside the pericarp . . . . .	582
36.5 Protective layers of floating fruits . . . . .	582
37 Pneumatocarpia and flying organs . . . . .	583
37.1 Pneumatocarpia . . . . .	583
37.2 Flying organs with airspaces . . . . .	585
37.3 Hairs serving as flying organs . . . . .	585
38 Dispersal organs in the form of hooks or burs . . . . .	587
39 Heterodiaspory . . . . .	587
39.1 Heterocarpy . . . . .	591
Heterocarpous Compositae . . . . .	594
Heterocarpy in other families . . . . .	597

39.2 Amphicarpy . . . . .	597
39.3 Heterocarpidy . . . . .	598
39.4 Heteromericarpy . . . . .	598
39.5 Seed pockets . . . . .	599
<b>H. Taxonomic use of the pericarp structure . . . . .</b>	<b>600</b>
<b>I. Phylogeny of the fruit . . . . .</b>	<b>603</b>
General view . . . . .	603
Special examples . . . . .	605
<b>K. Fruits of cultivated plants and polyploidy . . . . .</b>	<b>609</b>
Bibliography . . . . .	615
Index of authors . . . . .	643
Index of latin plant and animal names . . . . .	649
General index . . . . .	666