

<b>1</b>	Biochemistry: An Evolving Science	1
<b>2</b>	Protein Composition and Structure	29
<b>3</b>	Exploring Proteins and Proteomes	69
<b>4</b>	DNA, RNA, and the Flow of Genetic Information	113
<b>5</b>	Exploring Genes and Genomes	145
<b>6</b>	Exploring Evolution and Bioinformatics	185
<b>7</b>	Hemoglobin: Portrait of a Protein in Action	207
<b>8</b>	Enzymes: Basic Concepts and Kinetics	233
<b>9</b>	Catalytic Strategies	273
<b>10</b>	Regulatory Strategies	309
<b>11</b>	Carbohydrates	341
<b>12</b>	Lipids and Cell Membranes	373
<b>13</b>	Membrane Channels and Pumps	403
<b>14</b>	Signal-Transduction Pathways	437
<b>15</b>	Metabolism: Basic Concepts and Design	463
<b>16</b>	Glycolysis and Gluconeogenesis	491
<b>17</b>	The Citric Acid Cycle	541
<b>18</b>	Oxidative Phosphorylation	573
<b>19</b>	The Light Reactions of Photosynthesis	619
<b>20</b>	The Calvin Cycle and the Pentose Phosphate Pathway	645
<b>21</b>	Glycogen Metabolism	679
<b>22</b>	Fatty Acid Metabolism	709
<b>23</b>	Protein Turnover and Amino Acid Catabolism	751
<b>24</b>	The Biosynthesis of Amino Acids	787
<b>25</b>	Nucleotide Biosynthesis	821
<b>26</b>	The Biosynthesis of Membrane Lipids and Steroids	849
<b>27</b>	The Integration of Metabolism	889
<b>28</b>	Drug Development	921
<b>29</b>	DNA Replication, Repair, and Recombination	949
<b>30</b>	RNA Synthesis and Processing	983
<b>31</b>	Protein Synthesis	1021
<b>32</b>	The Control of Gene Expression in Prokaryotes	1057
<b>33</b>	The Control of Gene Expression in Eukaryotes	1075
<b>34</b>	Sensory Systems	1097
<b>35</b>	The Immune System	1119
<b>36</b>	Molecular Motors	1151

⊙ Chapters online only

PREFACE	V
CHAPTER 1	
<b>Biochemistry: An Evolving Science</b>	<b>1</b>
1.1 Biochemical Unity Underlies Biological Diversity	2
1.2 DNA Illustrates the Interplay Between Form and Function	4
DNA is constructed from four building blocks	4
Two single strands of DNA combine to form a double helix	5
DNA structure explains heredity and the storage of information	6
1.3 Concepts from Chemistry Explain the Properties of Biological Molecules	6
The formation of the DNA double helix as a key example	6
The double helix can form from its component strands	6
Covalent and noncovalent bonds are important for the structure and stability of biological molecules	7
The double helix is an expression of the rules of chemistry	10
The laws of thermodynamics govern the behavior of biochemical systems	11
Heat is released in the formation of the double helix	13
Acid–base reactions are central in many biochemical processes	14
Acid–base reactions can disrupt the double helix	15
Buffers regulate pH in organisms and in the laboratory	16
1.4 The Genomic Revolution Is Transforming Biochemistry, Medicine, and Other Fields	17
Genome sequencing has transformed biochemistry and other fields	18
Environmental factors influence human biochemistry	20
Genome sequences encode proteins and patterns of expression	21
APPENDIX	
<b>Visualizing Molecular Structures: Small Molecules</b>	<b>23</b>
APPENDIX:	
<b>Functional Groups</b>	<b>24</b>
CHAPTER 2	
<b>Protein Composition and Structure</b>	<b>29</b>
2.1 Proteins Are Built from a Repertoire of 20 Amino Acids	31
2.2 Primary Structure: Amino Acids Are Linked by Peptide Bonds to Form Polypeptide Chains	37

Proteins have unique amino acid sequences specified by genes	39	A protein purification scheme can be quantitatively evaluated	75
Polypeptide chains are flexible yet conformationally restricted	40	Ultracentrifugation is valuable for separating biomolecules and determining their masses	81
<b>2.3 Secondary Structure: Polypeptide Chains Can Fold into Regular Structures Such As the Alpha Helix, the Beta Sheet, and Turns and Loops</b>	42	Protein purification can be made easier with the use of recombinant DNA technology	82
The alpha helix is a coiled structure stabilized by intrachain hydrogen bonds	42	<b>3.2 Immunology Provides Important Techniques with Which to Investigate Proteins</b>	83
Beta sheets are stabilized by hydrogen bonding between polypeptide strands	44	Antibodies to specific proteins can be generated	83
Polypeptide chains can change direction by making reverse turns and loops	46	Monoclonal antibodies with virtually any desired specificity can be readily prepared	85
<b>2.4 Tertiary Structure: Proteins Can Fold into Globular or Fibrous Structures</b>	46	Proteins can be detected and quantified by using an enzyme-linked immunosorbent assay	85
Fibrous proteins provide structural support for cells and tissues	49	Western blotting permits the detection of proteins separated by gel electrophoresis	87
<b>2.5 Quaternary Structure: Polypeptide Chains Can Assemble into Multisubunit Structures</b>	51	Co-immunoprecipitation enables the identification of binding partners of a protein	88
<b>2.6 The Amino Acid Sequence of a Protein Determines Its Three-Dimensional Structure</b>	52	Fluorescent markers make the visualization of proteins in the cell possible	89
Amino acids have different propensities for forming $\alpha$ helices, $\beta$ sheets, and turns	54	<b>3.3 Mass Spectrometry Is a Powerful Technique for the Identification of Peptides and Proteins</b>	89
Protein folding is a highly cooperative process	55	Peptides can be sequenced by mass spectrometry	92
Proteins fold by progressive stabilization of intermediates rather than by random search	56	Proteins can be specifically cleaved into small peptides to facilitate analysis	93
Prediction of three-dimensional structure from sequence remains a great challenge	57	Genomic and proteomic methods are complementary	94
Some proteins are inherently unstructured and can exist in multiple conformations	58	The amino acid sequence of a protein provides valuable information	95
Protein misfolding and aggregation are associated with some neurological diseases	59	Individual proteins can be identified by mass spectrometry	96
Posttranslational modifications confer new capabilities to proteins	60	<b>3.4 Peptides Can Be Synthesized by Automated Solid-Phase Methods</b>	97
<b>APPENDIX</b>		<b>3.5 Three-Dimensional Protein Structure Can Be Determined by X-ray Crystallography, NMR Spectroscopy, and Cryo-Electron Microscopy</b>	100
<b>Visualizing Molecular Structures: Proteins</b>	<b>64</b>	X-ray crystallography reveals three-dimensional structure in atomic detail	100
<b>CHAPTER 3</b>		Nuclear magnetic resonance spectroscopy can reveal the structures of proteins in solution	102
<b>Exploring Proteins and Proteomes</b>	<b>69</b>	Cryo-electron microscopy is an emerging method of protein structure determination	105
The proteome is the functional representation of the genome	70	<b>APPENDIX</b>	
<b>3.1 The Purification of Proteins Is an Essential First Step in Understanding Their Function</b>	70	<b>Problem-Solving Strategies</b>	<b>108</b>
The assay: How do we recognize the protein we are looking for?	71	<b>CHAPTER 4</b>	
Proteins must be released from the cell to be purified	71	<b>DNA, RNA, and the Flow of Genetic Information</b>	<b>113</b>
Proteins can be purified according to solubility, size, charge, and binding affinity	72	<b>4.1 A Nucleic Acid Consists of Four Kinds of Bases Linked to a Sugar-Phosphate Backbone</b>	114
Proteins can be separated by gel electrophoresis and displayed	75	RNA and DNA differ in the sugar component and one of the bases	114

Nucleotides are the monomeric units of nucleic acids	115	Restriction fragments can be separated by gel electrophoresis and visualized	148
DNA molecules are very long and have directionality	116	DNA can be sequenced by controlled termination of replication	149
<b>4.2 A Pair of Nucleic Acid Strands with Complementary Sequences Can Form a Double-Helical Structure</b>	<b>117</b>	DNA probes and genes can be synthesized by automated solid-phase methods	150
The double helix is stabilized by hydrogen bonds and van der Waals interactions	117	Selected DNA sequences can be greatly amplified by the polymerase chain reaction	151
DNA can assume a variety of structural forms	119	PCR is a powerful technique in medical diagnostics, forensics, and studies of molecular evolution	152
Some DNA molecules are circular and supercoiled	120	The tools for recombinant DNA technology have been used to identify disease-causing mutations	153
Single-stranded nucleic acids can adopt elaborate structures	121	<b>5.2 Recombinant DNA Technology Has Revolutionized All Aspects of Biology</b>	<b>154</b>
<b>4.3 The Double Helix Facilitates the Accurate Transmission of Hereditary Information</b>	<b>122</b>	Restriction enzymes and DNA ligase are key tools in forming recombinant DNA molecules	154
Differences in DNA density established the validity of the semiconservative replication hypothesis	122	Plasmids and $\lambda$ phage are choice vectors for DNA cloning in bacteria	155
The double helix can be reversibly melted	123	Bacterial and yeast artificial chromosomes	157
Unusual circular DNA exists in the eukaryotic nucleus	124	Specific genes can be cloned from digests of genomic DNA	158
<b>4.4 DNA Is Replicated by Polymerases That Take Instructions from Templates</b>	<b>125</b>	Complementary DNA prepared from mRNA can be expressed in host cells	159
DNA polymerase catalyzes phosphodiester-bridge formation	125	Proteins with new functions can be created through directed changes in DNA	160
The genes of some viruses are made of RNA	126	Recombinant methods enable the exploration of the functional effects of disease-causing mutations	163
<b>4.5 Gene Expression Is the Transformation of DNA Information into Functional Molecules</b>	<b>127</b>	<b>5.3 Complete Genomes Have Been Sequenced and Analyzed</b>	<b>163</b>
Several kinds of RNA play key roles in gene expression	127	The genomes of organisms ranging from bacteria to multicellular eukaryotes have been sequenced	164
All cellular RNA is synthesized by RNA polymerases	128	The sequence of the human genome has been completed	165
RNA polymerases take instructions from DNA templates	129	Next-generation sequencing methods enable the rapid determination of a complete genome sequence	166
Transcription begins near promoter sites and ends at terminator sites	130	Comparative genomics has become a powerful research tool	168
Transfer RNAs are the adaptor molecules in protein synthesis	131	<b>5.4 Eukaryotic Genes Can Be Quantitated and Manipulated with Considerable Precision</b>	<b>169</b>
<b>4.6 Amino Acids Are Encoded by Groups of Three Bases Starting from a Fixed Point</b>	<b>132</b>	Gene-expression levels can be comprehensively examined	169
Major features of the genetic code	133	New genes inserted into eukaryotic cells can be efficiently expressed	171
Messenger RNA contains start and stop signals for protein synthesis	134	Transgenic animals harbor and express genes introduced into their germ lines	172
The genetic code is nearly universal	135	Gene disruption and genome editing provide clues to gene function and opportunities for new therapies	172
<b>4.7 Most Eukaryotic Genes Are Mosaics of Introns and Exons</b>	<b>135</b>	RNA interference provides an additional tool for disrupting gene expression	176
RNA processing generates mature RNA	136	Tumor-inducing plasmids can be used to introduce new genes into plant cells	177
Many exons encode protein domains	136	Human gene therapy holds great promise for medicine	178
<b>APPENDIX</b>		<b>APPENDIX</b>	
<b>Problem-Solving Strategies</b>	<b>140</b>	<b>Biochemistry in Focus</b>	<b>180</b>
<b>CHAPTER 5</b>			
<b>Exploring Genes and Genomes</b>	<b>145</b>		
<b>5.1 The Exploration of Genes Relies on Key Tools</b>	<b>146</b>		
Restriction enzymes split DNA into specific fragments	147		

CHAPTER 6		
<b>Exploring Evolution and Bioinformatics</b>	<b>185</b>	
6.1 Homologs Are Descended from a Common Ancestor	186	
6.2 Statistical Analysis of Sequence Alignments Can Detect Homology	187	
The statistical significance of alignments can be estimated by shuffling	189	
Distant evolutionary relationships can be detected through the use of substitution matrices	190	
Databases can be searched to identify homologous sequences	193	
6.3 Examination of Three-Dimensional Structure Enhances Our Understanding of Evolutionary Relationships	194	
Tertiary structure is more conserved than primary structure	194	
Knowledge of three-dimensional structures can aid in the evaluation of sequence alignments	195	
Repeated motifs can be detected by aligning sequences with themselves	196	
Convergent evolution illustrates common solutions to biochemical challenges	196	
Comparison of RNA sequences can be a source of insight into RNA secondary structures	197	
6.4 Evolutionary Trees Can Be Constructed on the Basis of Sequence Information	198	
Horizontal gene transfer events may explain unexpected branches of the evolutionary tree	199	
6.5 Modern Techniques Make the Experimental Exploration of Evolution Possible	200	
Ancient DNA can sometimes be amplified and sequenced	200	
Molecular evolution can be examined experimentally	200	
APPENDIX		
<b>Biochemistry in Focus</b>	<b>203</b>	
APPENDIX:		
<b>Problem-Solving Strategies</b>	<b>204</b>	
CHAPTER 7		
<b>Hemoglobin: Portrait of a Protein in Action</b>	<b>207</b>	
7.1 Binding of Oxygen by Heme Iron	208	
Changes in heme electronic structure upon oxygen binding are the basis for functional imaging studies	210	
The structure of myoglobin prevents the release of reactive oxygen species	210	
Human hemoglobin is an assembly of four myoglobin-like subunits	211	
7.2 Hemoglobin Binds Oxygen Cooperatively	212	
Oxygen binding markedly changes the quaternary structure of hemoglobin	213	
Hemoglobin cooperativity can be potentially explained by several models	214	
Structural changes at the heme groups are transmitted to the $\alpha_1\beta_1-\alpha_2\beta_2$ interface	215	
2,3-Bisphosphoglycerate in red cells is crucial in determining the oxygen affinity of hemoglobin	216	
Carbon monoxide can disrupt oxygen transport by hemoglobin	217	
7.3 Hydrogen Ions and Carbon Dioxide Promote the Release of Oxygen: The Bohr Effect	217	
7.4 Mutations in Genes Encoding Hemoglobin Subunits Can Result in Disease	220	
Sickle-cell anemia results from the aggregation of mutated deoxyhemoglobin molecules	220	
Thalassemia is caused by an imbalanced production of hemoglobin chains	221	
The accumulation of free $\alpha$ -hemoglobin chains is prevented	222	
Additional globins are encoded in the human genome	223	
APPENDIX		
<b>Binding Models Can Be Formulated in Quantitative Terms: The Hill Plot and the Concerted Model</b>	<b>225</b>	
APPENDIX		
<b>Biochemistry in Focus</b>	<b>227</b>	
CHAPTER 8		
<b>Enzymes: Basic Concepts and Kinetics</b>	<b>233</b>	
8.1 Enzymes Are Powerful and Highly Specific Catalysts	234	
Many enzymes require cofactors for activity	235	
Enzymes can transform energy from one form into another	236	
8.2 Gibbs Free Energy Is a Useful Thermodynamic Function for Understanding Enzymes	236	
The free-energy change provides information about the spontaneity but not the rate of a reaction	237	
The standard free-energy change of a reaction is related to the equilibrium constant	237	
Enzymes alter only the reaction rate and not the reaction equilibrium	239	
8.3 Enzymes Accelerate Reactions by Facilitating the Formation of the Transition State	240	
The formation of an enzyme-substrate complex is the first step in enzymatic catalysis	241	
The active sites of enzymes have some common features	242	
The binding energy between enzyme and substrate is important for catalysis	243	

<b>8.4</b>	<b>The Michaelis–Menten Model Accounts for the Kinetic Properties of Many Enzymes</b>	<b>244</b>	<b>9.2</b>	<b>Carbonic Anhydrases Make a Fast Reaction Faster</b>	<b>286</b>
	Kinetics is the study of reaction rates	244		Carbonic anhydrase contains a bound zinc ion essential for catalytic activity	287
	The steady-state assumption facilitates a description of enzyme kinetics	245		Catalysis entails zinc activation of a water molecule	288
	Variations in $K_M$ can have physiological consequences	247		A proton shuttle facilitates rapid regeneration of the active form of the enzyme	290
	$K_M$ and $V_{max}$ values can be determined by several means	247	<b>9.3</b>	<b>Restriction Enzymes Catalyze Highly Specific DNA-Cleavage Reactions</b>	<b>291</b>
	$K_M$ and $V_{max}$ values are important enzyme characteristics	248		Cleavage is by in-line displacement of 3'-oxygen from phosphorus by magnesium-activated water	292
	$k_{cat}/K_M$ is a measure of catalytic efficiency	249		Restriction enzymes require magnesium for catalytic activity	294
	Most biochemical reactions include multiple substrates	251		The complete catalytic apparatus is assembled only within complexes of cognate DNA molecules, ensuring specificity	295
	Allosteric enzymes do not obey Michaelis–Menten kinetics	252		Host-cell DNA is protected by the addition of methyl groups to specific bases	297
<b>8.5</b>	<b>Enzymes Can Be Inhibited by Specific Molecules</b>	<b>253</b>		Type II restriction enzymes have a catalytic core in common and are probably related by horizontal gene transfer	298
	The different types of reversible inhibitors are kinetically distinguishable	254	<b>9.4</b>	<b>Myosins Harness Changes in Enzyme Conformation to Couple ATP Hydrolysis to Mechanical Work</b>	<b>298</b>
	Irreversible inhibitors can be used to map the active site	256		ATP hydrolysis proceeds by the attack of water on the gamma phosphoryl group	299
	Penicillin irreversibly inactivates a key enzyme in bacterial cell-wall synthesis	258		Formation of the transition state for ATP hydrolysis is associated with a substantial conformational change	300
	Transition-state analogs are potent inhibitors of enzymes	260		The altered conformation of myosin persists for a substantial period of time	301
	Enzymes have impact outside the laboratory or clinic	261		Scientists can watch single molecules of myosin move	303
<b>8.6</b>	<b>Enzymes Can Be Studied One Molecule at a Time</b>	<b>261</b>		Myosins are a family of enzymes containing P-loop structures	303
APPENDIX					
	<b>Enzymes Are Classified on the Basis of the Types of Reactions That They Catalyze</b>	<b>264</b>	APPENDIX		
APPENDIX					
	<b>Biochemistry in Focus</b>	<b>265</b>	<b>Problem-Solving Strategies</b>		
APPENDIX					
	<b>Problem-Solving Strategies</b>	<b>266</b>	APPENDIX		
CHAPTER 9					
	<b>Catalytic Strategies</b>	<b>273</b>	<b>Problem-Solving Strategies</b>		
	A few basic catalytic principles are used by many enzymes	274	<b>306</b>		
<b>9.1</b>	<b>Proteases Facilitate a Fundamentally Difficult Reaction</b>	<b>275</b>	CHAPTER 10		
	Chymotrypsin possesses a highly reactive serine residue	276	<b>Regulatory Strategies</b>		
	Chymotrypsin action proceeds in two steps linked by a covalently bound intermediate	277	<b>309</b>		
	Serine is part of a catalytic triad that also includes histidine and aspartate	278	<b>10.1</b>	<b>Aspartate Transcarbamoylase Is Allosterically Inhibited by the End Product of Its Pathway</b>	<b>310</b>
	Catalytic triads are found in other hydrolytic enzymes	281		Allosterically regulated enzymes do not follow Michaelis–Menten kinetics	311
	The catalytic triad has been dissected by site-directed mutagenesis	282		ATCase consists of separable catalytic and regulatory subunits	312
	Cysteine, aspartyl, and metalloproteases are other major classes of peptide-cleaving enzymes	283		Allosteric interactions in ATCase are mediated by large changes in quaternary structure	312
	Protease inhibitors are important drugs	285		Allosteric regulators modulate the T-to-R equilibrium	316
			<b>10.2</b>	<b>Isozymes Provide a Means of Regulation Specific to Distinct Tissues and Developmental Stages</b>	<b>317</b>

<b>10.3</b>	<b>Covalent Modification Is a Means of Regulating Enzyme Activity</b>	<b>318</b>	Cellulose, a structural component of plants, is made of chains of glucose	350
	Kinases and phosphatases control the extent of protein phosphorylation	319	Human milk oligosaccharides protect newborns from infection	352
	Phosphorylation is a highly effective means of regulating the activities of target proteins	321	<b>11.3</b> Carbohydrates Can Be Linked to Proteins to Form Glycoproteins	<b>352</b>
	Cyclic AMP activates protein kinase A by altering the quaternary structure	321	Carbohydrates can be linked to proteins through asparagine ( <i>N</i> -linked) or through serine or threonine ( <i>O</i> -linked) residues	352
	Mutations in protein kinase A can cause Cushing's syndrome	322	The glycoprotein erythropoietin is a vital hormone	353
	Exercise modifies the phosphorylation of many proteins	323	Glycosylation functions in nutrient sensing	354
<b>10.4</b>	<b>Many Enzymes Are Activated by Specific Proteolytic Cleavage</b>	<b>323</b>	Proteoglycans, composed of polysaccharides and protein, have important structural roles	354
	Chymotrypsinogen is activated by specific cleavage of a single peptide bond	324	Proteoglycans are important components of cartilage	355
	Proteolytic activation of chymotrypsinogen leads to the formation of a substrate-binding site	325	Mucins are glycoprotein components of mucus	356
	The generation of trypsin from trypsinogen leads to the activation of other zymogens	325	Chitin can be processed to a molecule with a variety of uses	357
	Some proteolytic enzymes have specific inhibitors	326	Protein glycosylation takes place in the lumen of the endoplasmic reticulum and in the Golgi complex	357
	Serpins can be degraded by a unique enzyme	328	Specific enzymes are responsible for oligosaccharide assembly	358
	Blood clotting is accomplished by a cascade of zymogen activations	328	Blood groups are based on protein glycosylation patterns	359
	Prothrombin must bind to Ca <sup>2+</sup> to be converted to thrombin	329	Errors in glycosylation can result in pathological conditions	359
	Fibrinogen is converted by thrombin into a fibrin clot	329	Oligosaccharides can be "sequenced"	360
	Vitamin K is required for the formation of $\gamma$ -carboxyglutamate	331	<b>11.4</b> Lectins Are Specific Carbohydrate-Binding Proteins	<b>361</b>
	The clotting process must be precisely regulated	332	Lectins promote interactions between cells and within cells	361
	Hemophilia revealed an early step in clotting	333	Lectins are organized into different classes	362
			Influenza virus binds to sialic acid residues	363
<b>APPENDIX</b>			<b>APPENDIX</b>	
	<b>Biochemistry in Focus</b>	<b>335</b>	<b>Biochemistry in Focus</b>	<b>365</b>
<b>APPENDIX</b>			<b>APPENDIX</b>	
	<b>Problem-Solving Strategies</b>	<b>336</b>	<b>Problem-Solving Strategies</b>	<b>366</b>
<b>CHAPTER 11</b>			<b>CHAPTER 12</b>	
	<b>Carbohydrates</b>	<b>341</b>	<b>Lipids and Cell Membranes</b>	<b>373</b>
<b>11.1</b>	<b>Monosaccharides Are the Simplest Carbohydrates</b>	<b>342</b>	Many Common Features Underlie the Diversity of Biological Membranes	374
	Many common sugars exist in cyclic forms	344	<b>12.1</b> Fatty Acids Are Key Constituents of Lipids	<b>374</b>
	Pyranose and furanose rings can assume different conformations	346	Fatty acid names are based on their parent hydrocarbons	374
	Glucose is a reducing sugar	347	Fatty acids vary in chain length and degree of unsaturation	375
	Monosaccharides are joined to alcohols and amines through glycosidic bonds	348	<b>12.2</b> There Are Three Common Types of Membrane Lipids	<b>376</b>
	Phosphorylated sugars are key intermediates in energy generation and biosyntheses	348	Phospholipids are the major class of membrane lipids	376
<b>11.2</b>	<b>Monosaccharides Are Linked to Form Complex Carbohydrates</b>	<b>349</b>		
	Sucrose, lactose, and maltose are the common disaccharides	349		
	Glycogen and starch are storage forms of glucose	350		

Membrane lipids can include carbohydrate moieties	378	Multidrug resistance highlights a family of membrane pumps with ATP-binding cassette domains	411
Cholesterol is a lipid based on a steroid nucleus	378		
Archaeal membranes are built from ether lipids with branched chains	379	<b>13.3</b> Lactose Permease Is an Archetype of Secondary Transporters That Use One Concentration Gradient to Power the Formation of Another	413
A membrane lipid is an amphipathic molecule containing a hydrophilic and a hydrophobic moiety	379		
<b>12.3</b> Phospholipids and Glycolipids Readily Form Bimolecular Sheets in Aqueous Media	380	<b>13.4</b> Specific Channels Can Rapidly Transport Ions Across Membranes	415
Lipid vesicles can be formed from phospholipids	381	Action potentials are mediated by transient changes in Na <sup>+</sup> and K <sup>+</sup> permeability	415
Lipid bilayers are highly impermeable to ions and most polar molecules	382	Patch-clamp conductance measurements reveal the activities of single channels	416
<b>12.4</b> Proteins Carry Out Most Membrane Processes	383	The structure of a potassium ion channel is an archetype for many ion-channel structures	417
Proteins associate with the lipid bilayer in a variety of ways	384	The structure of the potassium ion channel reveals the basis of ion specificity	418
Proteins interact with membranes in a variety of ways	384	The structure of the potassium ion channel explains its rapid rate of transport	420
Some proteins associate with membranes through covalently attached hydrophobic groups	388	Voltage gating requires substantial conformational changes in specific ion-channel domains	421
Transmembrane helices can be accurately predicted from amino acid sequences	388	A channel can be inactivated by occlusion of the pore: the ball-and-chain model	422
<b>12.5</b> Lipids and Many Membrane Proteins Diffuse Rapidly in the Plane of the Membrane	390	The acetylcholine receptor is an archetype for ligand-gated ion channels	423
The fluid mosaic model allows lateral movement but not rotation through the membrane	391	Action potentials integrate the activities of several ion channels working in concert	424
Membrane fluidity is controlled by fatty acid composition and cholesterol content	391	Disruption of ion channels by mutations or chemicals can be potentially life-threatening	426
Lipid rafts are highly dynamic complexes formed between cholesterol and specific lipids	392	<b>13.5</b> Gap Junctions Allow Ions and Small Molecules to Flow Between Communicating Cells	427
All biological membranes are asymmetric	393		
<b>12.6</b> Eukaryotic Cells Contain Compartments Bounded by Internal Membranes	393	<b>13.6</b> Specific Channels Increase the Permeability of Some Membranes to Water	429
APPENDIX		APPENDIX	
<b>Biochemistry in Focus</b>	<b>399</b>	<b>Biochemistry in Focus</b>	<b>431</b>
CHAPTER 13		APPENDIX	
<b>Membrane Channels and Pumps</b>	<b>403</b>	<b>Problem-Solving Strategies</b>	<b>432</b>
The expression of transporters largely defines the metabolic activities of a given cell type	404		
<b>13.1</b> The Transport of Molecules Across a Membrane May Be Active or Passive	404	CHAPTER 14	
Many molecules require protein transporters to cross membranes	405	<b>Signal-Transduction Pathways</b>	<b>437</b>
Free energy stored in concentration gradients can be quantified	405	Signal transduction depends on molecular circuits	438
<b>13.2</b> Two Families of Membrane Proteins Use ATP Hydrolysis to Pump Ions and Molecules Across Membranes	406	<b>14.1</b> Epinephrine and Angiotensin II Signaling: Heterotrimeric G Proteins Transmit Signals and Reset Themselves	439
P-type ATPases couple phosphorylation and conformational changes to pump calcium ions across membranes	407	Ligand binding to 7TM receptors leads to the activation of heterotrimeric G proteins	440
Digitalis specifically inhibits the Na <sup>+</sup> – K <sup>+</sup> pump by blocking its dephosphorylation	410	Activated G proteins transmit signals by binding to other proteins	442
P-type ATPases are evolutionarily conserved and play a wide range of roles	410	Cyclic AMP stimulates the phosphorylation of many target proteins by activating protein kinase A	443
		G proteins spontaneously reset themselves through GTP hydrolysis	443
		Some 7TM receptors activate the phosphoinositide cascade	444

Calcium ion is a widely used second messenger	445	Phosphoryl-transfer potential is an important form of cellular energy transformation	470
Calcium ion often activates the regulatory protein calmodulin	447	ATP may have roles other than in energy and signal transduction	472
<b>14.2 Insulin Signaling: Phosphorylation Cascades Are Central to Many Signal-Transduction Processes</b>	447	<b>15.3 The Oxidation of Carbon Fuels Is an Important Source of Cellular Energy</b>	472
The insulin receptor is a dimer that closes around a bound insulin molecule	448	Compounds with high phosphoryl-transfer potential can couple carbon oxidation to ATP synthesis	473
Insulin binding results in the cross-phosphorylation and activation of the insulin receptor	448	Ion gradients across membranes provide an important form of cellular energy that can be coupled to ATP synthesis	474
The activated insulin-receptor kinase initiates a kinase cascade	449	Phosphates play a prominent role in biochemical processes	474
Insulin signaling is terminated by the action of phosphatases	451	Energy from foodstuffs is extracted in three stages	474
<b>14.3 EGF Signaling: Signal-Transduction Pathways Are Poised to Respond</b>	451	<b>15.4 Metabolic Pathways Contain Many Recurring Motifs</b>	475
EGF binding results in the dimerization of the EGF receptor	451	Activated carriers exemplify the modular design and economy of metabolism	475
The EGF receptor undergoes phosphorylation of its carboxyl-terminal tail	452	Many activated carriers are derived from vitamins	478
EGF signaling leads to the activation of Ras, a small G protein	453	Key reactions are reiterated throughout metabolism	480
Activated Ras initiates a protein kinase cascade	453	Metabolic processes are regulated in three principal ways	483
EGF signaling is terminated by protein phosphatases and the intrinsic GTPase activity of Ras	454	Aspects of metabolism may have evolved from an RNA world	485
<b>14.4 Many Elements Recur with Variation in Different Signal-Transduction Pathways</b>	454	APPENDIX	
<b>14.5 Defects in Signal-Transduction Pathways Can Lead to Cancer and Other Diseases</b>	455	<b>Problem-Solving Strategies</b>	<b>487</b>
Monoclonal antibodies can be used to inhibit signal-transduction pathways activated in tumors	456	CHAPTER 16	
Protein kinase inhibitors can be effective anticancer drugs	457	<b>Glycolysis and Gluconeogenesis</b>	<b>491</b>
Cholera and whooping cough are the result of altered G-protein activity	457	Glucose is generated from dietary carbohydrates	492
APPENDIX		Glucose is an important fuel for most organisms	493
<b>Biochemistry in Focus</b>	<b>459</b>	<b>16.1 Glycolysis Is an Energy-Conversion Pathway in Many Organisms</b>	493
CHAPTER 15		The enzymes of glycolysis are associated with one another	493
<b>Metabolism: Basic Concepts and Design</b>	<b>463</b>	Glycolysis can be divided into two parts	493
<b>15.1 Metabolism Is Composed of Many Coupled, Interconnecting Reactions</b>	464	Hexokinase traps glucose in the cell and begins glycolysis	495
Metabolism consists of energy-yielding and energy-requiring reactions	464	Fructose 1,6-bisphosphate is generated from glucose 6-phosphate	496
A thermodynamically unfavorable reaction can be driven by a favorable reaction	465	The six-carbon sugar is cleaved into two three-carbon fragments	497
<b>15.2 ATP Is the Universal Currency of Free Energy in Biological Systems</b>	466	Mechanism: Triose phosphate isomerase salvages a three-carbon fragment	498
ATP hydrolysis is exergonic	466	The oxidation of an aldehyde to an acid powers the formation of a compound with high phosphoryl-transfer potential	499
ATP hydrolysis drives metabolism by shifting the equilibrium of coupled reactions	468	Mechanism: Phosphorylation is coupled to the oxidation of glyceraldehyde 3-phosphate by a thioester intermediate	500
The high phosphoryl potential of ATP results from structural differences between ATP and its hydrolysis products	469	ATP is formed by phosphoryl transfer from 1,3-bisphosphoglycerate	501
		Additional ATP is generated with the formation of pyruvate	503



Two ATP molecules are formed in the conversion of glucose into pyruvate	504	APPENDIX	
NAD <sup>+</sup> is regenerated from the metabolism of pyruvate	505	<b>BIOCHEMISTRY IN FOCUS</b>	<b>532</b>
Fermentations provide usable energy in the absence of oxygen	507	<b>Biochemistry in Focus 1</b>	<b>532</b>
Fructose is converted into glycolytic intermediates by fructokinase	508	<b>Biochemistry in Focus 2</b>	<b>533</b>
Excessive fructose consumption can lead to pathological conditions	508	APPENDIX	
Galactose is converted into glucose 6-phosphate	509	<b>Problem-Solving Strategies</b>	<b>533</b>
Many adults are intolerant of milk because they are deficient in lactase	510	CHAPTER 17	
Galactose is highly toxic if the transferase is missing	511	<b>The Citric Acid Cycle</b>	<b>541</b>
<b>16.2 The Glycolytic Pathway Is Tightly Controlled</b>	<b>511</b>	The citric acid cycle harvests high-energy electrons	542
Glycolysis in muscle is regulated to meet the need for ATP	512	<b>17.1 The Pyruvate Dehydrogenase Complex Links Glycolysis to the Citric Acid Cycle</b>	<b>543</b>
The regulation of glycolysis in the liver illustrates the biochemical versatility of the liver	513	Mechanism: The synthesis of acetyl coenzyme A from pyruvate requires three enzymes and five coenzymes	544
A family of transporters enables glucose to enter and leave animal cells	516	Flexible linkages allow lipoamide to move between different active sites	546
Aerobic glycolysis is a property of rapidly growing cells	517	<b>17.2 The Citric Acid Cycle Oxidizes Two-Carbon Units</b>	<b>548</b>
Cancer and endurance training affect glycolysis in a similar fashion	518	Citrate synthase forms citrate from oxaloacetate and acetyl coenzyme A	548
<b>16.3 Glucose Can Be Synthesized from Noncarbohydrate Precursors</b>	<b>519</b>	Mechanism: The mechanism of citrate synthase prevents undesirable reactions	549
Gluconeogenesis is not a reversal of glycolysis	519	Citrate is isomerized into isocitrate	550
The conversion of pyruvate into phosphoenolpyruvate begins with the formation of oxaloacetate	521	Isocitrate is oxidized and decarboxylated to alpha-ketoglutarate	551
Oxaloacetate is shuttled into the cytoplasm and converted into phosphoenolpyruvate	522	Succinyl coenzyme A is formed by the oxidative decarboxylation of alpha-ketoglutarate	552
The conversion of fructose 1,6-bisphosphate into fructose 6-phosphate and orthophosphate is an irreversible step	523	A compound with high phosphoryl-transfer potential is generated from succinyl coenzyme A	552
The generation of free glucose is an important control point	523	Mechanism: Succinyl coenzyme A synthetase transforms types of biochemical energy	553
Six high-transfer-potential phosphoryl groups are spent in synthesizing glucose from pyruvate	524	Oxaloacetate is regenerated by the oxidation of succinate	554
<b>16.4 Gluconeogenesis and Glycolysis Are Reciprocally Regulated</b>	<b>525</b>	The citric acid cycle produces high-transfer-potential electrons, ATP, and CO <sub>2</sub>	555
Energy charge determines whether glycolysis or gluconeogenesis will be most active	525	<b>17.3 Entry to the Citric Acid Cycle and Metabolism Through It Are Controlled</b>	<b>557</b>
The balance between glycolysis and gluconeogenesis in the liver is sensitive to blood-glucose concentration	526	The pyruvate dehydrogenase complex is regulated allosterically and by reversible phosphorylation	557
Substrate cycles amplify metabolic signals and produce heat	528	The citric acid cycle is controlled at several points	559
Lactate and alanine formed by contracting muscle are used by other organs	528	Defects in the citric acid cycle contribute to the development of cancer	559
Glycolysis and gluconeogenesis are evolutionarily intertwined	530	An enzyme in lipid metabolism is hijacked to inhibit pyruvate dehydrogenase activity	560
		<b>17.4 The Citric Acid Cycle Is a Source of Biosynthetic Precursors</b>	<b>561</b>
		The citric acid cycle must be capable of being rapidly replenished	561

The disruption of pyruvate metabolism is the cause of beriberi and poisoning by mercury and arsenic	562		
The citric acid cycle may have evolved from preexisting pathways	563		
<b>17.5 The Glyoxylate Cycle Enables Plants and Bacteria to Grow on Acetate</b>	<b>564</b>		
<b>APPENDIX</b>			
<b>BIOCHEMISTRY IN FOCUS</b>	<b>566</b>		
<b>Biochemistry in Focus 1</b>	<b>566</b>		
<b>Biochemistry in Focus 2</b>	<b>567</b>		
<b>APPENDIX</b>			
<b>Problem-Solving Strategies</b>	<b>568</b>		
<b>CHAPTER 18</b>			
<b>Oxidative Phosphorylation</b>	<b>573</b>		
<b>18.1 Eukaryotic Oxidative Phosphorylation Takes Place in Mitochondria</b>	<b>574</b>		
Mitochondria are bounded by a double membrane	574		
Mitochondria are the result of an endosymbiotic event	575		
<b>18.2 Oxidative Phosphorylation Depends on Electron Transfer</b>	<b>576</b>		
The electron-transfer potential of an electron is measured as redox potential	576		
Electron flow from NADH to molecular oxygen powers the formation of a proton gradient	579		
<b>18.3 The Respiratory Chain Consists of Four Complexes: Three Proton Pumps and a Physical Link to the Citric Acid Cycle</b>	<b>579</b>		
Iron-sulfur clusters are common components of the electron-transport chain	581		
The high-potential electrons of NADH enter the respiratory chain at NADH-Q oxidoreductase	582		
Ubiquinol is the entry point for electrons from FADH <sub>2</sub> of flavoproteins	584		
Electrons flow from ubiquinol to cytochrome <i>c</i> through Q-cytochrome <i>c</i> oxidoreductase	584		
The Q cycle funnels electrons from a two-electron carrier to a one-electron carrier and pumps protons	584		
Cytochrome <i>c</i> oxidase catalyzes the reduction of molecular oxygen to water	586		
Most of the electron-transport chain is organized into a complex called the respirasome	589		
Toxic derivatives of molecular oxygen such as superoxide radicals are scavenged by protective enzymes	590		
Electrons can be transferred between groups that are not in contact	591		
The conformation of cytochrome <i>c</i> has remained essentially constant for more than a billion years	592		
<b>18.4 A Proton Gradient Powers the Synthesis of ATP</b>	<b>592</b>		
ATP synthase is composed of a proton-conducting unit and a catalytic unit	594		
Proton flow through ATP synthase leads to the release of tightly bound ATP: The binding-change mechanism	595		
Rotational catalysis is the world's smallest molecular motor	597		
Proton flow around the <i>c</i> ring powers ATP synthesis	597		
ATP synthase and G proteins have several common features	599		
<b>18.5 Many Shuttles Allow Movement Across Mitochondrial Membranes</b>	<b>600</b>		
Electrons from cytoplasmic NADH enter mitochondria by shuttles	600		
The entry of ADP into mitochondria is coupled to the exit of ATP by ATP-ADP translocase	601		
Mitochondrial transporters for metabolites have a common tripartite structure	602		
<b>18.6 The Regulation of Cellular Respiration Is Governed Primarily by the Need for ATP</b>	<b>603</b>		
The complete oxidation of glucose yields about 30 molecules of ATP	603		
The rate of oxidative phosphorylation is determined by the need for ATP	603		
ATP synthase can be regulated	604		
Regulated uncoupling leads to the generation of heat	605		
Reintroduction of UCP-1 into pigs may be economically valuable	607		
Oxidative phosphorylation can be inhibited at many stages	607		
Mitochondrial diseases are being discovered	608		
Mitochondria play a key role in apoptosis	609		
Power transmission by proton gradients is a central motif of bioenergetics	609		
<b>APPENDIX</b>			
<b>Biochemistry in Focus</b>	<b>611</b>		
<b>APPENDIX</b>			
<b>Problem-Solving Strategies</b>	<b>612</b>		
<b>CHAPTER 19</b>			
<b>The Light Reactions of Photosynthesis</b>	<b>619</b>		
Photosynthesis converts light energy into chemical energy	620		
<b>19.1 Photosynthesis Takes Place in Chloroplasts</b>	<b>621</b>		
The primary events of photosynthesis take place in thylakoid membranes	621		
Chloroplasts arose from an endosymbiotic event	622		
<b>19.2 Light Absorption by Chlorophyll Induces Electron Transfer</b>	<b>622</b>		
A special pair of chlorophylls initiate charge separation	623		

Cyclic electron flow reduces the cytochrome of the reaction center	625	Three ATP and two NADPH molecules are used to bring carbon dioxide to the level of a hexose	654
<b>19.3 Two Photosystems Generate a Proton Gradient and NADPH in Oxygenic Photosynthesis</b>	<b>626</b>	Starch and sucrose are the major carbohydrate stores in plants	654
Photosystem II transfers electrons from water to plastoquinone and generates a proton gradient	626	<b>20.2 The Activity of the Calvin Cycle Depends on Environmental Conditions</b>	<b>655</b>
Cytochrome <i>bf</i> links photosystem II to photosystem I	629	Rubisco is activated by light-driven changes in proton and magnesium ion concentrations	655
Photosystem I uses light energy to generate reduced ferredoxin, a powerful reductant	629	Thioredoxin plays a key role in regulating the Calvin cycle	655
Ferredoxin–NADP <sup>+</sup> reductase converts NADP <sup>+</sup> into NADPH	630	The C <sub>4</sub> pathway of tropical plants accelerates photosynthesis by concentrating carbon dioxide	656
<b>19.4 A Proton Gradient across the Thylakoid Membrane Drives ATP Synthesis</b>	<b>631</b>	Crassulacean acid metabolism permits growth in arid ecosystems	659
The ATP synthase of chloroplasts closely resembles those of mitochondria and prokaryotes	632	<b>20.3 The Pentose Phosphate Pathway Generates NADPH and Synthesizes Five-Carbon Sugars</b>	<b>659</b>
The activity of chloroplast ATP synthase is regulated	633	Two molecules of NADPH are generated in the conversion of glucose 6-phosphate into ribulose 5-phosphate	659
Cyclic electron flow through photosystem I leads to the production of ATP instead of NADPH	634	The pentose phosphate pathway and glycolysis are linked by transketolase and transaldolase	660
The absorption of eight photons yields one O <sub>2</sub> , two NADPH, and three ATP molecules	634	Mechanism: Transketolase and transaldolase stabilize carbanionic intermediates by different mechanisms	662
<b>19.5 Accessory Pigments Funnel Energy into Reaction Centers</b>	<b>635</b>	<b>20.4 The Metabolism of Glucose 6-Phosphate by the Pentose Phosphate Pathway Is Coordinated with Glycolysis</b>	<b>665</b>
Resonance energy transfer allows energy to move from the site of initial absorbance to the reaction center	636	The rate of the oxidative phase of the pentose phosphate pathway is controlled by the level of NADP <sup>+</sup>	665
The components of photosynthesis are highly organized	637	The flow of glucose 6-phosphate depends on the need for NADPH, ribose 5-phosphate, and ATP	666
Many herbicides inhibit the light reactions of photosynthesis	638	The pentose phosphate pathway is required for rapid cell growth	668
<b>19.6 The Ability to Convert Light into Chemical Energy Is Ancient</b>	<b>638</b>	Through the looking-glass: The Calvin cycle and the pentose phosphate pathway are mirror images	668
Artificial photosynthetic systems may provide clean, renewable energy	639	<b>20.5 Glucose 6-Phosphate Dehydrogenase Plays a Key Role in Protection Against Reactive Oxygen Species</b>	<b>668</b>
APPENDIX		Glucose 6-phosphate dehydrogenase deficiency causes a drug-induced hemolytic anemia	669
<b>Biochemistry in Focus</b>	<b>641</b>	A deficiency of glucose 6-phosphate dehydrogenase confers an evolutionary advantage in some circumstances	670
APPENDIX		APPENDIX	
<b>Problem-Solving Strategies</b>	<b>641</b>	<b>BIOCHEMISTRY IN FOCUS</b>	<b>672</b>
CHAPTER 20		<b>Biochemistry in Focus 1</b>	<b>672</b>
<b>The Calvin Cycle and the Pentose Phosphate Pathway</b>	<b>645</b>	<b>Biochemistry in Focus 2</b>	<b>672</b>
<b>20.1 The Calvin Cycle Synthesizes Hexoses from Carbon Dioxide and Water</b>	<b>646</b>	APPENDIX	
Carbon dioxide reacts with ribulose 1,5-bisphosphate to form two molecules of 3-phosphoglycerate	646	<b>Problem-Solving Strategies</b>	<b>673</b>
Rubisco activity depends on magnesium and carbamate	648		
Rubisco activase is essential for rubisco activity	649		
Rubisco also catalyzes a wasteful oxygenase reaction: Catalytic imperfection	649		
Hexose phosphates are made from phosphoglycerate, and ribulose 1,5-bisphosphate is regenerated	651		

CHAPTER 21		A biochemical understanding of glycogen-storage diseases is possible	700
<b>Glycogen Metabolism</b>	<b>679</b>		
Glycogen metabolism is the regulated release and storage of glucose	680		
<b>21.1 Glycogen Breakdown Requires the Interplay of Several Enzymes</b>	<b>681</b>		
Phosphorylase catalyzes the phosphorolytic cleavage of glycogen to release glucose 1-phosphate	681		
Mechanism: Pyridoxal phosphate participates in the phosphorolytic cleavage of glycogen	682		
A debranching enzyme also is needed for the breakdown of glycogen	684		
Phosphoglucomutase converts glucose 1-phosphate into glucose 6-phosphate	685		
The liver contains glucose 6-phosphatase, a hydrolytic enzyme absent from muscle	685		
<b>21.2 Phosphorylase Is Regulated by Allosteric Interactions and Reversible Phosphorylation</b>	<b>686</b>		
Liver phosphorylase produces glucose for use by other tissues	686		
Muscle phosphorylase is regulated by the intracellular energy charge	687		
Biochemical characteristics of muscle fiber types differ	688		
Phosphorylation promotes the conversion of phosphorylase <i>b</i> to phosphorylase <i>a</i>	688		
Phosphorylase kinase is activated by phosphorylation and calcium ions	689		
An isozymic form of glycogen phosphorylase exists in the brain	690		
<b>21.3 Epinephrine and Glucagon Signal the Need for Glycogen Breakdown</b>	<b>690</b>		
G proteins transmit the signal for the initiation of glycogen breakdown	690		
Glycogen breakdown must be rapidly turned off when necessary	692		
The regulation of glycogen phosphorylase became more sophisticated as the enzyme evolved	692		
<b>21.4 Glycogen Synthesis Requires Several Enzymes and Uridine Diphosphate Glucose</b>	<b>693</b>		
UDP-glucose is an activated form of glucose	693		
Glycogen synthase catalyzes the transfer of glucose from UDP-glucose to a growing chain	693		
A branching enzyme forms $\alpha$ -1,6 linkages	694		
Glycogen synthase is the key regulatory enzyme in glycogen synthesis	695		
Glycogen is an efficient storage form of glucose	695		
<b>21.5 Glycogen Breakdown and Synthesis Are Reciprocally Regulated</b>	<b>696</b>		
Protein phosphatase 1 reverses the regulatory effects of kinases on glycogen metabolism	696		
Insulin stimulates glycogen synthesis by inactivating glycogen synthase kinase	698		
Glycogen metabolism in the liver regulates the blood-glucose concentration	698		
		APPENDIX	
		<b>Biochemistry in Focus</b>	<b>702</b>
		APPENDIX	
		<b>Problem-Solving Strategies</b>	<b>703</b>
		CHAPTER 22	
		<b>Fatty Acid Metabolism</b>	<b>709</b>
		Fatty acid degradation and synthesis mirror each other in their chemical reactions	710
		<b>22.1 Triacylglycerols Are Highly Concentrated Energy Stores</b>	<b>710</b>
		Dietary lipids are digested by pancreatic lipases	712
		Dietary lipids are transported in chylomicrons	713
		<b>22.2 The Use of Fatty Acids as Fuel Requires Three Stages of Processing</b>	<b>713</b>
		Triacylglycerols are hydrolyzed by hormone-stimulated lipases	713
		Free fatty acids and glycerol are released into the blood	714
		Fatty acids are linked to coenzyme A before they are oxidized	715
		Carnitine carries long-chain activated fatty acids into the mitochondrial matrix	716
		Acetyl CoA, NADH, and FADH <sub>2</sub> are generated in each round of fatty acid oxidation	717
		The complete oxidation of palmitate yields 106 molecules of ATP	718
		<b>22.3 Unsaturated and Odd-Chain Fatty Acids Require Additional Steps for Degradation</b>	<b>719</b>
		An isomerase and a reductase are required for the oxidation of unsaturated fatty acids	719
		Odd-chain fatty acids yield propionyl CoA in the final thiolysis step	720
		Vitamin B <sub>12</sub> contains a corrin ring and a cobalt atom	721
		Mechanism: Methylmalonyl CoA mutase catalyzes a rearrangement to form succinyl CoA	721
		Fatty acids are also oxidized in peroxisomes	723
		Some fatty acids may contribute to the development of pathological conditions	724
		<b>22.4 Ketone Bodies Are a Fuel Source Derived from Fats</b>	<b>724</b>
		Ketone bodies are a major fuel in some tissues	725
		Animals cannot convert fatty acids into glucose	727
		<b>22.5 Fatty Acids Are Synthesized by Fatty Acid Synthase</b>	<b>728</b>
		Fatty acids are synthesized and degraded by different pathways	728
		The formation of malonyl CoA is the committed step in fatty acid synthesis	729
		Intermediates in fatty acid synthesis are attached to an acyl carrier protein	729

Fatty acid synthesis consists of a series of condensation, reduction, dehydration, and reduction reactions	729	Mechanism: Pyridoxal phosphate forms Schiff-base intermediates in aminotransferases	759
Fatty acids are synthesized by a multifunctional enzyme complex in animals	731	Aspartate aminotransferase is an archetypal pyridoxal-dependent transaminase	761
The synthesis of palmitate requires 8 molecules of acetyl CoA, 14 molecules of NADPH, and 7 molecules of ATP	733	Blood levels of aminotransferases serve a diagnostic function	761
Citrate carries acetyl groups from mitochondria to the cytoplasm for fatty acid synthesis	733	Pyridoxal phosphate enzymes catalyze a wide array of reactions	762
Several sources supply NADPH for fatty acid synthesis	734	Serine and threonine can be directly deaminated	763
Fatty acid metabolism is altered in tumor cells	735	Peripheral tissues transport nitrogen to the liver	763
Triacylglycerols may become an important renewal energy source	735	<b>23.4 Ammonium Ion Is Converted into Urea in Most Terrestrial Vertebrates</b>	<b>764</b>
<b>22.6 The Elongation and Unsaturation of Fatty Acids Are Accomplished by Accessory Enzyme Systems</b>	<b>736</b>	The urea cycle begins with the formation of carbamoyl phosphate	764
Membrane-bound enzymes generate unsaturated fatty acids	736	Carbamoyl phosphate synthetase is the key regulatory enzyme for urea synthesis	764
Eicosanoid hormones are derived from polyunsaturated fatty acids	737	Carbamoyl phosphate reacts with ornithine to begin the urea cycle	765
Variations on a theme: Polyketide and nonribosomal peptide synthetases resemble fatty acid synthase	738	The urea cycle is linked to gluconeogenesis	766
<b>22.7 Acetyl CoA Carboxylase Plays a Key Role in Controlling Fatty Acid Metabolism</b>	<b>739</b>	Urea-cycle enzymes are evolutionarily related to enzymes in other metabolic pathways	767
Acetyl CoA carboxylase is regulated by conditions in the cell	739	Inherited defects of the urea cycle cause hyperammonemia and can lead to brain damage	767
Acetyl CoA carboxylase is regulated by a variety of hormones	740	Urea is not the only means of disposing of excess nitrogen	768
AMP-activated protein kinase is a key regulator of metabolism	740	<b>23.5 Carbon Atoms of Degraded Amino Acids Emerge as Major Metabolic Intermediates</b>	<b>769</b>
APPENDIX		Pyruvate is an entry point into metabolism for a number of amino acids	770
<b>Biochemistry in Focus</b>	<b>743</b>	Oxaloacetate is an entry point into metabolism for aspartate and asparagine	770
APPENDIX		Alpha-ketoglutarate is an entry point into metabolism for five-carbon amino acids	770
<b>Problem-Solving Strategies</b>	<b>744</b>	Succinyl coenzyme A is a point of entry for several amino acids	771
CHAPTER 23		Methionine degradation requires the formation of a key methyl donor, S-adenosylmethionine	772
<b>Protein Turnover and Amino Acid Catabolism</b>	<b>751</b>	Threonine deaminase initiates the degradation of threonine	772
<b>23.1 Proteins Are Degraded to Amino Acids</b>	<b>752</b>	The branched-chain amino acids yield acetyl CoA, acetoacetate, or propionyl CoA	772
The digestion of dietary proteins begins in the stomach and is completed in the intestine	752	Oxygenases are required for the degradation of aromatic amino acids	773
Cellular proteins are degraded at different rates	753	Protein metabolism helps to power the flight of migratory birds	775
<b>23.2 Protein Turnover Is Tightly Regulated</b>	<b>753</b>	<b>23.6 Inborn Errors of Metabolism Can Disrupt Amino Acid Degradation</b>	<b>776</b>
Ubiquitin tags proteins for destruction	754	Phenylketonuria is one of the most common metabolic disorders	777
The proteasome digests the ubiquitin-tagged proteins	755	Determining the basis of the neurological symptoms of phenylketonuria is an active area of research	777
The ubiquitin pathway and the proteasome have prokaryotic counterparts	756	APPENDIX	
Protein degradation can be used to regulate biological function	757	<b>Biochemistry in Focus</b>	<b>779</b>
<b>23.3 The First Step in Amino Acid Degradation Is the Removal of Nitrogen</b>	<b>758</b>	APPENDIX	
Alpha-amino groups are converted into ammonium ions by the oxidative deamination of glutamate	758	<b>Problem-Solving Strategies</b>	<b>780</b>

CHAPTER 24		APPENDIX	
<b>The Biosynthesis of Amino Acids</b>	<b>787</b>	<b>Biochemistry in Focus</b>	<b>815</b>
Amino acid synthesis requires solutions to three key biochemical problems	788	APPENDIX	
		<b>Problem-Solving Strategies</b>	<b>816</b>
<b>24.1 Nitrogen Fixation: Microorganisms Use ATP and a Powerful Reductant to Reduce Atmospheric Nitrogen to Ammonia</b>	<b>788</b>	CHAPTER 25	
The iron–molybdenum cofactor of nitrogenase binds and reduces atmospheric nitrogen	790	<b>Nucleotide Biosynthesis</b>	<b>821</b>
Ammonium ion is assimilated into an amino acid through glutamate and glutamine	791	Nucleotides can be synthesized by de novo or salvage pathways	822
<b>24.2 Amino Acids Are Made from Intermediates of the Citric Acid Cycle and Other Major Pathways</b>	<b>793</b>	<b>25.1 The Pyrimidine Ring Is Assembled de Novo or Recovered by Salvage Pathways</b>	<b>822</b>
Human beings can synthesize some amino acids but must obtain others from their diet	793	Bicarbonate and other oxygenated carbon compounds are activated by phosphorylation	823
Aspartate, alanine, and glutamate are formed by the addition of an amino group to an alpha-ketoacid	794	The side chain of glutamine can be hydrolyzed to generate ammonia	823
A common step determines the chirality of all amino acids	795	Intermediates can move between active sites by channeling	824
The formation of asparagine from aspartate requires an adenylated intermediate	796	Orotate acquires a ribose ring from PRPP to form a pyrimidine nucleotide and is converted into uridylylate	824
Glutamate is the precursor of glutamine, proline, and arginine	796	Nucleotide mono-, di-, and triphosphates are interconvertible	825
3-Phosphoglycerate is the precursor of serine, cysteine, and glycine	797	CTP is formed by amination of UTP	825
Tetrahydrofolate carries activated one-carbon units at several oxidation levels	798	Salvage pathways recycle pyrimidine bases	826
S-Adenosylmethionine is the major donor of methyl groups	799	<b>25.2 Purine Bases Can Be Synthesized de Novo or Recycled by Salvage Pathways</b>	<b>826</b>
Cysteine is synthesized from serine and homocysteine	801	The purine ring system is assembled on ribose phosphate	827
High homocysteine levels correlate with vascular disease	801	The purine ring is assembled by successive steps of activation by phosphorylation followed by displacement	827
Shikimate and chorismate are intermediates in the biosynthesis of aromatic amino acids	801	AMP and GMP are formed from IMP	829
Tryptophan synthase illustrates substrate channeling in enzymatic catalysis	804	Enzymes of the purine synthesis pathway associate with one another in vivo	830
		Salvage pathways economize intracellular energy expenditure	830
<b>24.3 Feedback Inhibition Regulates Amino Acid Biosynthesis</b>	<b>805</b>	<b>25.3 Deoxyribonucleotides Are Synthesized by the Reduction of Ribonucleotides Through a Radical Mechanism</b>	<b>831</b>
Branched pathways require sophisticated regulation	805	Mechanism: A tyrosyl radical is critical to the action of ribonucleotide reductase	831
The sensitivity of glutamine synthetase to allosteric regulation is altered by covalent modification	807	Stable radicals other than tyrosyl radical are employed by other ribonucleotide reductases	833
<b>24.4 Amino Acids Are Precursors of Many Biomolecules</b>	<b>808</b>	Thymidylate is formed by the methylation of deoxyuridylylate	834
Glutathione, a gamma-glutamyl peptide, serves as a sulfhydryl buffer and an antioxidant	809	Dihydrofolate reductase catalyzes the regeneration of tetrahydrofolate, a one-carbon carrier	835
Nitric oxide, a short-lived signal molecule, is formed from arginine	809	Several valuable anticancer drugs block the synthesis of thymidylate	835
Amino acids are precursors for a number of neurotransmitters	810	<b>25.4 Key Steps in Nucleotide Biosynthesis Are Regulated by Feedback Inhibition</b>	<b>836</b>
Porphyrins are synthesized from glycine and succinyl coenzyme A	810	Pyrimidine biosynthesis is regulated by aspartate transcarbamoylase	836
Porphyrins accumulate in some inherited disorders of porphyrin metabolism	814		

The synthesis of purine nucleotides is controlled by feedback inhibition at several sites	837		
The synthesis of deoxyribonucleotides is controlled by the regulation of ribonucleotide reductase	837		
<b>25.5 Disruptions in Nucleotide Metabolism Can Cause Pathological Conditions</b>	<b>838</b>		
The loss of adenosine deaminase activity results in severe combined immunodeficiency	839		
Gout is induced by high serum levels of urate	839		
Lesch–Nyhan syndrome is a dramatic consequence of mutations in a salvage-pathway enzyme	840		
Folic acid deficiency promotes birth defects such as spina bifida	841		
APPENDIX			
<b>Biochemistry in Focus</b>	<b>842</b>		
APPENDIX			
<b>Problem-Solving Strategies</b>	<b>843</b>		
CHAPTER 26			
<b>The Biosynthesis of Membrane Lipids and Steroids</b>	<b>849</b>		
<b>26.1 Phosphatidate Is a Common Intermediate in the Synthesis of Phospholipids and Triacylglycerols</b>	<b>850</b>		
The synthesis of phospholipids requires an activated intermediate	851		
Some phospholipids are synthesized from an activated alcohol	852		
Phosphatidylcholine is an abundant phospholipid	853		
Excess choline is implicated in the development of heart disease	853		
Base-exchange reactions can generate phospholipids	853		
Sphingolipids are synthesized from ceramide	854		
Gangliosides are carbohydrate-rich sphingolipids that contain acidic sugars	855		
Sphingolipids confer diversity on lipid structure and function	856		
Respiratory distress syndrome and Tay–Sachs disease result from the disruption of lipid metabolism	856		
Ceramide metabolism stimulates tumor growth	857		
Phosphatidic acid phosphatase is a key regulatory enzyme in lipid metabolism	857		
<b>26.2 Cholesterol Is Synthesized from Acetyl Coenzyme A in Three Stages</b>	<b>858</b>		
The synthesis of mevalonate, which is activated as isopentenyl pyrophosphate, initiates the synthesis of cholesterol	858		
Squalene (C <sub>30</sub> ) is synthesized from six molecules of isopentenyl pyrophosphate (C <sub>5</sub> )	859		
Squalene cyclizes to form cholesterol	860		
<b>26.3 The Complex Regulation of Cholesterol Biosynthesis Takes Place at Several Levels</b>	<b>861</b>		
Lipoproteins transport cholesterol and triacylglycerols throughout the organism	863		
Low-density lipoproteins play a central role in cholesterol metabolism	866		
The absence of the LDL receptor leads to hypercholesterolemia and atherosclerosis	867		
Mutations in the LDL receptor prevent LDL release and result in receptor destruction	868		
Inability to transport cholesterol from the lysosome causes Niemann–Pick disease	869		
Cycling of the LDL receptor is regulated	869		
HDL appears to protect against atherosclerosis	870		
The clinical management of cholesterol levels can be understood at a biochemical level	870		
<b>26.4 Important Biochemicals Are Synthesized from Cholesterol and Isoprene</b>	<b>871</b>		
Letters identify the steroid rings and numbers identify the carbon atoms	872		
Steroids are hydroxylated by cytochrome P450 monooxygenases that use NADPH and O <sub>2</sub>	873		
The cytochrome P450 system is widespread and performs a protective function	874		
Pregnenolone, a precursor of many other steroids, is formed from cholesterol by cleavage of its side chain	875		
Progesterone and corticosteroids are synthesized from pregnenolone	875		
Androgens and estrogens are synthesized from pregnenolone	875		
Vitamin D is derived from cholesterol by the ring-splitting activity of light	877		
Five-carbon units are joined to form a wide variety of biomolecules	878		
Some isoprenoids have industrial applications	880		
APPENDIX			
<b>Biochemistry in Focus</b>	<b>882</b>		
APPENDIX			
<b>Problem-Solving Strategies</b>	<b>882</b>		
CHAPTER 27			
<b>The Integration of Metabolism</b>	<b>889</b>		
<b>27.1 Caloric Homeostasis Is a Means of Regulating Body Weight</b>	<b>890</b>		
<b>27.2 The Brain Plays a Key Role in Caloric Homeostasis</b>	<b>892</b>		
Signals from the gastrointestinal tract induce feelings of satiety	892		
Leptin and insulin regulate long-term control over caloric homeostasis	893		
Leptin is one of several hormones secreted by adipose tissue	894		

Leptin resistance may be a contributing factor to obesity	895	Screening libraries of synthetic compounds expands the opportunity for identification of drug leads	933
Dieting is used to combat obesity	896	Drugs can be designed on the basis of three-dimensional structural information about their targets	935
<b>27.3 Diabetes Is a Common Metabolic Disease Often Resulting from Obesity</b>	<b>896</b>	<b>28.3 Genomic Analyses Can Aid Drug Discovery</b>	<b>937</b>
Insulin initiates a complex signal-transduction pathway in muscle	897	Potential targets can be identified in the human proteome	937
Metabolic syndrome often precedes type 2 diabetes	898	Animal models can be developed to test the validity of potential drug targets	938
Excess fatty acids in muscle modify metabolism	899	Potential targets can be identified in the genomes of pathogens	939
Insulin resistance in muscle facilitates pancreatic failure	899	Genetic differences influence individual responses to drugs	939
Metabolic derangements in type 1 diabetes result from insulin insufficiency and glucagon excess	901	<b>28.4 The Clinical Development of Drugs Proceeds Through Several Phases</b>	<b>940</b>
<b>27.4 Exercise Beneficially Alters the Biochemistry of Cells</b>	<b>901</b>	Clinical trials are time-consuming and expensive	941
Mitochondrial biogenesis is stimulated by muscular activity	902	The evolution of drug resistance can limit the utility of drugs for infectious agents and cancer	942
Fuel choice during exercise is determined by the intensity and duration of activity	902		
<b>27.5 Food Intake and Starvation Induce Metabolic Changes</b>	<b>905</b>	APPENDIX	
The starved-fed cycle is the physiological response to a fast	905	<b>Biochemistry in Focus</b>	<b>945</b>
Metabolic adaptations in prolonged starvation minimize protein degradation	907	CHAPTER 29	
<b>27.6 Ethanol Alters Energy Metabolism in the Liver</b>	<b>909</b>	<b>DNA Replication, Repair, and Recombination</b>	<b>949</b>
Ethanol metabolism leads to an excess of NADH	910	<b>29.1 DNA Replication Proceeds by the Polymerization of Deoxyribonucleoside Triphosphates Along a Template</b>	<b>951</b>
Excess ethanol consumption disrupts vitamin metabolism	911	DNA polymerases require a template and a primer	951
		All DNA polymerases have structural features in common	951
APPENDIX		Two bound metal ions participate in the polymerase reaction	952
<b>BIOCHEMISTRY IN FOCUS</b>	<b>914</b>	The specificity of replication is dictated by complementarity of shape between bases	952
<b>Biochemistry in Focus 1</b>	<b>914</b>	An RNA primer synthesized by primase enables DNA synthesis to begin	953
<b>Biochemistry in Focus 2</b>	<b>915</b>	One strand of DNA is made continuously, whereas the other strand is synthesized in fragments	953
APPENDIX		DNA ligase joins ends of DNA in duplex regions	954
<b>Problem-Solving Strategies</b>	<b>916</b>	The separation of DNA strands requires specific helicases and ATP hydrolysis	955
CHAPTER 28		<b>29.2 DNA Unwinding and Supercoiling Are Controlled by Topoisomerases</b>	<b>956</b>
<b>Drug Development</b>	<b>921</b>	The linking number of DNA, a topological property, determines the degree of supercoiling	956
<b>28.1 Compounds Must Meet Stringent Criteria to Be Developed into Drugs</b>	<b>923</b>	Topoisomerases prepare the double helix for unwinding	958
Drugs must be potent and selective	923	Type I topoisomerases relax supercoiled structures	959
Drugs must have suitable properties to reach their targets	924	Type II topoisomerases can introduce negative supercoils through coupling to ATP hydrolysis	960
Toxicity can limit drug effectiveness	929	<b>29.3 DNA Replication Is Highly Coordinated</b>	<b>961</b>
<b>28.2 Drug Candidates Can Be Discovered by Serendipity, Screening, or Design</b>	<b>930</b>	DNA replication requires highly processive polymerases	962
Serendipitous observations can drive drug development	930		
Natural products are a valuable source of drugs and drug leads	932		



The leading and lagging strands are synthesized in a coordinated fashion	962	Some messenger RNAs directly sense metabolite concentrations	992
DNA replication in <i>Escherichia coli</i> begins at a unique site and proceeds through initiation, elongation, and termination	964	The <i>rho</i> protein helps to terminate the transcription of some genes	992
DNA synthesis in eukaryotes is initiated at multiple sites	965	Some antibiotics inhibit transcription	994
Telomeres are unique structures at the ends of linear chromosomes	967	Precursors of transfer and ribosomal RNA are cleaved and chemically modified after transcription in prokaryotes	995
Telomeres are replicated by telomerase, a specialized polymerase that carries its own RNA template	968	<b>30.2 Transcription in Eukaryotes Is Highly Regulated</b>	<b>996</b>
<b>29.4 Many Types of DNA Damage Can Be Repaired</b>	<b>968</b>	Three types of RNA polymerase synthesize RNA in eukaryotic cells	997
Errors can arise in DNA replication	969	Three common elements can be found in the RNA polymerase II promoter region	998
Bases can be damaged by oxidizing agents, alkylating agents, and light	969	The TFIID protein complex initiates the assembly of the active transcription complex	999
DNA damage can be detected and repaired by a variety of systems	970	Multiple transcription factors interact with eukaryotic promoters	1000
The presence of thymine instead of uracil in DNA permits the repair of deaminated cytosine	972	Enhancer sequences can stimulate transcription at start sites thousands of bases away	1001
Some genetic diseases are caused by the expansion of repeats of three nucleotides	973	<b>30.3 The Transcription Products of Eukaryotic Polymerases Are Processed</b>	<b>1001</b>
Many cancers are caused by the defective repair of DNA	973	RNA polymerase I produces three ribosomal RNAs	1001
Many potential carcinogens can be detected by their mutagenic action on bacteria	975	RNA polymerase III produces transfer RNA	1002
<b>29.5 DNA Recombination Plays Important Roles in Replication, Repair, and Other Processes</b>	<b>975</b>	The product of RNA polymerase II, the pre-mRNA transcript, acquires a 5' cap and a 3' poly(A) tail	1002
RecA can initiate recombination by promoting strand invasion	976	Small regulatory RNAs are cleaved from larger precursors	1004
Some recombination reactions proceed through Holliday-junction intermediates	976	RNA editing changes the proteins encoded by mRNA	1004
		Sequences at the ends of introns specify splice sites in mRNA precursors	1005
		Splicing consists of two sequential transesterification reactions	1006
APPENDIX		Small nuclear RNAs in spliceosomes catalyze the splicing of mRNA precursors	1007
<b>Biochemistry in Focus</b>	<b>980</b>	Transcription and processing of mRNA are coupled	1010
		Mutations that affect pre-mRNA splicing cause disease	1010
		Most human pre-mRNAs can be spliced in alternative ways to yield different proteins	1011
CHAPTER 30		<b>30.4 The Discovery of Catalytic RNA Was Revealing in Regard to Both Mechanism and Evolution</b>	<b>1012</b>
<b>RNA Synthesis and Processing</b>	<b>983</b>		
RNA synthesis comprises three stages: Initiation, elongation, and termination	984	APPENDIX	
<b>30.1 RNA Polymerases Catalyze Transcription</b>	<b>985</b>	<b>Biochemistry in Focus</b>	<b>1017</b>
RNA chains are formed de novo and grow in the 5'-to-3' direction	986		
RNA polymerases backtrack and correct errors	988	CHAPTER 31	
RNA polymerase binds to promoter sites on the DNA template to initiate transcription	988	<b>Protein Synthesis</b>	<b>1021</b>
Sigma subunits of RNA polymerase recognize promoter sites	989		
RNA polymerases must unwind the template double helix for transcription to take place	990	<b>31.1 Protein Synthesis Requires the Translation of Nucleotide Sequences into Amino Acid Sequences</b>	<b>1022</b>
Elongation takes place at transcription bubbles that move along the DNA template	991	The synthesis of long proteins requires a low error frequency	1022
Sequences within the newly transcribed RNA signal termination	991		

Transfer RNA molecules have a common design	1023	APPENDIX	
Some transfer RNA molecules recognize more than one codon because of wobble in base-pairing	1025	<b>Problem-Solving Strategies</b>	<b>1051</b>
<b>31.2 Aminoacyl Transfer RNA Synthetases Read the Genetic Code</b>	<b>1026</b>	CHAPTER 32	
Amino acids are first activated by adenylation	1027	<b>The Control of Gene Expression in Prokaryotes</b>	<b>1057</b>
Aminoacyl-tRNA synthetases have highly discriminating amino acid activation sites	1028	<b>32.1 Many DNA-Binding Proteins Recognize Specific DNA Sequences</b>	<b>1058</b>
Proofreading by aminoacyl-tRNA synthetases increases the fidelity of protein synthesis	1029	The helix-turn-helix motif is common to many prokaryotic DNA-binding proteins	1059
Synthetases recognize various features of transfer RNA molecules	1029	<b>32.2 Prokaryotic DNA-Binding Proteins Bind Specifically to Regulatory Sites in Operons</b>	<b>1060</b>
Aminoacyl-tRNA synthetases can be divided into two classes	1030	An operon consists of regulatory elements and protein-encoding genes	1061
<b>31.3 The Ribosome Is the Site of Protein Synthesis</b>	<b>1031</b>	The <i>lac</i> repressor protein in the absence of lactose binds to the operator and blocks transcription	1061
Ribosomal RNAs (5S, 16S, and 23S rRNA) play a central role in protein synthesis	1031	Ligand binding can induce structural changes in regulatory proteins	1062
Ribosomes have three tRNA-binding sites that bridge the 30S and 50S subunits	1032	The operon is a common regulatory unit in prokaryotes	1063
The start signal is usually AUG preceded by several bases that pair with 16S rRNA	1034	Transcription can be stimulated by proteins that contact RNA polymerase	1064
Bacterial protein synthesis is initiated by formylmethionyl transfer RNA	1035	<b>32.3 Regulatory Circuits Can Result in Switching Between Patterns of Gene Expression</b>	<b>1065</b>
Formylmethionyl-tRNA <sub>f</sub> is placed in the P site of the ribosome in the formation of the 70S initiation complex	1035	The $\lambda$ repressor regulates its own expression	1065
Elongation factors deliver aminoacyl-tRNA to the ribosome	1036	A circuit based on the $\lambda$ repressor and Cro forms a genetic switch	1066
Peptidyl transferase catalyzes peptide-bond synthesis	1037	Many prokaryotic cells release chemical signals that regulate gene expression in other cells	1066
The formation of a peptide bond is followed by the GTP-driven translocation of tRNAs and mRNA	1038	Biofilms are complex communities of prokaryotes	1067
Protein synthesis is terminated by release factors that read stop codons	1040	<b>32.4 Gene Expression Can Be Controlled at Posttranscriptional Levels</b>	<b>1068</b>
<b>31.4 Eukaryotic Protein Synthesis Differs from Bacterial Protein Synthesis Primarily in Translation Initiation</b>	<b>1041</b>	Attenuation is a prokaryotic mechanism for regulating transcription through the modulation of nascent RNA secondary structure	1068
Mutations in initiation factor 2 cause a curious pathological condition	1042	APPENDIX	
<b>31.5 A Variety of Antibiotics and Toxins Can Inhibit Protein Synthesis</b>	<b>1043</b>	<b>Biochemistry in Focus</b>	<b>1072</b>
Some antibiotics inhibit protein synthesis	1043	CHAPTER 33	
Diphtheria toxin blocks protein synthesis in eukaryotes by inhibiting translocation	1044	<b>The Control of Gene Expression in Eukaryotes</b>	<b>1075</b>
Some toxins modify 28S ribosomal RNA	1045	<b>33.1 Eukaryotic DNA Is Organized into Chromatin</b>	<b>1077</b>
<b>31.6 Ribosomes Bound to the Endoplasmic Reticulum Manufacture Secretory and Membrane Proteins</b>	<b>1045</b>	Nucleosomes are complexes of DNA and histones	1077
Protein synthesis begins on ribosomes that are free in the cytoplasm	1046	DNA wraps around histone octamers to form nucleosomes	1078
Signal sequences mark proteins for translocation across the endoplasmic reticulum membrane	1046	<b>33.2 Transcription Factors Bind DNA and Regulate Transcription Initiation</b>	<b>1079</b>
Transport vesicles carry cargo proteins to their final destination	1048	A range of DNA-binding structures are employed by eukaryotic DNA-binding proteins	1080
APPENDIX		Activation domains interact with other proteins	1081
<b>Biochemistry in Focus</b>	<b>1051</b>	Multiple transcription factors interact with eukaryotic regulatory regions	1081

Enhancers can stimulate transcription in specific cell types	1082	Light absorption induces a specific isomerization of bound 11- <i>cis</i> -retinal	1108
Induced pluripotent stem cells can be generated by introducing four transcription factors into differentiated cells	1082	Light-induced lowering of the calcium level coordinates recovery	1109
<b>33.3 The Control of Gene Expression Can Require Chromatin Remodeling</b>	<b>1083</b>	Color vision is mediated by three cone receptors that are homologs of rhodopsin	1110
The methylation of DNA can alter patterns of gene expression	1084	Rearrangements in the genes for the green and red pigments lead to “color blindness”	1111
Steroids and related hydrophobic molecules pass through membranes and bind to DNA-binding receptors	1084	<b>34.4 Hearing Depends on the Speedy Detection of Mechanical Stimuli</b>	<b>1111</b>
Nuclear hormone receptors regulate transcription by recruiting coactivators to the transcription complex	1085	Hair cells use a connected bundle of stereocilia to detect tiny motions	1112
Steroid-hormone receptors are targets for drugs	1086	Mechanosensory channels have been identified in <i>Drosophila</i> and vertebrates	1113
Chromatin structure is modulated through covalent modifications of histone tails	1087	<b>34.5 Touch Includes the Sensing of Pressure, Temperature, and Other Factors</b>	<b>1113</b>
Transcriptional repression can be achieved through histone deacetylation and other modifications	1089	Studies of capsaicin reveal a receptor for sensing high temperatures and other painful stimuli	1114
<b>33.4 Eukaryotic Gene Expression Can Be Controlled at Posttranscriptional Levels</b>	<b>1090</b>	APPENDIX	
Genes associated with iron metabolism are translationally regulated in animals	1090	<b>Biochemistry in Focus</b>	<b>1116</b>
Small RNAs regulate the expression of many eukaryotic genes	1092	☉ CHAPTER 35	
APPENDIX		<b>The Immune System</b>	<b>1119</b>
<b>Biochemistry in Focus</b>	<b>1094</b>	Innate immunity is an evolutionarily ancient defense system	1120
☉ CHAPTER 34		The adaptive immune system responds by using the principles of evolution	1122
<b>Sensory Systems</b>	<b>1097</b>	<b>35.1 Antibodies Possess Distinct Antigen-Binding and Effector Units</b>	<b>1124</b>
<b>34.1 A Wide Variety of Organic Compounds Are Detected by Olfaction</b>	<b>1098</b>	<b>35.2 Antibodies Bind Specific Molecules Through Hypervariable Loops</b>	<b>1126</b>
Olfaction is mediated by an enormous family of seven-transmembrane-helix receptors	1099	The immunoglobulin fold consists of a beta-sandwich framework with hypervariable loops	1127
Odorants are decoded by a combinatorial mechanism	1101	X-ray analyses have revealed how antibodies bind antigens	1127
<b>34.2 Taste Is a Combination of Senses That Function by Different Mechanisms</b>	<b>1102</b>	Large antigens bind antibodies with numerous interactions	1128
Sequencing of the human genome led to the discovery of a large family of 7TM bitter receptors	1103	<b>35.3 Diversity Is Generated by Gene Rearrangements</b>	<b>1129</b>
A heterodimeric 7TM receptor responds to sweet compounds	1105	J (joining) genes and D (diversity) genes increase antibody diversity	1130
Umami, the taste of glutamate and aspartate, is mediated by a heterodimeric receptor related to the sweet receptor	1105	More than 10 <sup>8</sup> antibodies can be formed by combinatorial association and somatic mutation	1131
Salty tastes are detected primarily by the passage of sodium ions through channels	1105	The oligomerization of antibodies expressed on the surfaces of immature B cells triggers antibody secretion	1132
Sour tastes arise from the effects of hydrogen ions (acids) on channels	1106	Different classes of antibodies are formed by the hopping of V <sub>H</sub> genes	1133
<b>34.3 Photoreceptor Molecules in the Eye Detect Visible Light</b>	<b>1106</b>	<b>35.4 Major-Histocompatibility-Complex Proteins Present Peptide Antigens on Cell Surfaces for Recognition by T-Cell Receptors</b>	<b>1134</b>
Rhodopsin, a specialized 7TM receptor, absorbs visible light	1107	Peptides presented by MHC proteins occupy a deep groove flanked by alpha helices	1135
		☉ Chapters online only	

T-cell receptors are antibody-like proteins containing variable and constant regions	1136	<b>36.2 Myosins Move Along Actin Filaments</b>	1156
CD8 on cytotoxic T cells acts in concert with T-cell receptors	1137	Actin is a polar, self-assembling, dynamic polymer	1157
Helper T cells stimulate cells that display foreign peptides bound to class II MHC proteins	1139	Myosin head domains bind to actin filaments	1158
Helper T cells rely on the T-cell receptor and CD4 to recognize foreign peptides on antigen-presenting cells	1139	Motions of single motor proteins can be directly observed	1159
MHC proteins are highly diverse	1141	Phosphate release triggers the myosin power stroke	1159
Human immunodeficiency viruses subvert the immune system by destroying helper T cells	1141	Muscle is a complex of myosin and actin	1160
<b>35.5 The Immune System Contributes to the Prevention and the Development of Human Diseases</b>	1142	The length of the lever arm determines motor velocity	1162
T cells are subjected to positive and negative selection in the thymus	1142	<b>36.3 Kinesin and Dynein Move Along Microtubules</b>	1162
Autoimmune diseases result from the generation of immune responses against self-antigens	1143	Microtubules are hollow cylindrical polymers	1163
The immune system plays a role in cancer prevention	1144	Kinesin motion is highly processive	1164
Vaccines are a powerful means to prevent and eradicate disease	1144	<b>36.4 A Rotary Motor Drives Bacterial Motion</b>	1166
		Bacteria swim by rotating their flagella	1166
		Proton flow drives bacterial flagellar rotation	1167
		Bacterial chemotaxis depends on reversal of the direction of flagellar rotation	1168
		APPENDIX	
APPENDIX		<b>Biochemistry in Focus</b>	1171
<b>Biochemistry in Focus</b>	1147	Answers to Problems	A1
		Selected Readings	B1
		Index	C1
⊙ CHAPTER 36			
<b>Molecular Motors</b>	1151		
<b>36.1 Most Molecular-Motor Proteins Are Members of the P-Loop NTPase Superfamily</b>	1152		
Molecular motors are generally oligomeric proteins with an ATPase core and an extended structure	1153		
ATP binding and hydrolysis induce changes in the conformation and binding affinity of motor proteins	1155		

⊙ Chapters online only