

# Contents

**Preface**    xiii

**Introduction**    xv

**1. Drought Disaster in Asia**    1

*Mathew Barlow, Heidi Cullen, Brad Lyon, and Olga Wilhelmi*

**2. Global Landslides Risk Case Study**    21

*Farrokh Nadim, Oddvar Kjekstad, Ulrik Domaas, Ramez Rafat, and Pascal Peduzzi*

**3. Storm Surges in Coastal Areas**    79

*Robert J. Nicholls*

**4. Natural Disaster Risks in Sri Lanka: Mapping Hazards and Risk Hotspots**    109

*Lareef Zubair and Vidhura Ralapanawe, Upamala Tennakoon, Zeenas Yahiya, and Ruvini Perera*

**5. Multihazard Risks in Caracas, República Bolivariana de Venezuela**    137

*Kristina R. Czuchlewski, Klaus H. Jacob, Arthur L. Lerner-Lam, Kevin Vranes, and Students of the Urban Planning Studio: "Disaster Resilient Caracas"*

**6. Reducing the Impacts of Floods through Early Warning and Preparedness: A Pilot Study for Kenya**    165

*Hussein Gadain, Nicolas Bidault, Linda Stephen, Ben Watkins, Maxx Dilley, and Nancy Mutunga*

**Tables**

Table 2.1.	Description of variables	25
Table 2.2.	Classification of slope factor " $S_r$ " for evaluation of susceptibility	26
Table 2.3.	Classification of lithology factor " $S_l$ " for evaluation of susceptibility	27
Table 2.4.	Classification of soil moisture factor " $S_h$ " for evaluation of susceptibility	28
Table 2.5.	Classification of precipitation trigger indicator " $T_p$ "	28
Table 2.6.	Classification of seismicity trigger indicator " $T_s$ "	31
Table 2.7.	Classification of landslide hazard potential based on the computed hazard index originally suggested by Mora and Vahrson (1994)	31
Table 2.8.	Classification of landslide hazard potential based on the computed hazard index used in this study	31
Table 2.9.	Classification of slope factor " $S_r$ " for snow avalanche susceptibility	31
Table 2.10.	Classification of precipitation factor " $T_p$ " for avalanche hazard evaluation	38

Table 2.11.	Classification of temperature factor “ $T_1$ ” for avalanche hazard analysis	38
Table 2.12.	Classification of snow avalanche hazard potential	38
Table 2.13.	Annual frequency of occurrence and typical return period (in years) for different classes of landslide and avalanche hazard	39
Table 2.A.1.	Classes of frequencies	66
Table 2.A.2.	Vulnerability indicators	72
Table 2.A.3.	Exponent and p-value for landslide multiple regression	74
Table 2.A.4.	Other exponents and p-values for landslide multiple regression	75
Table 3.1.	Hurricane characteristics and indicative surge magnitudes based on the Saffir-Simpson scale	81
Table 3.2.	Some major coastal cities and human-induced subsidence during the 20th century	82
Table 3.3.	Generic approaches to hazard reduction based on purposeful adjustment.	85
Table 3.4.	Regional contributions to coastal flooding in 1990 and the 2020s based on the analysis of Nicholls (2004).	87
Table 3.5.	The range of scenarios used by Nicholls (2004)	88
Table 3.6.	Estimates of the global exposure and incidence of flooding under the four SRES scenarios in the 2080s, plus 1990 estimates as a reference	88
Table 3.7.	Global-mean sea-level rise scenarios (cm) used by Nicholls (2004) (referenced to 1990), including the IS92a GGal scenario as a reference	90
Table 3.8.	The SRES Socioeconomic Scenarios for the 2080s: A Global Summary	90
Table 3.9.	Deaths associated with major hurricanes, cyclones, and typhoons (MC) and extra-tropical storm (ETS) disasters (>1,000 deaths) since 1700.	92
Table 3.10.	Deaths in storm surges around the North Sea from the 11th to the 18th centuries. All surges were due to extra-tropical storms	93
Table 3.11.	An expert synthesis of storm surge hotspots around the world.	98
Table 3.12.	Potential and actual hotspots vulnerable to flooding by the storm surge.	99
Table 5.1.	Critical Facilities and Systems (Categories and Definitions)	142
Table 5.2.	Studio estimates for the order of magnitude of losses for a generic city whose assets are valued at US \$100 billion.	161
Table 6.1.	Flood scenarios for a worst case and a moderate case	177
Table 6.2.	Characteristics of the different livelihood zones analyzed	182
Table 6.3.	Percentage of livestock contribution to cash income and food consumption of the livelihood zones analyzed	182

## Figures

Figure 1.1.	Total Annual Precipitation, in millimeters.	3
Figure 1.2.	Total number of drought disasters for all Asian countries with geo-referenced boundaries available	4
Figure 1.3.	Number of drought disasters with month specified, for all countries listed in the Asia category in EM-DAT	5
Figure 1.4.	Number of drought disasters for Asia and the maritime continent, summed by year and over all countries in the region	5
Figure 1.5.	Number of drought disasters with months specified for Asia and the maritime continent	6

Figure 1.6.	Number of drought disasters for non-Asia countries in the EM-DAT database	6
Figure 1.7.	Precipitation anomalies for the 1999–2001 period, divided by yearly standard deviation to facilitate comparison over diverse climate regimes	7
Figure 1.8.	Reported drought disasters, 1999–2001	8
Figure 1.9.	Match between drought disaster and climatic measure of drought (3 consecutive months with precipitation deficits meeting a set threshold).	10
Figure 1.10.	Match between drought disaster and climatic measure of drought (4 out of 6 months with precipitation deficits meeting a set threshold).	10
Figure 1.11.	Match between drought disaster and climatic measure of drought (12-month average of Weighted Anomaly of Standardized Precipitation (WASP)).	11
Figure 1.12.	Number of matches for 12-month WASP compared to the total number of drought disaster reports (with monthly data).	11
Figure 1.13.	Correlation between the 12-month WASP calculated from two different precipitation data sets: the University of East Anglia (UEA) precipitation data and the CPC's Merged Analysis of Precipitation (CMAP).	13
Figure 1.14.	Time series of drought disasters and climatic drought events (based on 12-month WASP)	13
Figure 1.15.	Climate anomalies (12-month WASP) for two periods: 1982–1983 (red) and 1999–2000 (blue)	14
Figure 1.16.	WASP estimate of climatic drought (shaded brown curve) and drought disaster declarations (red bars) for Central-Southwest Asia countries.	15
Figure 1.17.	WASP estimate of climatic drought (shaded brown curve) and drought disaster declarations (red bars) for Laos and India.	16
Figure 1.A.1.	Persistent deficit of precipitation	18
Figure 2.1.	General approach for landslide hazard and risk evaluation	23
Figure 2.2.	Global soil moisture index: 1961–1990	29
Figure 2.3.	Expected monthly extreme values for a 100-years event.	30
Figure 2.4.	Expected PGA with a return period of 475 years	32
Figure 2.5.	Variation of slope factor, $S_r$ , in Tajikistan and its neighboring regions	33
Figure 2.6.	Variation of lithology factor, $S_l$ , in Tajikistan and its neighboring regions	34
Figure 2.7.	Variation of seismic trigger indicator, $T_s$ , in Tajikistan and its neighboring regions	35
Figure 2.8.	Variation of soil moisture factor, $S_h$ , in Tajikistan and its neighboring regions	36
Figure 2.9.	Landslide hazard zonation map obtained for Tajikistan and its neighboring regions	37
Figure 2.10.	Example landslide hazard map for Central American and Caribbean countries	40
Figure 2.11.	Example landslide risk map for parts of Central and South America	41
Figure 2.12.	Historical rock avalanche events in Møre & Romsdal and Sogn & Fjordane Counties extracted from Norway's historical database (NGU/Astor Furseth)	42
Figure 2.13.	Regional hazard zonation in Møre & Romsdal County in western Norway.	43
Figure 2.14.	Landslide hazard map (landslide and rock fall hazards) for the western part of Norway based on the simplified model	44
Figure 2.15.	Snow avalanche hazard zones for Norway based on the avalanche hazard model	45
Figure 2.16.	Map of Armenia	47
Figure 2.17.	Comparison of global landslide hazard mapping in Armenia using NGI model with the GEORISK landslide inventory	48
Figure 2.18.	Major landslide events in Nepal during a 30-year time period (1971–2000)	49
Figure 2.19.	Landslide hazard in Nepal predicted by the NGI model in this study	49

- Figure 2.20. Population density map of Nepal in 1995. 50
- Figure 2.21. Landslide hazard in Georgia predicted by the model developed in this study 52
- Figure 2.22. Snow avalanche hazard in Georgia predicted by the model developed in this study 53
- Figure 2.23. Observed landslides in Sri Lanka between 1947 and 2003 and prediction of landslide hazard in Sri Lanka by the model developed in this study. 54
- Figure 2.24. Historical landslide data in Jamaica 55
- Figure 2.25. Prediction of landslide hazard in Jamaica with the model developed in this study 55
- Figure 2.26. Global hotspot landslide hazard zonation for the world 56
- Figure 2.27. Global hotspot landslide hazard zonation for Central Asia and the Middle East 58
- Figure 2.28. Global hotspot landslide hazard zonation for Central American and Caribbean countries 59
- Figure 2.29. Hotspot landslide risk zonation for Central America and Jamaica 61
- Figure 2.30. Hotspot landslide risk zonation for Central Asia 62
- Figure 2.31. Global hotspot snow avalanche hazard zonation for Central Asia 63
- Figure 2.A.1. Distribution of risk utilizing a vulnerability proxy in Central America 69
- Figure 2.A.2. Distribution of risk using a vulnerability proxy in South America 70
- Figure 2.A.3. Distribution of risk utilizing a vulnerability proxy in Central Asia 71
- Figure 2.A.4. Transformation for variables ranging between 0 and 1 73
- Figure 2.A.5. Predicted killed versus observed for landslide 74
- 
- Figure 3.1. Areas in the southwest Netherlands flooded by the 1953 storm surge, February 1, 1953 (from Edwards 1953) 81
- Figure 3.2. A simplified reconstruction of the November 1970 storm surge in Bangladesh. 82
- Figure 3.3. Areas in Tokyo that are below normal high-water and surge levels with and without a 1-m rise in sea level. 83
- Figure 3.4. Coasts affected by tropical cyclones 84
- Figure 3.5. People at risk (that is, people potentially flooded) versus people in the flood hazard zone in 1990 for 20 global regions. 88
- Figure 3.6. People at risk (that is, people potentially flooded) versus people in the flood hazard zone in the 2080s for 20 global regions. 91
- Figure 3.7. Deaths by major hurricanes, cyclones, and typhoons (MC) and extra-tropical storms (ETS) from 1700 to 2000. 94
- Figure 3.8. Number of “significant” events based on two thresholds of deaths: > 50,000 deaths, and > 20,000 deaths, as well as all events (>1,000 deaths) 94
- Figure 3.9. Annual deaths due to surges, averaged over 50-year periods using the data in Table 3.8 95
- Figure 3.10. Deaths per event for hurricanes making landfall in the United States 96
- Figure 3.11. Flooding of the East Coast of England during the 1953 storm surge 100
- Figure 3.12. Deaths per surge event in Bangladesh from 1800 to 2000 using the data in Table 3.9. 102
- Figure 3.13. Surges on the U.S. Gulf and East Coast. Relative storm-surge potential (a), and surge graphs for six Atlantic coast locations (b), the hurricane of September 14–15, 1944. 103
- Figure 3.14. Subsidence from the 1920s to the 1990s in Shanghai, China 104
- Figure 4.1. The district boundaries of Sri Lanka are shown over the topography 110
- Figure 4.2.a. The average annual rainfall climatology estimated based on data from 284 stations in the period between 1960 and 1990. Homogenous climatological regions as proposed by Puvaneswaran and Smithson (1993) are overlaid. 111

Figure 4.2.b.	The average monthly rainfall between 1869 and 1998 for Sri Lanka	112
Figure 4.3.	The density of population in each of the 323 Divisional Secretariat Divisions based on data from the census of 2001	114
Figure 4.4.	The food insecurity index of Divisional Secretariat Divisions (DSDs) as estimated by the World Food Programme	115
Figure 4.5.a.	Sectoral breakdown of the GDP for 2001	117
Figure 4.5.b.	Sectoral breakdown of the labor force for 2001	117
Figure 4.6.	The gross domestic product (GDP) by province for 1995	118
Figure 4.7.	The estimate of industrial output in the districts in 1995	119
Figure 4.8.	Infrastructure density index estimated for each district, as described in the text	121
Figure 4.9.	The drought hazard was estimated using a modified WASP index.	122
Figure 4.10.	Drought disaster incidence frequency was constructed by aggregating the numbers of droughts that have been recorded in each district.	123
Figure 4.11.	The flood hazard estimate based on the frequency of months of extreme rainfall using data between 1960 and 2000.	124
Figure 4.12.	Frequency map of flood disaster incidence created by aggregating the numbers of floods recorded in each district between 1957 and 1995.	125
Figure 4.13.	EM-DAT data on floods from 1975 to 2001 were used to estimate the monthly frequency of floods in the Western Slopes and Eastern Slopes regions.	127
Figure 4.14.	A landslide hazard risk index was estimated based on frequency of incidence.	128
Figure 4.15.	The storm and cyclone tracks for the last 100 years were used to create a cyclone hazard risk map.	129
Figure 4.16.	The monthly count of storms and cyclones between 1887 and 2000	130
Figure 4.17.	Multihazard index constructed by aggregating the hazard indices and scaling the result to range between 0 and 100	131
Figure 4.18.	Multihazard risk estimated by weighting each hazard index by its frequency from 1948 to 1992 and rescaling the result to range from 0 to 100.	132
Figure 4.19.	Multihazard risk estimated by weighting each hazard index by incidence frequency.	133
Figure 4.20.	Multihazard risk estimated by weighting each hazard index by the associated relief expenditure between 1948–1992.	134
Figure 5.1.	Regional elevation map of Caracas and Vargas State	139
Figure 5.2.	Map of the Petare barrio of Caracas, illustrating the dual nature of the city. On the left is the open spacing of the planned, “formal” city. On the right are the densely packed squatter barrios of the “informal” city.	140
Figure 5.3.	An example of the multihazard map produced by the Urban Planning studio.	145
Figure 5.4.	Reserved open space in the Caracas Valley.	147
Figure 5.5.	Hospitals, police stations, and fire stations in the Caracas Valley	147
Figure 5.6.	Proposed interventions in a section of Petare to improve disaster preparedness with the allocation of reserved space	148
Figure 5.7.	The Caracas water supply system, showing key infrastructure crossing fault lines	150
Figure 5.8.	Seismic hazard affecting the city’s transportation network.	152
Figure 5.9.	Debris flows affecting the city’s transportation network.	153
Figure 5.10.	Flood hazard affecting the city’s transportation network. The main east-west thoroughfare through the city is paralleled by the main, channelized river.	154

- Figure 6.1. Location map of Tana River basin in Kenya with the river gauging stations 166
- Figure 6.2. Location map of the Tana River and Garissa Districts with coverage of the Tana River basin in the Garissa District 168
- Figure 6.3. Rainfall for selected stations during El Niño 1997–98 170
- Figure 6.4. Data used in the creation of the DEM for flood-hazard mapping. 171
- Figure 6.5. Variability of river stages at Garissa Town (1933–2001) with special focus on El Niño 1997–98 heights 172
- Figure 6.6. Stream flow modeling at Garissa (1995–1999) 173
- Figure 6.7. Flood hazard map for the 1961 flood (the case of a severe flood before construction of the dams) 176
- Figure 6.8. Flood-hazard map for the El Niño 1997/98 flood (a worst-case scenario after the construction of the dams) 177
- Figure 6.9. (a) Livelihood zones overlaid on El Niño floods case 178
- Figure 6.9. (b) Livelihood zones overlaid on El Niño floods case 178
- Figure 6.10. Livelihood zones overlaid on El Niño flood cases 180
- Figure 6.11. Impacts of floods on market prices and livestock: (a) commodity prices (b) livestock losses 181