

Contents

List of tables	xiii
List of figures	xix
Foreword	xxii
Preface to the first edition	xxv
Preface to the second edition	xxviii
1 Uncertainty in forensic science	1
1.1 Introduction	1
1.2 Statistics and the law	2
1.3 Uncertainty in scientific evidence	5
1.3.1 The frequentist method	6
1.3.2 Stains of body fluids	7
1.3.3 Glass fragments	9
1.4 Terminology	12
1.5 Types of data	15
1.6 Probability	16
1.6.1 Introduction	16
1.6.2 A standard for uncertainty	18
1.6.3 Events	20
1.6.4 Subjective probability	21
1.6.5 Laws of probability	23
1.6.6 Dependent events and background information	25
1.6.7 Law of total probability	29
1.6.8 Updating of probabilities	32
2 Variation	35
2.1 Populations	35
2.2 Samples and estimates	37
2.3 Counts	40
2.3.1 Probabilities	40
2.3.2 Summary measures	41
2.3.3 Binomial distribution	43
2.3.4 Multinomial distribution	44
2.3.5 Hypergeometric distribution	45
2.3.6 Poisson distribution	48
2.3.7 Beta-binomial distribution	51
2.4 Measurements	52
2.4.1 Summary statistics	52
2.4.2 Normal distribution	53

2.4.3	Student's <i>t</i> -distribution	60
2.4.4	Beta distribution	62
2.4.5	Dirichlet distribution	63
2.4.6	Multivariate Normal and correlation	64
3	The evaluation of evidence	69
3.1	Odds	69
3.1.1	Complementary events	69
3.1.2	Examples	70
3.1.3	Definition	70
3.2	Bayes' Theorem	72
3.2.1	Statement of the theorem	72
3.2.2	Examples	73
3.3	Errors in interpretation	78
3.3.1	Fallacy of the transposed conditional	79
3.3.2	Source probability error	81
3.3.3	Ultimate issue error	82
3.3.4	Defender's fallacy	82
3.3.5	Probability (another match) error	83
3.3.6	Numerical conversion error	84
3.3.7	False positive fallacy	85
3.3.8	Uniqueness	86
3.3.9	Other difficulties	87
3.3.10	Empirical evidence of errors in interpretation	89
3.4	The odds form of Bayes' theorem	95
3.4.1	Likelihood ratio	95
3.4.2	Logarithm of the likelihood ratio	99
3.5	The value of evidence	101
3.5.1	Evaluation of forensic evidence	101
3.5.2	Summary of competing propositions	105
3.5.3	Qualitative scale for the value of the evidence	107
3.5.4	Misinterpretations	111
3.5.5	Explanation of transposed conditional and defence fallacies	112
3.5.6	The probability of guilt	116
3.6	Summary	118
4	Historical review	119
4.1	Early history	119
4.2	The Dreyfus case	122
4.3	Statistical arguments by early twentieth-century forensic scientists	125
4.4	People v. Collins	126
4.5	Discriminating power	129
4.5.1	Derivation	129
4.5.2	Evaluation of evidence by discriminating power	130
4.5.3	Finite samples	133
4.5.4	Combination of independent systems	135
4.5.5	Correlated attributes	136
4.6	Significance probabilities	141
4.6.1	Calculation of significance probabilities	141
4.6.2	Relationship to likelihood ratio	144
4.6.3	Combination of significance probabilities	147

4.7	Coincidence probabilities	149
4.7.1	Introduction	149
4.7.2	Comparison stage	151
4.7.3	Significance stage	151
4.8	Likelihood ratio	153

5 Bayesian inference 157

5.1	Introduction	157
5.2	Bayesian inference for a Bernoulli probability	160
5.3	Estimation with zero occurrences in a sample	162
5.4	Estimation of products in forensic identification	165
5.5	Bayesian inference for a Normal mean	166
5.6	Interval estimation	170
5.6.1	Confidence intervals	170
5.6.2	Highest posterior density intervals	172
5.6.3	Bootstrap intervals	172
5.6.4	Likelihood intervals	173
5.7	Odds ratios	175

6 Sampling 179

6.1	Introduction	179
6.2	Choice of sample size	182
6.2.1	Large consignments	182
6.2.2	Small consignments	186
6.3	Quantity estimation	190
6.3.1	Frequentist approach	190
6.3.2	Bayesian approach	191
6.4	Misleading evidence	197

7 Interpretation 205

7.1	Concepts and court cases	205
7.1.1	Relevant population	205
7.1.2	Consideration of odds	206
7.1.3	Combination of evidence	208
7.1.4	Specific cases	208
7.2	Pre-assessment and relevant propositions	214
7.2.1	Levels of proposition	214
7.2.2	Pre-assessment of the case	217
7.2.3	Pre-assessment of the evidence	220
7.3	Assessment of value of various evidential types	221
7.3.1	Earprints	221
7.3.2	Firearms and toolmarks	223
7.3.3	Fingerprints	226
7.3.4	Speaker recognition	228
7.3.5	Hair	229
7.3.6	Documents	231
7.3.7	Envelopes	233
7.3.8	Handwriting	235
7.3.9	Paint	239
7.4	Pre-data and post-data questions	239

8	Transfer evidence	245
8.1	The likelihood ratio	245
8.1.1	Probability of guilt	245
8.1.2	Justification	246
8.1.3	Combination of evidence and comparison of more than two propositions	248
8.2	Correspondence probabilities	254
8.3	Direction of transfer	255
8.3.1	Transfer of evidence from the criminal to the scene	255
8.3.2	Transfer of evidence from the scene to the criminal	260
8.3.3	Transfer probabilities	261
8.3.4	Two-way transfer	270
8.3.5	Presence of non-matching evidence	271
8.4	Grouping	271
8.5	Relevant populations	274
9	Discrete data	283
9.1	Notation	283
9.2	Single sample	283
9.2.1	Introduction	283
9.2.2	General population	286
9.2.3	Particular population	286
9.2.4	Examples	286
9.3	Two samples	288
9.3.1	Two stains, two offenders	288
9.3.2	DNA profiling	291
9.4	Many samples	292
9.4.1	Many different profiles	292
9.4.2	General cases	293
9.5	Relevance of evidence and relevant material	295
9.5.1	Introduction	295
9.5.2	Subjective probabilities	296
9.5.3	Association propositions	296
9.5.4	Intermediate association propositions	297
9.5.5	Examples	298
9.5.6	Two stains, one offender	302
9.6	Summary	304
9.6.1	Stain known to have been left by offenders	304
9.6.2	Relevance: stain may not have been left by offenders	305
9.6.3	Relevance and the crime level	307
9.7	Missing persons	308
9.7.1	Case 1 (Kuo, 1982)	309
9.7.2	Case 2 (Ogino and Gregonis, 1981)	309
9.7.3	Calculation of the likelihood ratio	310
9.8	Paternity: combination of likelihood ratios	312
9.8.1	Likelihood of paternity	314
9.8.2	Probability of exclusion in paternity	317

10	Continuous data	319
10.1	The likelihood ratio	319
10.2	Normal distribution for between-source data	321
10.2.1	Sources of variation	322
10.2.2	Derivation of the marginal distribution	322
10.2.3	Approximate derivation of the likelihood ratio	324
10.2.4	Lindley's approach	326
10.2.5	Interpretation of result	327
10.2.6	Examples	328
10.3	Estimation of a probability density function	330
10.4	Kernel density estimation for between-source data	337
10.4.1	Application to medullary widths of cat hairs	339
10.4.2	Refractive index of glass	339
10.5	Probabilities of transfer	342
10.5.1	Introduction	342
10.5.2	Single fragment	342
10.5.3	Two fragments	345
10.5.4	A practical approach to glass evaluation	349
10.5.5	Graphical models for the assessment of transfer probabilities	352
10.6	Approach based on <i>t</i> -distribution	353
10.7	Appendix Derivation of <i>V</i> when the between-source measurements are assumed normally distributed	357
11	Multivariate analysis	359
11.1	Introduction	359
11.2	Description of example	360
11.3	Univariate <i>t</i> -tests	362
11.4	Hotelling's T^2	363
11.5	Univariate Normality, two sources of variation	365
11.6	Multivariate Normality, two sources of variation	366
11.7	Caveat lector	371
11.8	Summary	372
11.9	Appendix	373
11.9.1	Matrix terminology	373
11.9.2	Determination of a likelihood ratio with an assumption of Normality	377
12	Fibres	381
12.1	Introduction	381
12.2	Likelihood ratios in scenarios involving fibres	381
12.2.1	Fibres evidence left by an offender	382
12.2.2	Comments on the fibres scenario	387
12.2.3	Fibres evidence not left by the offender	388
12.2.4	Cross-transfer	389
12.3	Pre-assessment in fibres scenarios	392
12.3.1	The case history	392
12.3.2	Formulation of the pairs of propositions and events	392
12.3.3	Assessment of the expected value of the likelihood ratio	394
12.4	The relevant population of fibres	396

13	DNA profiling	399
13.1	Introduction	399
13.2	Hardy–Weinberg equilibrium	401
13.3	DNA likelihood ratio	404
13.4	Uncertainty	405
13.5	Variation in sub-population allele frequencies	405
13.6	Related individuals	409
13.7	More than two propositions	412
13.8	Database searching	414
	13.8.1 Search and selection effect (double counting error)	418
13.9	Island problem	419
13.10	Mixtures	421
13.11	Error rate	424
14	Bayesian networks	429
14.1	Introduction	429
14.2	Bayesian networks	430
	14.2.1 The construction of Bayesian networks	431
14.3	Evidence at the crime level	437
	14.3.1 Preliminaries	437
	14.3.2 Description of probabilities required	437
14.4	Missing evidence	439
	14.4.1 Preliminaries	439
	14.4.2 Determination of a structure for a Bayesian network	440
	14.4.3 Comments	442
14.5	Error rates	442
	14.5.1 Preliminaries	442
	14.5.2 Determination of a structure for a Bayesian network	443
14.6	Transfer evidence	444
	14.6.1 Preliminaries	444
	14.6.2 Determination of a structure for a Bayesian network	444
	14.6.3 Comment on the <i>transfer</i> node	446
14.7	Combination of evidence	447
14.8	Cross-transfer evidence	449
	14.8.1 Description of nodes	451
	14.8.2 Probabilities for nodes	452
14.9	Factors to consider	452
	14.9.1 Parameter choice	453
14.10	Summary	453
References		455
Notation		485
Cases		489
Author index		491
Subject index		499

List of Tables

1.1	Genotypic frequencies for locus <i>LDLR</i> amongst Caucasians in Chicago based on a sample of size 200 (from Johnson and Peterson, 1999).	8
1.2	Refractive index measurements.	10
1.3	Genotype probabilities, assuming Hardy-Weinberg equilibrium, for a diallelic system with allele probabilities p and q .	27
1.4	The proportion of people in a population who fall into the four possible categories of genetic markers.	29
2.1	Probabilities for the number of sixes, X , in four rolls of a fair six-sided die	41
2.2	Intermediate calculations for the variance of the number of sixes, x , in four rolls of a fair six-sided die	43
2.3	Probabilities that a random variable X with a Poisson distribution with mean 4 takes values 0, 1, 2, 3, 4 and greater than 4	50
2.4	Values of cumulative distribution function $\Phi(z)$ and its complement $1 - \Phi(z)$ for the standard Normal distribution for given values of z .	57
2.5	Probabilities for absolute values from the standard Normal distribution function	57
2.6	Percentage points $t_{(n-1)}(P)$ for the t -distribution for given values of sample size n , degrees of freedom $(n - 1)$ and P , and the corresponding point $z(P)$ for the standard Normal distribution.	61
3.1	Two-by-two contingency table for frequencies for the tabulation of patients with or without a disease (S or \bar{S}) given a blood test positive or negative (R or \bar{R}).	74
3.2	Two-by-two contingency table for probabilities for the tabulation of patients with or without a disease (S or \bar{S}) given a blood test positive or negative (R or \bar{R}).	75
3.3	Hypothetical results for deaths amongst a population.	76

3.4	Probability θ of at least one match, given a frequency of the trace evidence of γ , in a population of size 1 million.	83
3.5	Evidence occurs with RMP γ . Smallest number ψ of people to be observed before a match with the evidence occurs with a given probability, $Pr(M) = 0.5, 0.9$; $\psi_5 = \log 0.5 / \log(1 - \gamma)$, $\psi_9 = \log 0.1 / \log(1 - \gamma)$, n_5 is the smallest integer greater than ψ_5 , n_9 is the smallest integer greater than ψ_9 .	85
3.6	The probability, θ' , of at least one match with the evidence, which occurs with RMP γ , when $n' = 1/\gamma$ people are tested.	85
3.7	Frequency of Kell and Duffy types by colour in a hypothetical area. (Reproduced by permission of The Forensic Science Society.).	97
3.8	Effect on prior odds in favour of H_p relative to H_d of evidence E with value V of 1 000. Reference to background information I is omitted.	104
3.9	Value of the evidence for each genotype.	107
3.10	Qualitative scale for reporting the value of the support of the evidence for H_p against H_d (Evetts <i>et al.</i> , 2000a).	107
3.11	The values of the logarithm of the posterior odds in favour of an issue determined from the values of the logarithm of the prior odds in favour of guilt ($\log(\text{Prior odds})$) and the logarithm of the likelihood ratio (Aitken and Taroni, 1998). The values in the body of the table are obtained by adding the appropriate row and column values. Logarithms are taken to base 10. The verbal description is taken from Calman and Royston (1997). (Reproduced by permission of The Forensic Science Society.).	110
3.12	Probability of guilt required for proof beyond reasonable doubt (Simon and Mahan, 1971).	116
4.1	Probabilities suggested by the prosecutor for various characteristics of the couple observed in the case of <i>People v. Collins</i> .	127
4.2	Allelic frequencies for <i>TPOX</i> locus for Swiss and NZ Caucasians and the probability Q_{TPOX} of a match.	136
4.3	Allelic frequencies for <i>TH01</i> locus for Swiss and NZ Caucasians and the probability Q_{TH01} of a match.	136
4.4	The calculation of discriminating power (DP) for Normal distributions of p dimensions.	139
4.5	Significance probabilities P for refractive index x of glass for mean $\theta_0 = 1.518458$ and standard deviation $\sigma = 4 \times 10^{-5}$ and decisions assuming a significance level of 5%	143
4.6	Variation in the likelihood ratio V , as given by (4.7), with sample size n , for a standardised distance $z_n = 2$, a result which is significant at the 5% level.	147

5.1	Probabilistic upper bounds θ_0 for the proportion θ of occurrences of an event in a population when there have been no occurrences in a sample of size n , for given values of $\epsilon = Pr(\theta > \theta_0)$, with a beta prior for which $\alpha = \beta = 1$.	164
5.2	Numbers of banknotes contaminated.	176
5.3	The value of evidence in comparing two binomial proportions for different values of sample size n and number of 'successes' z , where the samples are banknotes and a 'success' is a contaminated banknote. The binomial proportion in the numerator is 0.83, estimated from banknotes known to be used for drug dealing, and the binomial proportion in the denominator is 0.57, estimated from banknotes known to be from general circulation.	178
6.1	The probability that the proportion of drugs in a large consignment is greater than 50% for various sample sizes m and prior parameters α and β . (Reprinted with permission from ASTM International.)	184
6.2	The sample size required to be 100% certain that the proportion of units in the consignment which contain drugs is greater than θ , when all the units inspected are found to contain drugs. The prior parameters $\alpha = \beta = 1$. (Reprinted with permission from ASTM International.)	185
6.3	Estimates of quantities q of drugs (in grams), in a consignment of $m + n$ units, according to various possible burdens of proof, expressed as percentages $P = 100 \times Pr(Q > q \mid m, z, n, \bar{x}, s)$ in 26 packages when 6 packages are examined ($m = 6, n = 20$) and $z = 6, 5$, or 4 are found to contain drugs. The mean (\bar{x}) and standard deviation (s) of the quantities found in the packages examined which contain drugs are 0.0425 g and 0.0073 g. The parameters for the beta prior are $\alpha = \beta = 1$. Numbers in brackets are the corresponding frequentist lower bounds using the fpc factor (6.2). (Reprinted with permission from ASTM International.)	194
6.4	Estimates of quantities q of drugs (in grams), in a consignment of $m + n$ units, according to various possible burdens of proof, expressed as percentages $P = 100 \times Pr(Q > q \mid m, z, n, \bar{x}, s)$ in 2600 packages when 6 packages are examined ($m = 6, n = 2594$) and $z = 6, 5$, or 4 are found to contain drugs. The mean (\bar{x}) and standard deviation (s) of the quantities found in the packages examined which contain drugs are 0.0425 g and 0.0073 g. The parameters for the beta prior are $\alpha = \beta = 1$. Numbers in brackets are the corresponding frequentist lower bounds without using the fpc factor (6.8). (Reprinted with permission from ASTM International.).	196

6.5	Probabilities of strong misleading evidence $M(n)$ and weak evidence $W(n)$ for boundary values k of 8 and 32 for strong evidence and sample sizes n of 5, 10 and 20.	203
7.1	Examples of the hierarchy of propositions. (Reproduced by permission of The Forensic Science Society.)	217
7.2	Hypothetical relative frequencies for the maximum CMS count y for bullets fired from the same gun $f(y SG)$ and for bullets fired from different guns $f(y DG)$, and likelihood ratios $V = f(y SG)/f(y DG)$. (Reprinted with permission from ASTM International.)	224
7.3	Poisson probabilities for the maximum CMS count y for bullets fired from the same gun $Pr(Y = y SG)$ (7.1) with mean $\lambda_s = 3.91$ and for bullets fired from different guns $Pr(Y = y DG)$ (7.2) with mean $\lambda_D = 1.325$ and likelihood ratio $V = Pr(Y = y SG)/Pr(Y = y DG)$. (Reprinted with permission from ASTM International.)	225
7.4	Values of a probabilistic γ upper bound for the probability ϕ_0 of a false match for different values of n and γ .	242
7.5	Values of a probabilistic γ lower bound for the likelihood ratio $1/\phi_0$ for different values of n and γ , with values rounded down.	242
8.1	Probability evidence in <i>State v. Klindt</i> (altered for illustrative purposes) from Lenth (1986).	249
8.2	Frequencies of Ruritanians and those of genotype Γ for locus <i>LDLR</i> in a hypothetical population.	256
8.3	Distribution of blood groups of innocently acquired bloodstains on clothing of people of type O , compared with the distribution in the general population.	268
8.4	Critical values $r(\alpha, n)$ for the range of a sample of size n from a standardised Normal distribution, from Owen (1962).	273
8.5	Critical values $\lambda(\alpha, n)$ for samples of size n for the divisive grouping algorithm from Triggs <i>et al.</i> (1997).	274
8.6	Probability that a criminal lived in a particular area, given the crime was committed in Auckland, New Zealand.	280
9.1	Gene frequencies for New Zealand in the <i>ABO</i> system.	287
9.2	Gene frequencies for Chinese in New Zealand in the <i>ABO</i> system.	287
9.3	Phenotypes for two cases of missing persons.	308
9.4	Gene frequencies and phenotypic incidences for Case 1 (Kuo, 1982).	309
9.5	Gene frequencies and phenotypic incidences for Case 2 (Ogino and Gregonis, 1981).	309
9.6	Frequencies for up to three codominant alleles.	310
9.7	Relative frequencies for <i>ABO</i> system.	310

9.8	Probabilities that the parents will pass the specified stain phenotype, Case 1.	311
9.9	Probabilities that the parents will pass the specified phenotypes, Case 2.	311
9.10	Two pieces of evidence on DNA markers.	312
9.11	Posterior probabilities of paternity for various prior probabilities for evidence for alleged father $E_1 = 11 - 13$, $E_2 = 18 - 18$.	316
9.12	Hummel's likelihood of paternity.	317
10.1	Likelihood ratio values for varying values of $(y - \bar{x})/\sigma$ and $(y - \mu)/\tau$.	326
10.2	Value of $\tau(2^{1/2}\sigma)^{-1} \exp(-\frac{1}{2}\lambda^2 + \frac{1}{2}\delta^2)$ (10.7) as a function of $\lambda = \bar{x} - \bar{y} /(2^{1/2}\sigma)$ and $\delta = z - \mu /\tau$ for $\tau/\sigma = 100$.	329
10.3	Medullary widths in microns of 220 cat hairs (Peabody <i>et al.</i> , 1983).	332
10.4	Value of the evidence for various values of \bar{x} and \bar{y} , the smoothing parameter λ and the within-cat standard deviation σ ; $m = n = 10$ throughout; $s = 23$ microns.	339
10.5	Refractive index of 2269 fragments of float glass from buildings, Lambert and Evett (1984).	340
10.6	Coincidence probability and value of the evidence (kernel and Lindley approaches) for various values of \bar{x} and \bar{y} and the smoothing parameter λ (for the kernel approach); $m = 10$, $n = 5$; within-window standard deviation $\sigma = 0.000\ 04$, between-window standard deviation $\tau = 0.004$; overall mean $\mu = 1.5182$.	341
10.7	Distributional parameters for glass problems.	344
10.8	Transfer probabilities for glass problems.	344
10.9	Some values for the likelihood ratio V for the single-fragment case, from Evett (1986).	345
10.10	Possible sources of two fragments.	345
10.11	Refractive indices of glass fragments for Johnston, recovered, and a control set with means, separate and pooled standard deviations (s.d). The number of recovered fragments $n_y = 11$ and the number of control fragments $n_x = 10$. Example presented in Walsh <i>et al.</i> (1996).	355
10.12	Summary statistics for concentration of dye CI 14720 in illicit pills (Goldmann <i>et al.</i> , 2004).	356
12.1	Findings from the analysis of the hair combings.	394
12.2	Events and probabilities relating to findings under H_p and H_d .	395
12.3	Likelihood ratios for the outcomes from Table 12.1 with $t_0 = 0.01$, $t_s = 0.04$, $t_l = 0.95$; $p_0 = 0.78$, $p_1 = 0.22$; $s_s = 0.92$, $s_l = 0.08$; $m = 0.05$, as proposed by Champod and Jackson (2000).	396
13.1	Hardy-Weinberg proportions for a locus with two alleles, A and a , with frequencies p and q such that $p + q = 1$.	402

13.2	Observed and expected frequencies of <i>HumTH01</i> genotypes based on 95 unrelated Turkish individuals (from Çakir <i>et al.</i> , 2001).	403
13.3	Effects of population structure, as represented by F_{ST} , on the likelihood ratio, the reciprocal of the conditional match probability (13.8) for heterozygotes between alleles with equal frequency p .	408
13.4	Match probability $Pr(G G, H_d, I)$ that a relative has the same genotype as the suspect and the corresponding value for V assuming allelic frequencies of 0.1, from Weir and Hill (1993). (Reproduced by permission of The Forensic Science Society.)	410
13.5	Effects of family relatedness on match probability, $Pr(G_c G_s, H_d, I)$, from Weir (2001a). Note the use of θ for F_{ST} for clarity. ©2001, John Wiley & Sons, Ltd. Reproduced with permission.	411
13.6	General match probability values recommended for use when reporting full SGM-plus profile matches, with an F_{ST} value of 0 in situation 6 and a value of 0.02 for situations 1–5, from Foreman and Evett (2001). (Reproduced by permission of Springer-Verlag.)	411
13.7	Likelihood ratio for three-allele mixed sample, heterozygous victim and homozygous suspect. The victim has alleles a, b , the suspect has allele c . There are three ethnic groups under consideration, labelled 1, 2 and 3. The co-ancestry coefficient F_{ST} for the ethnic group, 1, of the unknown donor is labelled θ_1 for clarity. The allelic frequencies are p_{a1}, p_{b1} and p_{c1} in ethnic group 1. Ethnicities of the unknown donor, the victim and the suspect are considered. From Fung and Hu (2002). (Reproduced by permission of Springer-Verlag.)	423
13.8	Posterior odds that a suspect is the source of a sample that reportedly has a matching DNA profile, as a function of prior odds, random match probability, and false positive probability. Extracted from Thompson <i>et al.</i> (2003). (Reprinted with permission from ASTM international.)	426
14.1	Conditional probabilities for transfer with one group. The features of the control object are X , and the features of the recovered fibres are Y , which may be from 0, 1 or 2 groups. These recovered fibres may be present by chance alone (b_1) or not (b_0), or have been transferred from the victim (t_1) or not (t_0).	445
14.2	Conditional probabilities for transfer with many groups, as illustrated in Section 12.3.3. The transfer may be of no fibres (t_0), a small amount (t_s) or a large amount (t_l). A group may be present (p_1) or not (p_0), and the group may be small (s_s) or large (s_l). The outcome O has five categories: no groups, one not-matching group, one small matching group, one large matching group and two groups.	447

List of Figures

2.1	Probability density function for a Normal distribution, with mean 0.7 and variance 0.005.	55
2.2	Selected tail area probabilities for a standard Normal random variable Z : (a) $Pr(Z > 1)$, (b) $Pr(Z > 2)$, (c) $Pr(Z < 2)$, (d) $Pr(Z > 2.5)$.	58
4.1	Discrimination between two groups, identified by Δ and by \circ , with two perfectly correlated variables.	140
5.1	Prior density function $f(\theta \alpha, \beta)$ with $\alpha = 61$, $\beta = 44$, likelihood function $L(\theta n, x)$ with $n = 372$, $x = 201$, and posterior density function $f(\theta x + \alpha, n - x + \beta)$ for a Bernoulli parameter. (Reproduced from Weir, 1996a, by permission of Sinauer Associates, Inc.)	162
5.2	Standardised likelihood function for the proportion of people of blood group Γ , from a sample of size 30 in which 6 were of group Γ . A likelihood interval, derived from the observed data, of fairly strong support for the values of γ included within it is indicated with the dotted lines: (0.081, 0.372).	174
6.1	The prior probability $1 - F(\theta)$ that the proportion of units in a consignment is greater than θ , for various choices of α and β : $\alpha = \beta = 1$ (dot-dashed curve), $\alpha = \beta = 0.5$ (solid), $\alpha = 0.065$, $\beta = 0.935$ (dotted). (Reprinted with permission from ASTM International.)	186
6.2	The posterior probability $1 - F(\theta)$ that the proportion of units in a consignment is greater than θ , for various choices of α and β : $\alpha = \beta = 1$ (dot-dashed curve), $\alpha = \beta = 0.5$ (solid), $\alpha = 0.065$, $\beta = 0.935$ (dotted), after observation of four units, all found to be illegal. The corresponding probabilities that at least 50% of the consignment contains illegal units are marked as 0.985 ($\alpha = \beta = 0.5$), 0.970 ($\alpha = \beta = 1$), 0.950 ($\alpha = 0.065$, $\beta = 0.935$). (Reprinted with permission from ASTM International.)	187

6.3	The probability that the total quantity Q of drugs (in grams) in a consignment of 26 packages is greater than q when 6 packages are examined and 6 (solid curve), 5 (dashed), or 4 (dot-dashed) are found to contain drugs. The mean and standard deviation of the quantities found in the packages examined which contain drugs are 0.0425 g and 0.0073 g. The parameters for the beta prior are $\alpha = \beta = 1$. (Reprinted with permission from ASTM International.)	195
6.4	The probability that the total quantity Q of drugs (in grams) in a consignment of 2600 packages is greater than q when 6 packages are examined and 6 (solid curve), 5 (dashed), or 4 (dot-dashed) are found to contain drugs. The mean and standard deviation of the quantities found in the packages examined which contain drugs are 0.0425 g and 0.0073 g. The parameters for the beta prior are $\alpha = \beta = 1$. (Reprinted with permission from ASTM International.)	196
6.5	Bump function for the probability of misleading evidence $Pr_1\left(\frac{f_{2n}}{f_{1n}} > k\right)$ for $k = 8$ and $k = 32$ as a function of c , the distance from the true mean to the alternative, in standard errors. (Royall, 2000; reprinted with permission from the <i>Journal of the American Statistical Association</i> . Copyright 2000 by the American Statistical Association. All rights reserved.)	201
9.1	Variation in the logarithm to base 10 of the likelihood ratio V of the evidence with p , the probability that the stain would have been left by the suspect even though he was innocent of the offence, for various values of r , the probability that the stain would have been left by one of the offenders. The number of offenders, k , equals 4 and the relative frequency of the profile γ is 0.001. Adapted from Evett (1993a), with the inclusion of a curve for $r = 0$. The dotted line at $\log(V) = 0$ indicates where the evidence is equally likely under both propositions.	301
10.1	Refractive index measurements from 2269 fragments of float glass from buildings (from Lambert and Evett, 1984).	323
10.2	Medullary width (in microns) of 220 cat hairs (from Peabody <i>et al.</i> , 1983).	331
10.3	Examples of kernel density estimates showing individual kernels. Smoothing parameter values are (a) $\lambda = 0.5$ and (b) $\lambda = 1$.	333
10.4	Medullary widths, in microns, of cat hairs (Peabody <i>et al.</i> , 1983) and associated kernel density estimate with smoothing parameter equal to 0.09.	335

10.5	Medullary widths, in microns, of cat hairs (Peabody <i>et al.</i> , 1983) and associated kernel density estimate with smoothing parameter equal to 0.50.	335
10.6	Kernel density estimate with smoothing parameter 0.025 of refractive index measurements from 2269 fragments of float glass from buildings (Lambert and Evett, 1984).	336
10.7	Kernel density estimate with smoothing parameter 0.25 of refractive index measurements from 2269 fragments of float glass from buildings (Lambert and Evett, 1984).	337
10.8	The difference in refractive index measurements (higher – lower) for each pair of fragments for individuals who had two fragments of glass on their clothing, from Harrison <i>et al.</i> (1985). (Reproduced from Evett, 1986, by permission of The Forensic Science Society.)	347
10.9	Graphs of $\log_{10} \phi(1)$, $\log_{10} \phi(2)$ and $\log_{10} V = \log_{10}(\phi(1) + \phi(2))$ against $ \bar{y} - \bar{x} / \sigma$ for $\bar{y} = \mu$, for the transfer of two fragments of glass from the scene of the crime to the criminal: (a) $ y_1 - y_2 = \sigma$, (b) $ y_1 - y_2 = 4\sigma$. The value of the evidence is $V = \phi(1) + \phi(2)$. The dotted line is $\log_{10} V$. (Adapted from Evett, 1986.)	348
14.1	Basic connections in Bayesian networks: (a) serial, (b) diverging and (c) converging.	432
14.2	Bayesian network for evidence E and proposition H .	435
14.3	Bayesian network for a serial connection for a reported match RM in a DNA profile, where M denotes a match and H a proposition.	436
14.4	Four-node network for evaluation of evidence at the crime level.	437
14.5	Bayesian network fragments representing the relation between (a) the variables E and H , and (b) the variables M and E ; (c) Bayesian network for missing evidence.	441
14.6	Bayesian network for error rates. The probabilistic dependencies are indicated on the right-hand side of each node.	443
14.7	Five-node network: no outcome.	445
14.8	Five-node network: outcome, converging and serial in Bayesian networks.	446
14.9	Complete network for evaluation of cross-transfer evidence in DNA profiles. (Reprinted from Aitken <i>et al.</i> , 2003, with permission from Elsevier.)	451