

Contents

List of Figures.....	ix
List of Tables.....	xxi
Preface.....	xxv

SECTION I

Introduction on arid zone soil

Ariele Singer.....	1
--------------------	---

Chapter 1

Arid zone soils: Nature and properties.....	3
---	---

SECTION II

Biogeochemistry of trace elements in arid environments

Fengxiang X. Han.....	45
-----------------------	----

Chapter 2

Trace element distribution in arid zone soils.....	47
--	----

Chapter 3

Solution chemistry of trace elements in arid zone soils.....	69
--	----

Chapter 4

Selective sequential dissolution for trace elements in arid zone soils.....	107
--	-----

Chapter 5

Binding and distribution of trace elements among solid-phase components in arid zone soils.....	131
--	-----

Chapter 6

Transfer fluxes of trace elements in arid zone soils – a case study: Israeli arid soils	169
--	-----

Chapter 7
Bioavailability of trace elements in arid zone soils.....221

Chapter 8
Trace element pollution in arid zone soils.....267

Chapter 9
Global perspectives of anthropogenic interferences
in the natural trace element distribution: Industrial age inputs
of trace elements into the pedosphere 303

References 329

Index 357

List of Figures

- Figure 1.1. (a) Climate categories based on mean annual precipitation; (b) Desert and steppe boundaries as a function of both mean annual precipitation and temperature according to the Koeppen system; arrows show how boundaries shift according to whether precipitation falls mainly in the summer or winter (after Monger et al., 2004. Reprinted from *Encyclopedia of Soils in the Environment*, D. Hillel, Tropical soils: Arid and Semi-arid, p183, Copyright (2004), with permission from Elsevier).
- Figure 1.2. Global land area by aridity zone (%) (after UNEP, 1993. Reprinted from *World Atlas of Desertification*, Figure of global land area by aridity zones, p5, Copyright (1993), with permission from UNEP).
- Figure 1.3. Distributions of arid and semi-arid areas (Reprinted from *Soils of Arid Regions*, Dregne H.E., p 6, Copyright (1976), with permission from Elsevier).
- Figure 1.4. Phase diagram for the system $K_2O-Al_2O_3-SiO_2-H_2O$ at 25 °C and 1 atmosphere. Open circles are analytical data for water from springs or seeps in a granitic terrain in the Sierra Nevada (from Feth and others) (%) (from Birkeland, 1984, after Feth and others, 1964. Reprinted from *Soils and Geomorphology*, p111, Copyright (1984), with permission from Oxford University Press, New York, Oxford).
- Figure 1.5. Schematic Reg soil pedon: the gravel is chalk, underneath the gravel cover is the porous vesicular A horizon (%) (from Singer, 2007, Reprinted from *The Soils of the Land of Israel*, Copyright (2007), with permission from Springer).
- Figure 1.6. The transport of fine particles by duststorm (after Singer, 2007. Reprinted from *The Soils of the Land of Israel*, Copyright (2007), with permission from Springer).
- Figure 1.7. Distribution of salts in chloride-sulfate Solonchak during the various seasons of the year (after Ravikovitch and Bidner-Bar Hava, 1948) (from Singer, 2007, after Ravikovitch and Bidner-BarHava, 1948. Reprinted from *The Soils of the Land of Israel*, Copyright (2007), with permission from Springer).
- Figure 1.8. Cracks, surface mulch and soil structure in a Vertisol during the dry season (from FAO, 2001. Reprinted from *World Soil Resources Reports 94*, P. Driessen, J. Deckers, O. Spaargaren, F. Nachtergaele, eds., p80, Copyright (2001), with permission from FAO).

- Figure 1.9. Illustration of soil taxonomy suborders and great groups that have arid moisture regimes (shaded) and their moisture counterparts that have ustic and xeric moisture regimes (after Monger, 2002. Reprinted from Encyclopedia of Soil Science, H. Tan, ed., Arid Soils, Monger, p86, Copyright (2002), with permission from Marcel Dekker).
- Figure 2.1. Distribution of chemical elements in the earth's crust (mg/kg).
- Figure 2.2. Elemental enrichment factors in soils, plotted on a log scale against the ionic potential of the elements.
- Figure 2.3. Elemental enrichment factors in bacterial and fungi, plotted on a log scale against the ionic potential of the elements (after Banin and Navrot, 1975. Reprinted from Science, 189, Banin A. and Navrot J., Origin of Life: Clues from relations between chemical compositions of living organisms and natural environments, pp 550-551, Copyright (1975), with permission from AAAS).
- Figure 2.4. Elemental enrichment factors in plants and land animals, plotted on a log scale against the ionic potential of the elements (after Banin and Navrot, 1975. Reprinted from Science, 189, Banin A. and Navrot J., Origin of Life: Clues from relations between chemical compositions of living organisms and natural environments, pp 550-551, Copyright (1975), with permission from AAAS).
- Figure 2.5. Elemental enrichment factors in seawater, related to the ionic potential of the elements (after Banin and Navrot, 1975. Reprinted from Science, 189, Banin A. and Navrot J., Origin of Life: Clues from relations between chemical compositions of living organisms and natural environments, pp 550-551, Copyright (1975), with permission from AAAS).
- Figure 2.6. Effects of soil pH on total Cu concentrations in Israeli arid and semi-arid soils (R: 0.50, N: 39).
- Figure 3.1. Changes of Zn activity in 10 Colorado soils with pH (data extracted from Ma and Lindsay, 1990).
- Figure 3.2. Changes of Cd speciation in soil solutions of a typical Israeli calcareous soil with pH 4 - 9 (after Hirsch and Banin, 1990, with permission from Soil Sci. Soc. Am).
- Figure 3.3. Changes of Cd speciation in soil solutions of Cd nitrate-treated Domino soil from California (data extracted from Candelaria and Chang, 1997).

- Figure 3.4. Effects of phosphate levels on Cd and Zn solution speciation in California soils that received sludge application (data extracted from Villarroel et al., 1993).
- Figure 3.5. Effects of Cl concentrations (NaCl or CaCl_2) on Cd solution speciation in a California soil (data extracted from Bingham et al., 1983).
- Figure 4.1. Removal of carbonate from Israeli arid soils as indicated by the X-ray diffractograms after extraction of the carbonate fraction by NaOAc-HOAc solutions at various pHs for 16 hours. C: calcite: $d = 3.04 \text{ \AA}$, and D: dolomite, $d = 2.89 \text{ \AA}$. Number 1, 2, 3, 4, 5, and 6 indicate non-treated soil (No. 1), treatments (No. 2-6) with NaOAc-HOAc solutions at pH 7.0, 6.0, 5.5, 5.0 and 4.0, respectively (after Han and Banin, 1995. Reprinted from *Commun Soil Sci Plant Anal*, 26, Han and Banin A., Selective sequential dissolution techniques for trace metals in arid-zone soils: The carbonate dissolution step, p 563, Copyright (1995), with permission from Taylor & Francis US).
- Figure 4.2. Dissolution of Ca from Israeli arid soils by NaOAc-HOAc solutions at various pHs after the extraction of the exchangeable fraction (after Han and Banin, 1995. Reprinted from *Commun Soil Sci Plant Anal*, 26, Han and Banin A., Selective sequential dissolution techniques for trace metals in arid-zone soils: The carbonate dissolution step, p 568, Copyright (1995), with permission from Taylor & Francis US).
- Figure 4.3. Dissolution kinetics of CaCO_3 by NaOAc-HOAc solutions at varying pHs. a: Dissolution of pure CaCO_3 by NaOAc-HOAc (0.1 g CaCO_3 , 25 mL buffer solution), b and c: Dissolution of soil CaCO_3 by NaOAc-HOAc at pH 5.0 (1.0 g soil, 25 ml buffer solution) (after Han and Banin, 1995. Reprinted from *Commun Soil Sci Plant Anal*, 26, Han and Banin A., Selective sequential dissolution techniques for trace metals in arid-zone soils: The carbonate dissolution step, p 565, Copyright (1995), with permission from Taylor & Francis US).
- Figure 4.4. Effects of pH of NaOAc-HOAc solutions for the CARB fraction on the subsequent fractions (Soil CaCO_3 : 33.3%) (after Han and Banin, 1995. Reprinted from *Commun Soil Sci Plant Anal*, 26, Han and Banin A., Selective sequential dissolution techniques for trace metals in arid-zone soils: The carbonate dissolution step, p 572, Copyright (1995), with permission from Taylor & Francis US).

- Figure 4.5. Cumulative sums of extractable elements in soil J3 (CaCO_3 : 33.3%) in consecutive steps of the selective sequential dissolution procedure as a function of the cumulative steps. 1: CARB (carbonate); 2: CARB (carbonate) + ERO (easily reducible oxide); 3: CARB + ERO + OM (organic matter); 4: CARB + ERO + OM + RO (reducible oxide) (after Han and Banin, 1995. Reprinted from *Commun Soil Sci Plant Anal*, 26, Han and Banin A., Selective sequential dissolution techniques for trace metals in arid-zone soils: The carbonate dissolution step, p 573, Copyright (1995), with permission from Taylor & Francis US).
- Figure 4.6. Cumulative sums of extractable Ca, Mg, and Mn in arid soil J3, B10, and S1 in consecutive steps of two sequential dissolution procedures.
- Figure 5.1. Correlation between Co and Fe contents in Israeli arid soils.
- Figure 5.2. Correlation between Cu and MnO contents in Israeli arid soils.
- Figure 5.3. Effects of soil pH on the Zn amount bound to the Fe oxide fraction (amorphous/crystalline Fe oxide and overall Fe oxide bound fractions) in soils from China with pH 3.73 - 8.1 and 0 - 14.7% CaCO_3 (after Han et al., 1995. Reprinted from *Geoderma*, 66, Han F.X., Hu A.T., Qi H.Y., Transformation and distribution of forms of zinc in acid, neutral and calcareous soils of China, p 128, Copyright (1995), with permission from Elsevier).
- Figure 5.4. Effects of pH on Zn adsorption on an untreated and treated soils with peroxide and dithionite.
- Figure 5.5. Kinetics of Zn adsorption on a soil and soil with removals of organic matter and Fe oxides (1 g soil reacted with 20 ml of 10 mg/L Zn solution concentration).
- Figure 5.6. Relationship between Co and Mn contents extracted from solid-phase of six Israeli arid-zone soils with sequential dissolution procedures (after Han et al., 2002b. Reprinted from *J Environ Sci Health, Part A*, 137, Han F.X., Banin A., Kingery W.L., Li Z.P., Pathways and kinetics of transformation of cobalt among solid-phase components in arid-zone soils, p 184, Copyright (2002), with permission from Taylor & Francis).
- Figure 6.1. The fractional loading isotherms of Cu in a contaminated Israeli loessial soil at an initial (one hour) period and after 48 weeks. The soil was treated with increasing levels of metal nitrates and was incubated under the field capacity regime. Horizontal solid line represents the native content of Cu in the nonamended soil (*Figure 6.1 – Figure 6.4*, after Han and Banin, 2001. Reprinted from *Commun Soil Sci Plant Anal*, 32, Han F.X and Banin A., The

fractional loading isotherm of heavy metals in an arid-zone soil, pp 2700-2703, Copyright (2001), with permission from Taylor & Francis)

- Figure 6.2. The fractional loading isotherms of Cr in a contaminated Israeli loessial soil at an initial (one hour) period and after 48 weeks.
- Figure 6.3. The fractional loading isotherms of Ni in a contaminated Israeli loessial soil at an initial (one hour) period and after 48 weeks.
- Figure 6.4. The fractional loading isotherms of Zn in a contaminated Israeli loessial soil at an initial (one hour) period and after 48 weeks.
- Figure 6.5. The initial reduced partition index, I_R , of six metals in two Israeli arid soils. Two soils were treated with metal nitrates at various loading levels. Soils were incubated under the field capacity moisture regime (modified after Han and Banin, 1999, with permission from Springer Science and Business Media).
- Figure 6.6. Long-term changes of U_{ts} of metals in a loessial soil from Israel. The soil received metal nitrates under the field capacity moisture regime (after Han and Banin, 1999. Reprinted from Water Air Soil Pollut, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 245, Copyright (1999), with permission from Springer Science and Business Media).
- Figure 6.7. Ranges (arithmetic mean and standard deviation) of the reduced partition index (I_R) of Cd, Cu, Cr, Ni and Zn in 45 Israeli arid-zone soils (after Han et al., 2003a. Reprinted from Adv Environ Res, 8, Han F.X., Banin A., Kingery W.L., Triplett G.B., Zhou L.X., Zheng S.J., Ding W.X., New approach to studies of redistribution of heavy metals in soils, p 118, Copyright (2003), with permission from Elsevier).
- Figure 6.8. Changes of Cd fractions in two Israeli soils during 336 days of incubation at the field capacity moisture regime (after Han and Banin, 1999. Reprinted from Water Air Soil Pollut, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 238, Copyright (1999), with permission from Springer Science and Business Media).

- Figure 6.9. Changes of Cr fractions in two Israeli soils during 336 days of incubation at the field capacity moisture regime (after Han and Banin, 1999. Reprinted from *Water Air Soil Pollut*, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 239, Copyright (1999), with permission from Springer Science and Business Media).
- Figure 6.10. Changes of Cu fractions in two Israeli soils during 336 days of incubation at the field capacity moisture regime (after Han and Banin, 1999. Reprinted from *Water Air Soil Pollut*, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 240, Copyright (1999), with permission from Springer Science and Business Media).
- Figure 6.11. Changes of Ni fractions in two Israeli soils during 336 days of incubation at the field capacity moisture regime (after Han and Banin, 1999. Reprinted from *Water Air Soil Pollut*, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 241, Copyright (1999), with permission from Springer Science and Business Media).
- Figure 6.12. Changes of Zn fractions in two Israeli soils during 336 days of incubation at the field capacity moisture regime (after Han and Banin, 1999. Reprinted from *Water Air Soil Pollut*, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 243, Copyright (1999), with permission from Springer Science and Business Media).
- Figure 6.13. Changes of Pb fractions in two Israeli soils during 336 days of incubation at the field capacity moisture regime (after Han and Banin, 1999. Reprinted from *Water Air Soil Pollut*, 114, Han F.X and Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. II: Incubation under field capacity conditions, p 242, Copyright (1999), with permission from Springer Science and Business Media).

- Figure 6.14. Changes of U_{is} of Cd, Cu and Zn in an Israeli sandy soil after 8 months of sludge application (T is the total metal content in the non-amended soil) (after Han et al., 2003a. Reprinted from *Adv Environ Res*, 8, Han F.X., Banin A., Kingery W.L., Triplett G.B., Zhou L.X., Zheng S.J., Ding W.X., New approach to studies of redistribution of heavy metals in soils, p 117, Copyright (2003), with permission from Elsevier).
- Figure 6.15. The average transfer fluxes of Cr and Cu among solid-phase components in Israeli soils. The soils received metal nitrates at the 3T level under the field capacity moisture regime (T: total metal content in non-amended soils).
- Figure 6.16. The average transfer fluxes of Ni and Zn among solid-phase components in Israeli soils. The soils received metal nitrates at the 3T level under the field capacity moisture regime (T: total metal content in non-amended soils).
- Figure 6.17. Relationships between the initial (the first day) fluxes of metals in the soluble plus exchangeable fraction and metal loading levels in Israeli soils. Soils received metal nitrates and were incubated in the saturated-paste (SP) and field capacity (FC) moisture regimes.
- Figure 6.18. Changes of the annual transfer fluxes (from one day to one year) among solid-phase components with metal loading levels in Israeli soils. Soils received metal nitrates and were incubated in the saturated-paste (SP) and field capacity (FC) moisture regimes.
- Figure 6.19. Distribution of Mn in solid-phase fractions in Israeli soils. Soils were incubated at the saturated paste regime (after Han and Banin, 1996. Reprinted from *Soil Sci Soc Am J*, 60, Han F.X., Banin A., Solid-phase manganese fractionation changes in saturated arid-zone soils: Pathways and kinetics, p 1075, Copyright (1996), with permission from Soil Sci Soc Am).
- Figure 6.20. Distribution of Mn in solid-phase fractions in Israeli soils. Soils were incubated at the field capacity regime.
- Figure 6.21. Initial and annual distribution and transformations of Mn in the major solid-phase fractions in two Israeli soils. Soils were incubated at the saturated-paste regime (after Han and Banin, 1996. Reprinted from *Soil Sci Soc Am J*, 60, Han F.X., Banin A., Solid-phase manganese fractionation changes in saturated arid-zone soils: Pathways and kinetics, p 1076, Copyright (1996), with permission from Soil Sci Soc Am).

- Figure 6.22. Distribution and transformations of Mn in the major solid-phase fractions in two Israeli soils. Soils were incubated at the field capacity regime.
- Figure 6.23. Changes of the redox parameter ($pe + pH$) in two Israeli arid soils during the saturated paste incubation (after Han and Banin, 1996. Reprinted from Soil Sci Soc Am J, 60, Han F.X., Banin A., Solid-phase manganese fractionation changes in saturated arid-zone soils: Pathways and kinetics, p 1076, Copyright (1996), with permission from Soil Sci Soc Am).
- Figure 6.24. Comparisons of Mn changes in the ERO (a) and the EXC and the CARB fractions (b) in an Israeli sandy soil during one year of saturation incubation and subsequent drying processes.
- Figure 6.25. Changes of R_{tf} of Mn fractions and U_{is} in Israeli sandy (a) and loessial (b) soils during one year of incubation at the saturated paste regime (after Han and Banin, 1996. Reprinted from Soil Sci Soc Am J, 60, Han F.X., Banin A., Solid-phase manganese fractionation changes in saturated arid-zone soils: Pathways and kinetics, pp 1077-1078, Copyright (1996), with permission from Soil Sci Soc Am).
- Figure 6.26. Kinetics of Mn transformations (described by the parabolic diffusion equation) among three main labile solid-phase fractions of Israeli arid soils. Soils were incubated at the saturated paste regime (after Han and Banin, 1996. Reprinted from Soil Sci Soc Am J, 60, Han F.X., Banin A., Solid-phase manganese fractionation changes in saturated arid-zone soils: Pathways and kinetics, p 1079, Copyright (1996), with permission from Soil Sci Soc Am).
- Figure 6.27. Distribution of Co in the solid-phase components in two Israeli arid soils incubated at the saturated paste regime (after Han et al., 2002b. Reprinted from J Environ Sci Health, Part A, 137, Han F.X., Banin A., Kingery W.L., Li Z.P., Pathways and kinetics of transformation of cobalt among solid-phase components in arid-zone soils, p 187, Copyright (2003), with permission from Taylor & Francis).
- Figure 6.28. Initial changes of Co concentrations in the main solid-phase fractions of two Israeli arid soils. Soils were incubated under the saturated paste regime (after Han et al., 2002b. Reprinted from J Environ Sci Health, Part A, 137, Han F.X., Banin A., Kingery W.L., Li Z.P., Pathways and kinetics of transformation of cobalt among solid-phase components in arid-zone soils, p 188, Copyright (2003), with permission from Taylor & Francis).

- Figure 6.29. Relationships between the Co and Mn contents in the CARB and the ERO fractions in Israeli soils during the saturated paste incubation.
- Figure 6.30. Kinetics of transformation of Co in the ERO fraction in two Israeli soils according to the two-constant rate model (a) and the simple Elovich model (b), respectively. Soils were incubated under the saturation paste regime (modified after Han et al., 2002b. Reprinted from *J Environ Sci Health, Part A*, 137, Han F.X., Banin A., Kingery W.L., Li Z.P., Pathways and kinetics of transformation of cobalt among solid-phase components in arid-zone soils, p 192, Copyright (2003), with permission from Taylor & Francis).
- Figure 7.1. Trace element concentrations in plants on California Donimo soil (pH 7.5) amended with metal sulfate-enriched sludge (Data from Mitchell et al., 1978).
- Figure 7.2. Ratios of DPTA-extractable trace elements in the rhizosphere over bulk Egyptian soils irrigated with sewage water for 10, 40, and 80 years (Data from El-Motaium and Badawy, 1999).
- Figure 7.3. Changes of Cd, Pb and Zn concentrations in soil solutions with pH from ten Colorado arid soils (data from Ma and Lindsay, 1990).
- Figure 7.4. Comparisons of decreases in NH_4NO_3 -extractable Zn, Ni and Cu in an Israeli loessial soil receiving metal nitrates and incubated under saturated paste, field capacity, and wetting/drying cycle moisture regimes (Han and Banin, 1997, 1999, and Han et al., 2001a).
- Figure 7.5. The AB-DTPA-extractable Cu and Zn in a semi-arid soil from Colorado after termination of five excessive biosolids applications. Open circle (fine, smectitic, mesic Aridic Argiustolls) and open triangle (fine, smectitic, mesic Aridic Paleustolls) represent control soil without biosolid application, while solid circle and solid triangle indicate biosolid applications (data extracted from Barbarick and Ippolito, 2003).
- Figure 7.6. Decreases in bioavailability (as NH_4NO_3 -extractable) of Cu, Ni and Zn in an Israeli loess soil receiving metal salts and incubated under field capacity regime for one year (data from Han and Banin, 1999).
- Figure 7.7. Changes of Cu and Zn concentrations in soil solution with DOC from South Australia (Data from Fotovat et al., 1997).
- Figure 7.8. Swiss chard Cd concentrations as affected by chloride levels in soils (modified from Weggler-Beaton et al., 2000).

- Figure 7.9. Effects of N rates on Cd concentrations in wheat shoots and grain (data from Mitchell et al., 2000).
- Figure 8.1. The accumulation of trace elements in two Californian soils. The soils received four-year sludge applications at the rate of 0, 22.5, 45, and 90 tons ha⁻¹ year⁻¹. Domino loam soil, fine-loamy, mixed, thermic Xerollic Calciorthid; Greenfield sandy loam soil, coarse-loamy, mixed, thermic Typic Haploxeralf (Data from Sposito et al., 1982).
- Figure 9.1. Measured and fitted global annual industrial age As production. (a) by mining; (b) released by coal burning; (c) released by petroleum burning; and (d) gross annual release into the environment (See explanations in the text) (after Han et al., 2003b. Reprinted from *Naturwissenschaften*, 90, Han F.X., Su Y., Monts D.L., Plodinec M.J., Banin A., Triplett G.B., Assessment of global industrial-age anthropogenic arsenic contamination, pp 396-397, Copyright (2003), with kind permission of Springer Science and Business Media).
- Figure 9.2. (a) The percentages of annual and cumulative As generated from the world-wide coal and petroleum industries over the global gross As production. (b) The percentages of annual and cumulative As generated from the world-wide petroleum industry over As generated from both coal and petroleum industries (after Han et al., 2003b. Reprinted from *Naturwissenschaften*, 90, Han F.X., Su Y., Monts D.L., Plodinec M.J., Banin A., Triplett G.B., Assessment of global industrial-age anthropogenic arsenic contamination, pp 396-398, Copyright (2003), with kind permission of Springer Science and Business Media).
- Figure 9.3. Measured and fitted global annual industrial age Hg mine production and Hg production from world coal industry (after Han et al., 2002a. Reprinted from *Naturwissenschaften*, 89, Han F.X., Banin A., Su Y., Monts D.L., Plodinec M.J., Kingery W.L., Triplett G.B., Industrial age anthropogenic inputs of heavy metals into the pedosphere, p 500, Copyright (2002), with kind permission of Springer Science and Business Media).

- Figure 9.4. Measured and fitted world annual metal production. The measured world annual metal production data were collected from U.S. Geological Survey-Minerals Information, 1997; Adriano, 1986; Woytinsky and Woytinsky, 1953. Chromium production was calculated from chromite production assuming an average of 27.0% Cr. The fitted Cd data before 1963, were calculated from the world annual Zn production (Cd mainly as a by-product) (after Han et al., 2002a. Reprinted from *Naturwissenschaften*, 89, Han F.X., Banin A., Su Y., Monts D.L., Plodinec M.J., Kingery W.L., Triplett G.B., Industrial age anthropogenic inputs of heavy metals into the pedosphere, p 499, Copyright (2002), with kind permission of Springer Science and Business Media).
- Figure 9.5. World cumulative metal productions since 1860 based on fitted world annual metal productions (after Han et al., 2002a. Reprinted from *Naturwissenschaften*, 89, Han F.X., Banin A., Su Y., Monts D.L., Plodinec M.J., Kingery W.L., Triplett G.B., Industrial age anthropogenic inputs of heavy metals into the pedosphere, p 499, Copyright (2002), with kind permission of Springer Science and Business Media).
- Figure 9.6. Global cumulative industrial age Hg mine production, Hg production from coal, and total global Hg production since 1860 based on estimated world annual Hg mine and coal production (inset: The percentage of Hg production from coal over the total global Hg production) (after Han et al., 2002a. Reprinted from *Naturwissenschaften*, 89, Han F.X., Banin A., Su Y., Monts D.L., Plodinec M.J., Kingery W.L., Triplett G.B., Industrial age anthropogenic inputs of heavy metals into the pedosphere, p 500, Copyright (2002), with kind permission of Springer Science and Business Media).
- Figure 9.7. Global cumulative industrial age As production from mining, As generated from coal and petroleum, and cumulative gross As production since 1850 (after Han et al., 2003b. Reprinted from *Naturwissenschaften*, 90, Han F.X., Su Y., Monts D.L., Plodinec M.J., Banin A., Triplett G.B., Assessment of global industrial-age anthropogenic arsenic contamination, p 397, Copyright (2003), with kind permission of Springer Science and Business Media).

List of Tables

- Table 1.1. Aridity zones by region ($\times 10^6$ ha) (after UNEP, 1993).
- Table 1.2. Land area by aridity zones in Africa ($\times 10^6$ ha) (after UNEP, 1993).
- Table 1.3. Global extent of arid and semi-arid soils (km^2) based on the Soil Taxonomy system) (after Monger et al., 2004).
- Table 2.1. Concentrations of trace elements in the earth's crust, rocks and world soils^a.
- Table 2.2. Concentrations of selected trace elements in representative arid and semi-arid soils of North China^a (from Liu, 1996).
- Table 2.3. Geometric means for selected trace elements in arid and semi-arid soils of the U.S.
- Table 2.4. Averages and ranges of concentrations of selected trace elements in arid and semi-arid soils of Israel^a.
- Table 3.1. The major ion composition of dispersed soil solutions and saturation extracts from selected soils in California, U.S.
- Table 3.2. Concentrations of trace elements in soil solutions of the California soils that received sludge applications.
- Table 3.3. Concentrations of trace elements in soil solutions extracted by saturated paste from two metal salt-spiked Israeli soils incubated at saturated regime.
- Table 3.4. Solution complexes of calcium and magnesium at 25°C^a .
- Table 3.5. Solution complexes of selected trace elements at 25°C^a .
- Table 3.6. The first hydrolysis constants of selected elements in soil chemistry^a.
- Table 3.7. Summary of major representative species of selected trace elements in arid soil solutions.
- Table 3.8. Possible soil solution solubility controls for selected trace elements^a.
- Table 3.9. Solubility product constants of common compounds and minerals of selected major and trace elements at 25°C^a .
- Table 4.1. Some selective sequential dissolution procedures employed to fractionate trace elements in soils^{a, b}.
- Table 4.2. Dissolution of some major elements extracted by NaOAc-HOAc solutions at various pHs after removing the exchangeable fraction^a.
- Table 4.3. Dissolution of some trace elements extracted by NaOAc-HOAc at various pHs after removing the exchangeable fraction (mg/Kg)^a.

- Table 4.4. Comparisons of two selective sequential dissolution procedures^a.
- Table 4.5. Extractants for trace elements bound to various solid-phase components in soils.
- Table 5.1. Percentages of trace metals bound in soil organic matter in two Israeli arid soils treated with metal nitrates^a.
- Table 5.2. Percentages of trace metals bound in the Mn and Fe oxides in two Israeli arid soils treated with metal nitrates^a.
- Table 5.3. Percentages of trace metals bound in carbonates in two Israeli arid soils treated with metal nitrates.
- Table 5.4. Distribution of trace elements among solid-phase components^a in 45 Israeli arid soils
- Table 5.5. Distribution of Zn among solid-phase components in some Chinese arid soils^a.
- Table 5.6. Some basic atomic properties of studied and related elements^a.
- Table 5.7. Solubility equilibria of selected minerals of related elements^a
- Table 6.1. The relevant soil properties.
- Table 6.2. The original metal concentration (HNO_3 -extractable) in native soils and levels of addition to the two Israeli arid soils incubated under the saturated paste regime condition.
- Table 6.3. Treatment levels of trace metals in the two Israeli arid soils incubated under the field capacity and wetting-drying cycle moisture regimes
- Table 6.4. Protocol for the selective sequential dissolution procedure.
- Table 6.5. Reduced partition index (I_R) of trace metals in arid-zone soils incubated under the saturated paste regime (after Han and Banin, 1997. Reprinted from Water Air Soil Pollut, 95, Han F.X., Banin A., Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils. I: Incubation under saturated conditions, p 411, Copyright (1997), with permission from Springer Science and Business Media).
- Table 6.6. Comparisons of the redistribution of metals in two Israeli soils at $3T^a$ treatment after one year of incubation under the saturated paste (SP), field capacity (FC), and wetting-drying cycle (Cycle) moisture regimes (% of the sum of fractions, as means of two replicates) (from Han et al., 2001a, with permission from Lippincott Williams & Wilkins).
- Table 6.7. The average residence time (hour) of heavy metals in Israeli soil solutions.

- Table 6.8. The kinetics of the initial changes of Mn concentration of three fractions in incubated Israeli soils under saturated regime (C: mg kg⁻¹; t: sec) (Han and Banin, 1996, with permission from Soil Sci Soc Am).
- Table 7.1. Concentrations of selected trace elements in plants grown on noncontaminated arid and semi-arid soils^a.
- Table 7.2. Extractants for bioavailable trace elements in arid and semi-arid soils.
- Table 7.3. Bioavailability of trace elements in arid and semi-arid soils.
- Table 7.4. Zinc sources and application rates for correcting zinc deficiency on crops^a.
- Table 7.5. Solubility product constants of common compounds/minerals of selected major and trace elements at 25 °C^a.
- Table 7.6. Concentrations of bioavailable trace elements in arid and semi-arid soils of selected countries^a.
- Table 7.7. Concentrations of bioavailable Zn (DTPA-extractable Zn), Cu (DTPA-extractable Cu), and B (water-soluble B) in arid and semi-arid soils of North China^a (from Liu, 1996).
- Table 7.8. Plants/animals and countries susceptible for micronutrient deficiency.
- Table 7.9. Diagnostic soil and plant tissue Cd criteria for a 25% and 50% yield decrements^a.
- Table 8.1. Agricultural irrigation water standards.
- Table 8.2. Trace element composition of sewage treatment plant effluents from the U.S., southern California, and Israel.
- Table 8.3. Concentrations of trace elements in Israeli arid soils (0-30 cm) irrigated with secondary sewage effluents (from Banin et al., 1981).
- Table 8.4. Trace element composition of sewage sludge from the U.S. and Israel.
- Table 8.5. Cumulative loading rates of trace elements for land applications of sewage sludge in selected countries^a.
- Table 8.6. Maximum annual loading rates of trace elements for land applications of sewage sludge in selected countries^a.
- Table 8.7. Maximum permissible trace element levels of sewage sludge for land applications in the U.S. and China^a.
- Table 8.8. Average concentrations of trace elements in phosphate rocks (PR)^a.
- Table 8.9. Concentrations of trace elements in vegetation and soils at Tharsis mined areas of semi-arid South West Spain^a.
- Table 8.10. Summary of the effects of Cd on microbial processes and enzyme activities in arid and semi-arid soils.

- Table 8.11. Chemicals and minerals for the remediation of trace element contaminated arid and semi-arid soils.
- Table 8.12. Plant species for the phytoremediation of trace element contaminated soils.
- Table 9.1. Global industrial age annual production of selected trace elements and heavy metals in 1880, 1900, 1950, 1990 and 2000 (million tons) (Data extracted from Han et al., 2002a, 2003b).
- Table 9.2. Global industrial age cumulative production of selected trace elements and heavy metals in 1880, 1900, 1950, 1990 and 2000 (million tons)^a.
- Table 9.3. Potential anthropogenic inputs in world soils and global metal burden per capita^a.
- Table 9.4. Trace element contents of the lithosphere and world soils (mg/kg) (Data from Han et al. 2002a, with kind permission of Springer Science and Business Media).
- Table 9.5. Arsenic contents of the lithosphere and world soils (Data from Han et al., 2003b, with kind permission of Springer Science and Business Media).