

CONTENTS

	ABSTRACT	5
	Acknowledgements	7
1	INTRODUCTION	8
2	CONCEPTUAL APPROACH	9
3	METHODS	12
	3.1 Study sites	12
	3.2 Description of the wind park	12
	3.3 Data collection	13
	3.3.1 Aerial surveys	13
	3.3.2 Ground survey and spatial distribution of sea ducks	13
	3.3.3 Benthic surveys	14
	3.3.4 On-Off experiment	15
	3.3.5 The exploitation experiment	16
	3.3.6 The decoy experiment	17
	3.4 Data analysis	18
	3.4.1 Assumptions of the statistical tests	18
	3.4.2 Comparing TK and RS	20
	3.4.3 Comparisons within TK	21
	3.4.4 Around the wind park	22
4	RESULTS AND DISCUSSION	23
	4.1 Seasonal abundance and bird diversity at TK	23
	4.2 Abundance of eiders: comparing TK and RS	24
	4.2.1 Aerial surveys	24
	4.2.2 Ground surveys	26
	4.3 Abundance of potential prey: comparing TK and RS	28
	4.3.1 Size of blue mussels	30
	4.3.2 Biomass of blue mussels	31
	4.3.3 Relationships between mussel biomass and abundance of eiders	35
	4.4 Abundance and spatial distribution of eiders within TK	35
	4.4.1 Abundance	37
	4.4.2 Spatial distribution	39
	4.5 Abundance of scoters: comparing TK and RS	41
	4.6 Investigations around the wind park	41
	4.6.1 The on-off experiment	41
	4.6.2 The exploitation experiment	46
	4.6.3 The decoys experiment	48
5	SUMMARY and CONCLUSIONS	49
	5.1 Comparison between TK and RS	50
	5.2 Comparison between years within TK	51

5.3	Around the wind park	52
5.4	Power of statistical tests	52
5.5	Comparison with other studies	54
5.6	Conclusions	54

6 ASPECTS NOT COVERED BY THIS STUDY 55

6.1	Other species and different phases of the annual cycle	55
6.2	Flock sizes	55
6.3	Flight behaviour and collision probabilities	56
6.4	Disturbance by the travelling ships	56
6.5	Size of the wind park	56
6.6	Mitigation and prediction of impacts applied to sea ducks	56

7 BIBLIOGRAPHY 58

ANNEX 61