

Contents

Contents	3
Abbreviations	6
Abstract	8
Zusammenfassung	10
I. Introduction	12
I.1. Organization of the immune system	
I.2. Major histocompatibility complex (MHC) molecules and their role in immunorecognition	
I.3. Interferon gamma (IFN γ) and regulation of MHC molecules	
I.4. Role of STAT1 and STAT3 signaling in biological systems	
I.5. Allorecognition of embryo and expression of MHC molecules in the course of embryonic development	
I.6. Embryonic stem (ES) cells	
I.7. Immunological properties of ES cells (ESCs)	
II. Aims	35
III. Results	36
III.1. Regulation of MHC class I molecules in ESCs	
III.1.1. Expression of MHC class I molecules on ESCs	
III.1.2. Expression of components of MHC class I processing machinery in ESCs and their differentiated derivatives	
III.1.3. Expression of IFN γ receptors in ESCs and their differentiated derivatives	
III.1.4. Expression and activity of STAT1 in ESCs	
III.1.5. Expression of regulatory components of STAT1 signaling pathway and their activation by IFN γ in murine ESCs	
III.1.6. The role of LIF in control of MHC class I expression in murine ESCs and its differentiated derivatives.	

III.1.7. The role of STAT3 in regulation of MHC class I molecules in undifferentiated ESCs

III.2. Epigenetic regulation of MHC class I molecules in murine ESCs

III.2.1. STAT3 regulates the expression of epigenetic modifier genes in ESCs

III.2.2. Dnamt knockout (KO) murine ES cell lines show enhanced MHC class I expression

III.2.3. Epigenetic modifiers 2,5-azacytidine and Trichostatin A increase the expression of MHC class I molecules on ESCs

III.3. Biological consequences of STAT3 KD in ESCs

III.3.1. Downregulation of STAT3 in murine ESCs reduces their lysis by syngenic NK cells

III.3.2. STAT3 partially inhibits the lysis of murine ESCs by CTLs

III.4. Regulation of MHC class I expression in ES cell-derived cardiomyocytes (ES-CMs)

III.4.1. STAT3 KD in ES-CMs does not induce the expression of MHC class I molecules on their surface

III.4.2. Effect of IFN γ on expression of components of MHC class I antigen processing machinery in ES-CMs

III.4.3. ES-CMs are partially lysed by CTLs only after induction of MHC class I molecules by IFN γ

IV. Discussion71

V. Materials and methods84

V.1. Materials

V.1.1. Cell lines

V.1.2. Mouse strains

V.1.3. Molecular biology reagents

V.1.4. Cell culture reagents

V.1.5. Radioactive material

- V.1.6. Common reagents
- V.1.7. Primers
- V.1.8. Antibodies used for immunoblotting
- V.1.9. Antibodies used for flow cytometry
- V.1.10. STAT3 gene short hairpin RNA (shRNA) sequence
- V.1.11. Glassware and Plastics
- V.1.12. Instruments and devices

V.2. Methods

- V.2.1. Culture of Undifferentiated ES/iPS cells and other cell lines
- V.2.2. RT-PCR and quantative RT-PCR
- V.2.3. STAT3 KD and its validation in ESCs
- V.2.4. Immunoblotting
- V.2.5. Flow cytometric analyses
- V.2.6. Luciferase reporter assay
- V.2.7. Teratoma formation by ES cells in mice
- V.2.8. Activation and isolation of ovalbumin-specific CTLs
- V.2.9. Activation and isolation of syngeneic NK cells
- V.2.10. ⁵¹Cr-release cytotoxicity assay
- V.2.11. Interferon-γ enzyme-linked immunosorbent assay
- V.2.12. Generation of ES cell-derived cardiomyocytes

VI. References	101
VII. Acknowledgements	114
VIII. Statement (Erklärung)	116