Table of Contents

| Sı | ımmary | Y | i |
|----|--------|--|-----|
| Zı | usamme | enfassung | ii |
| 1 | Intro | duction | 1 |
| | 1.1 Sk | in: structure and functions | 1 |
| | | itaneous wound healing and cutaneous fibrosis | |
| | 1.2.1 | Cutaneous wound healing | |
| | 1.2.3 | | |
| | 1.2.3 | 1.2 Tissue regenerative phase/Proliferative phase | 5 |
| | 1.2. | | 5 |
| | 1.2. | | |
| | | mal wound repair | |
| | 1.2.2 | Cutaneous fibrosis | |
| | | ole of macrophages and fibroblasts in tissue repair, fibrosis and extracellul | |
| | - | production | |
| | | Fβ-TGFβRII signaling | |
| | 1.4.1 | La colore de la co | |
| | 1.4.2 | The role of TGFβ1-TGFβRII mediated functions in macrophages | |
| | 1.5 Cr | e-LoxP system and LysMCre mouse strain | .15 |
| 3 | Pocul | ts | 17 |
| _ | | eneration of myeloid cell specific TGFβRII deficient mice | |
| | 3.1.1 | Myeloid cell specific deletion of TGFβRII | |
| | | Loss of TGFβ1–TGFβRII signaling in TGFβRII ^{fl/fl} LysMCre macrophages | |
| | 3.1.2 | | |
| | | fect of TGFBRII deletion in myeloid cells on thioglycollate induced peritoni | |
| | | itro macrophage chemotaxis | |
| | 3.2.1 | Abolishment of anti-inflammatory effect in TGFβRII ^{n/n} LysMCre mice during an | |
| | | matory model of thioglycollate induced peritonitis | |
| | 3.2.2 | Impact of TGFβRII deficiency on <i>in vitro</i> random migration of macrophages | |
| | | pact of TGFβRII deficiency in macrophages on the repair process | |
| | 3.3.1 | TGFβRII deficiency does not impair kinetics of wound closure but delays wour | |
| | | action | |
| | 3.3.2 | Impaired myofibroblast differentiation and impaired deposition of collagen at | |
| | | d site of TGFβRII ^{n/n} LysMCre mice | |
| | 3.3.3 | Similar expression of TGF-β1 in the wounds of control and TGFβRII ^{n/n} LysMCre | |
| | mice | | |
| | 3.3.4 | Loss of TGFβRII in macrophages does not impact neo-angiogenesis | |
| | 3.3.5 | Loss of TGFBRII in macrophages does not impact infiltration of macrophage in | |
| | the wo | ound site | 29 |

| 3.3.6 Loss of TGFBRII in macrophages has no influence on inflammatory cytokine |
|--|
| induction32 |
| 3.3.7 Loss of TGFβRII in macrophages does not impact pro-fibrotic, pro-inflammatory |
| and pro-angiogenic cytokine regulation in isolated wound macrophages32 |
| 3.4 Impact of loss of TGFβRII signaling in macrophages on bleomycin induced |
| fibrosis35 |
| 3.4.1 Abrogation of TGFβRII in macrophages results in reduced bleomycin induced |
| fibrosis |
| 3.4.2 Abrogation of TGF β RII in macrophages results in impaired collagen deposition in |
| bleomycin induced fibrosis37 |
| 3.4.3 TGFβRII deficiency does not affect the number of myofibroblast and endothelial |
| cells in bleomycin induced fibrosis38 |
| 3.4.4 Possible mechanisms involved in the reduced bleomycin induced fibrosis of |
| TGFβRII ⁿ /nLysMCre mice39 |
| 3.4.4.1 Reduced infiltration of myeloid cells in bleomycin induced fibrotic skin of |
| TGFβRII ^a /aLysMCre mice |
| 3.4.4.2 Altered dermal inflammatory cytokine profile in bleomycin induced fibrotic skin of TGFBRII ^{n/n} LysMCre mice41 |
| 3.4.4.3 Similar expression of pro-fibrotic factors in control and TGFβRII ^{#/‡} LysMCre fibrotic |
| dermis44 |
| 3.4.4.4 TNF-α expression in fibrotic skin direct correlates with TNF-α expression in TGFβRII |
| deficient macrophage isolated from fibrotic skin46 |
| 3.4.4.5 Possible mechanisms of reduced fibrosis and less extracellular matrix deposition in |
| TGFβRII ^{n/n} LysMCre mice46 |
| |
| 4 Discussion50 |
| 4.1 Macrophage specific deficiency of TGFβRII results in context-dependent effect |
| on cell migration under in vitro and in vivo condition |
| _ |
| |
| |
| 4.2.2 TGFβRII mediated signaling in macrophage does not impact cell recruitment in wound tissue |
| |
| 4.3 Impact of TGFβRII mediated signaling in macrophage on fibrosis54 |
| 4.3.1 TGFβRII mediated signaling in macrophage generates fibrotic response54 |
| 4.3.2 TGFβRII mediated signaling in macrophages results in reduced expression of pro- |
| inflammatory factors |
| 4.3.3 Loss of TGFβRII mediated signaling in macrophage results in reduced expression |
| of TNF-α |
| 4.4 Hypothetical model for fibrosis mediated by TGFβRII signaling in macrophages |
| 59 |

| 5 Mat | terials | and Methods | 60 |
|-------|----------|--|----|
| 5.1 | Materia | ıls | 60 |
| 5.1. | 1 Buf | fers used | 60 |
| 5.1. | .2 Che | micals and reagents | 61 |
| 5.1. | .3 Con | nmercial Kits | 62 |
| 5.1. | | ombinant proteins and enzymes | |
| 5.1. | | hnical equipments and softwares | |
| 5.1. | | er materials | |
| 5.1. | | nary antibodies | |
| 5.1. | | ondary antibodies | |
| 5.1. | | onucleotides | |
| | | s | |
| 5.2. | | v cytometry | |
| | | Single cell suspension | |
| 3 | 5.2.1.1. | | |
| | 5.2.1.1. | • | |
| | 5.2.1.1. | • | |
| 5. | .2.1.2 | FACS staining and analysis | 67 |
| 5.2. | | ation of genomic DNA | |
| 5.2. | 3 Poly | merase chain reaction (PCR) | 68 |
| 5.2. | 4 RN/ | extraction, reverse transcription PCR and quantitative real time PCR | 68 |
| 5.2. | | aration of epidermis and dermis | |
| 5.2. | .6 Cult | ure and maintenance of peritoneal macrophages | 69 |
| 5.2. | | tro random migration and analysis | |
| 5.2. | | thern blot | |
| | | DNA digestion | |
| 5. | | Fransfer of DNA on nitrocellulose membrane and pre-hybridization | |
| 5 | | Generation of TGFβRII specific probe and purification | |
| 5 | .2.8.4 | Radioactive probe labeling and purification | 72 |
| 5 | .2.8.5 | Hybridization and detection | 72 |
| 5.2. | 9 Wes | tern blot | 72 |
| | | Sample preparation and SDS-PAGE (Sodiumdodecyl-sulphate-polyacrylamide-goresis | |
| | • | Protein-transfer and detection | |
| 5.2. | | stology | |
| | .2.10.1 | Histochemistry, Sirius red and Giemsa staining | |
| | .2.10.2 | F4/80 staining on cryosections. | |
| | .2.10.3 | α-SMA/CD31 staining | |
| 5. | .2.10.4 | Gr-1 staining | 74 |
| 5.2. | 11 Mc | rphometric analysis | |
| 5. | .2.11.1 | Quantification of wound healing parameters and bleomycin induced fibrosis | 75 |
| 5. | .2.11.2 | Quantification of histochemical stainings | 75 |
| 5. | .2.11.3 | Quantification of immunohistochemical stainings | 75 |
| 5.2. | 12 In. | situ zymography | 75 |
| 5.2. | 13 In | vivo experiments | 76 |
| 5. | .2.13.1 | Mouse strains | 76 |
| | .2.13.2 | Genotyping | |
| | .2.13.3 | Thioglycollate-induced peritonitis and peritoneal cell harvest | |
| | .2.13.4 | Wounding | |
| 5. | .2.13.5 | Bleomycin induced fibrosis | 77 |

| | 5.2.14 | Statistical analysis | 78 |
|----|---------|----------------------|-----|
| 6 | Refere | nces | .79 |
| 7 | Abbrev | viations | .85 |
| 8 | Acknov | wledgments | .88 |
| 9 | Erklärı | ung | .89 |
| 10 | Curricu | ılum Vitae | .90 |