TABLE OF CONTENTS

	PAGE
Abbreviations	1
1 Introduction	4
1.1 Components, classification and cost of pain	4
1.1.1 Components of pain	4
1.1.2 Classification of pain	5
1.1.2.1 Adaptive pain	5
1.1.2.2 Maladaptive pain	6
1.1.3 Cost of pain	6
1.2 Nociception	8
1.3 Chronic pain	9
1.3.1 Neuropathic pain	9
1.3.2 Inflammatory pain	10
1.3.3 Mechanism for chronic pain	11
1.3.3.1 Peripheral sensitization	12
1.3.3.2 Central sensitization	13
1.3.3.3 Long term depression of inhibition	14
1.3.3.4 Neuro-glial interactions	
1.4 Neuronal plasticity and chronic pain	16
1.4.1 Biochemical profile changes	17
1.4.2 Functional plasticity	18
1.4.3 Structural plasticity	18
1.5 Dendritic spine morphogenesis	19
1.5.1 Molecular candidates in modulating spine morphogenesis	20
1.5.1.1 Ion channels	21
1.5.1.2 Kinases	22
1.5.1.3 Structural protein	22
1.5.1.4 Small G-protein family	2 3
1.5.1.5 Guanine-nucleotide exchange factor, GEF	2 5
1.6 Kal7	27
1.7 Multiple sclerosis	29
1.7.1 Types of MS	30
1.7.2 Cause of MS	
1.7.3 Pathological evidence and putative mechanisms	
1.7.4 Clinical manifestation and MS-related pain	
1.8 EAE, an animal model for MS	33
1.8.1 Pain in EAE	33
1.9 Aims of this doctoral thesis	35
2 Materials and Methods	
2.1 Materials	
2.1.1 Chemicals and reagents	37
2.1.2 Nucleic acid and protein size standards	39



2.1.3 Enzymes	39
2.1.4 Commercial kits	
2.1.5 Technical equipment4	40
2.1.6 Analytical software	40
2.1.7 Oligonucleotides4	41
2.1.8 List for antibodies	41
2.1.9 List for peptides	42
2.1.10 Buffers and solutions	42
2.1.11 Consumables	43
2.1.12 Animals	44
2.2 Methods	44
2.2.1 Bioinformatic analysis	44
2.2.1.1 Gene analysis of kalrn gene and RT-PCR strategy	44
2.2.2 Molecular biology analysis	45
2.2.2.1 Total RNA isolation	45
2.2.2.2 Reverse transcription (RT)	46
2.2.2.3 Polymerase chain reaction (PCR) amplification	46
2.2.2.4 Molecular cloning	
2.2.2.5 Transformation of chemical competent bacterial cells	47
2.2.2.6 Plasmid DNA preparation	47
2.2.2.7 Purification of DNA	48
2.2.2.8 Precipitation of nucleic acid	48
2.2.2.9 Restriction digestion	48
2.2.2.10 DNA-agarose gel electrophoresis	48
2.2.2.11 Quantitative Real-Time PCR	
2.2.3 Protein biochemistry	49
2.2.3.1 Protein isolation	49
2.2.3.2 SDS-Polyacrylamide gel electrophoresis (SDS-PAGE)	.50
2.2.3.3 Immunoblot	
2.2.4 Histology	.51
2.2.4.1 Tissue preparation for histology studies	.51
2.2.4.2 Chromogenic immunohistochemistry	.52
2.2.4.3 Immunoflourescence	
2.2.4.4 Immunocytochemistry	.53
2.2.5 Animal work	
2.2.5.1 Animal handling	.54
2.2.5.2 Spinal cord injection	
2.2.5.3 Induction of EAE via active immunization	
2.2.5.4 Monitoring of EAE development	.56
2.2.5.5 Nociceptive testing	
2.2.6 Electrophysiology	
2.2.6.1 Spinal cord slice preparation and patch-clamp recordings	
2.2.6.2 Afferent recordings in skin-nerve preparation	
2.2.7 Morphological analysis and imaging	

2.2.7.2 Confocal imaging and densitometry 60 2.2.8 Cell culture work 60 2.2.8.1 Primary spinal neuron preparation 60 2.2.9 Statistics 60 3 Results 60	1 1 2
2.2.8 Cell culture work	1 1 2
2.2.9 Statistics 6	2
3 Results6	
	2
	J
3.1 Analysis of Kaiirin expression in the mouse spinal cord6	3
3.1.1 Gene structure of Kalrn6	3
3.1.2 Basal expression of spinal Kalirin isoforms at mRNA and protein level6	4
3.1.2.1 Kaiirin mRNA transcripts expression in the mouse spinal cord6	4
3.1.2.2 Expression of particular kalirin isoforms as determined by 3' RACE6	6
3.1.2.3 Kal7 is expressed in neuronal population in the spinal cord in culture.6	7
3.1.2.4 Kalirin protein expression in the dorsal horn from the naive mouse	е
spinal cord6	8
3.1.3 Distribution of Kalirins in the naive mouse spinal cord6	9
3.2 Conditional knockout of Kalirin in the mouse lumbar spinal cord7	0
3.2.1 Strategy for Kalrn gene knockout7	0
3.2.2 Spinal delivery of AAV virus, validation and optimization7	0
3.2.2.1 Spinal injection of the AAV results in faithful and robust expression of	
gene of interest in neuronal cell population7	0
3.2.2.2 Viral expression distribution, quantification and optimization7	1
3.2.3 Expression of Cre recombinase within mouse spinal dorsal horn an	d
characterization7	3
3.2.3.1 Microinjection of Cre-expressing AAV virus led to selective gen	е
manipulations7	3
3.2.3.2 AAV-Cre successfully targeted most of the spinal dorsal horn neuron	S
in the nociceptive laminae7	'3
3.2.4 Spinal Knockout of kalirin via microinjection of AAV-Cre7	5
3.3 Effects of Kalirin on pain-related behavior7	6
3.3.1 Effects of Kalirin on the maintenance of formalin-induced nocifensive	e
behavior7	7
3.3.2 Effects of Kalirin on the maintenance of chronic inflammation-relate	d
pain behavior7	'9
3.3.2.1 Effects of spinal knockout of total Kalirins on the developmer	
/maintenance of mechanical hypersensitivity induced by intraplantar CF	
injection8	30
3.3.2.2 Deletion of total Kalirin did not affect CFA-induced therma	al
hypersensitivity8	30
3.3.2.3 Effects of spinal knockout of Kal7 on the development/maintenance	
mechanical hypersensitivity induced by intraplantar CFA injection	
3.3.2.4 Absence of Kal7 did not affect CFA-induced thermal hypersensitivity a	
compared to control group	
3.4 Effects of Kalirin on neuronal spine morphogenesis	

3.4.1 Spinal-neuron-specific deletion of Kal7 suppressed spine morphogenesis
resulting from peripheral inflammatory insult84
3.4.2 Conditional knockout of Kaiirin reduced the density of spine-like
structures in cultured spinal neurons85
3.5 Physiological and behavioral consequences of transient interruption
of Kal7 signaling87
3.5.1 Kal7-signaling-related electrophysiological properties in spino-PAG
neuron87
3.5.1.1 Kal7 C terminus (CT) peptide blocked LFS-induced spinal LTP in
spino-PAG lamina I neurons88
spino-PAG lamina I neurons88 3.5.1.2 Kal7 interfering peptide did not affect the basic membrane properties of
the spino-PAG lamina I neuron89
3.5.2 Physiological significance of transient block of Kal7-mediated signaling in
the spinal cord90
the spinal cord90 3.5.2.1 Trying to understand the role of Kal7 signaling during development of
CFA-induced mechanical hypersensitivity91
3.5.2.2 Kal7 transient blockade did not exert effect over CFA-induced thermal
hypersensitivity91
3.6 Effects of Rac1 over pain-related behavior93
3.6.1 Genetic manipulation of Rac1 expression within the spinal dorsal horn.93
3.6.2 Alterations in Rac1 expression and formalin-induced acute nocifensive
behavior94
3.6.2.1 Absence of Rac1 reduced second phase of formalin-induced
nocifensive behavior94
3.6.2.2 Overexpression of Rac1 exaggerated second phase of
formalin-induced nocifensive behavior94
3.6.3 Role of spinal Rac1 in the development of inflammatory pain behavior95
3.6.3.1 Absence of Rac1 reduced the CFA-induced mechanical
hypersensitivity96
3.6.3.2 Absence of Rac1 did not affect the CFA-induced thermal
hypersensitivity96
3.6.3.3 Overexpression of Rac1 altered the basal threshold of mechanical
perception but did not enhance the CFA-induced mechanical
hypersensitivity97
3.6.3.4 Overexpression of Rac1 has no effect on CFA-induced thermal
hypersensitivity98
3.7 Effect of Rac1 on spine morphogenesis in dorsal spinal neurons99
3.7.1 Spinal-deletion of Rac1 reduced spine formation in vivo99
3.7.2 Spinal-overexpression of Rac1 increased dendritic spine formation100
3.8 Immunization and behavioral analysis in different EAE mouse
models101
3.8.1 Induction of EAE via active immunization in C57BL6 and SJL mice102
3.8.2 Analysis of pain behavior and general locomotion103
3.8.2.1 Pain behavior in SJL-EAE and C57-EAE mice103

3.8.2.2 General locomotion in EAE animals106
3.9 Characterization of the sensory abnormalities in PNS in SJL-EAE
animals108
3.9.1 Electrophysiological analysis of peripheral nerve activity in SJL-EAE
mice108
3.9.2 Immunohistological analysis of the central terminals patterning of primary
afferents in the spinal cord of SJL-EAE mice110
3.10 Characterization of the spinal cord pathology in SJL-EAE
animals112
3.10.1 Verification of IB4-positive cells in the spinal dorsal horn
3.10.2 Analysis of the loss of NeuN-positive signals in the spinal cord113
3.10.3 Glia cells activation in spinal cord during EAE disease116
3.10.3.1 Immunohistochemical analysis of spinal glia activation in SJL-EAE
mice
3.10.3.2 Quantification of activated glia cells in the superficial dorsal horn in
SJL-EAE and C57-EAE mice117
4 Discussion122
4.1 Possible activators for Kal7 during chronic pain122
4.2 Are NR2R-containing NMDARs unstream of Kal7 in modulation of
4.2 Are NR2B-containing NMDARs upstream of Kal7 in modulation of
pain?127
pain?127 4.3 Kal7, Rac1 Dendritic spines and chronic pain131
pain?127 4.3 Kal7, Rac1 Dendritic spines and chronic pain131 4.4 Rac1 vs. RhoG132
pain?
pain? 127 4.3 Kal7, Rac1 Dendritic spines and chronic pain 131 4.4 Rac1 vs. RhoG 132 4.5 Pain in MS and EAE 134 4.6 Reduction of NeuN-positive cells in the spinal cord in EAE 137 4.7 Neuronal-glial interaction in EAE 138 4.8 Peripheral nerve alteration/ demyelination 141 4.9 Outlook 143 5 Summary 144 6 Reference 146 7 Publications 166
pain?
pain? 127 4.3 Kal7, Rac1 Dendritic spines and chronic pain 131 4.4 Rac1 vs. RhoG 132 4.5 Pain in MS and EAE 134 4.6 Reduction of NeuN-positive cells in the spinal cord in EAE 137 4.7 Neuronal-glial interaction in EAE 138 4.8 Peripheral nerve alteration/ demyelination 141 4.9 Outlook 143 5 Summary 144 6 Reference 146 7 Publications 166

.