

Vladimir Manea

Binding Energy of Strongly Deformed Radionuclides

Penning-Trap Mass Spectrometry and
Mean-Field Theoretical Studies

Doctoral Thesis accepted by
the Université Paris-Sud, France

Contents

1	Nuclear Observables	1
1.1	Trends of Binding Energies	1
1.2	Mass Filters	4
1.2.1	Estimators of the Single-Particle Energies	4
1.2.2	Estimators of the Odd-Even Staggering	11
1.2.3	Residual Effects	13
1.3	Complementary Nuclear Data	16
	References	19
2	Experimental Method and Data Analysis	21
2.1	Charged-Particle Traps	21
2.2	Penning Trap	22
2.2.1	Dynamics of a Trapped Ion	22
2.2.2	Driving the Ion's Motion	25
2.2.3	Detecting the Ion's Motion	27
2.3	From Cyclotron Frequency to Mass: Procedure, Precision, Systematic Errors	31
2.4	Production and Preparation of the Ion Ensemble	36
2.5	Complementary Applications of the MR-TOF MS	44
2.6	Experimental Results	47
2.6.1	Neutron-Rich Rubidium Isotopes	47
2.6.2	Neutron-Deficient Gold Isotopes	52
	References	57
3	Nuclear-Theory Concepts	61
3.1	Many-Body Calculations	61
3.2	The Hartree-Fock-Bogoliubov Approach	63
3.2.1	Hartree-Fock Field	63
3.2.2	Pairing Field	66
3.3	Competition Between Particle-Particle and Particle-Hole Correlations in Nuclei	69

3.4	Theoretical Analysis of the Measured Nuclear Data	71
3.4.1	Aim	71
3.4.2	HFODD Code	73
3.4.3	Tests of the Method	75
	References	80
4	Self-consistent Mean-Field Calculations	83
4.1	Neutron-Rich $A \approx 100$ Nuclei	83
4.2	Neutron-Deficient Gold-Thallium Nuclei	92
4.3	Odd-Even Staggering of Mercury Isotopes	99
4.4	Summary	103
	References	107
5	Conclusions and Outlook	109
	References	112
	Appendix A: Finite-Difference Operators	113
	Appendix B: Motion of a Charged Particle in a Penning Trap	117